



Evaluation of Palm Print Sign and Prayer Sign in Prediction of Difficult Laryngoscopy in Diabetic Patients

Amirreza Vakilian¹, Masoomeh Tabari^{1*}, Maryam Emadzadeh² and Ghasem Soltani¹

¹Department of Anesthesiology, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

²Clinical Research Development Unit, Ghaem Hospital, Mashhad University of Medical Sciences, Mashhad, Iran

*Corresponding author: Department of Anesthesiology, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran. Email: tabarim@mums.ac.ir

Received 2022 August 14; Revised 2022 December 18; Accepted 2022 December 26.

Abstract

Background: Laryngoscopy or endotracheal intubation in diabetic patients due to impaired mobility of the atlanto-occipital joint due to non-enzymatic glycosylation in connective tissue has always been a problem. A difficult laryngoscopy can disrupt the intubation process; therefore, it is valuable to predict difficult laryngoscopy using some anatomical criteria before general anesthesia.

Objectives: The present study discussed the diagnostic value of two tests, palm print sign and prayer sign, in the prediction of laryngoscopy difficulty.

Methods: Design: A diagnostic test and cross-sectional analytical design was used in this study. Setting: The study population included 200 patients with type 2 diabetes who were candidates for surgery under general anesthesia. Before surgery, the patients were examined regarding the airway status, Mallampati classification, head extension rate, thyromental distance, body mass index, upper lip biting test, and two palm print sign and prayer sign tests. All the diagnostic tests were compared to the Cormack test result for difficult airways regarding their sensitivity and specificity in difficult laryngoscopy.

Results: The highest sensitivity was related to the Mallampati test, prayer sign test, and palm print sign test (100%). Furthermore, the mouth-opening test had the highest specificity (100%). The highest accuracy was reported for Mallampati, palm print sign, and prayer sign tests (> 86%).

Conclusions: Among the tests studied to predict difficulty in laryngoscopy in diabetic patients, Mallampati and palm print sign tests have good sensitivity, specificity, and accuracy. Studies with a larger sample size are suggested to obtain more accurate results.

Keywords: Intubation, Diabetic Patients, General Anesthesia

1. Background

Intratracheal intubation is one of the most common methods of airway control. One of the risk factors that increase laryngoscopic and intubation stress is the difficulty of intubation in some patients (1). In general, inadequate ventilation, esophageal intubation, and difficult endotracheal intubation are the most common causes of severe respiratory consequences in the field of anesthesia (2). It is important to consider the anatomical signs and clinical factors associated with a difficult airway in patients with a potentially difficult airway (3). These indices include the Mallampati test, measurement of the thyromental distance (TMD), and the degree of head extension. Most of these indices do not have enough sensitivity and specificity (4). To predict easy or difficult airways, a series of common predictive tests are used that are easy, free, and non-invasive (2, 3). The Mallampati grading test is one of these methods used to predict the severity of intubation, which

has a sensitivity within the range of 42 - 81% (5, 6).

Diabetes is one of the most common endocrine diseases, which is especially important for the management of the airway for anesthesiologists. Diabetic individuals might be at greater risk for airway problems than the general population (1, 7, 8). Regarding the causes related to the underlying disease (i.e., diabetes), studies have shown that the presence of stiff joint syndrome or limited joint mobility, which is commonly observed in diabetic patients, can cause airway problems (6, 9). In some preliminary studies, the prevalence of airway problems in type 2 diabetic patients was reported to be 18.7%, compared to 2.5% in non-diabetic patients (10). In these studies, difficult laryngoscopy was defined as a three- or four-grade based on the Cormack-Lehane criteria (11).

Studies have also evaluated the relationship between difficult laryngoscopy and patients' ability to approach the palmar surfaces of their phalangeal joints, formerly

referred to as the “prayer sign” or the “palm print” sign. In palm print, the individual cannot extend the joints between the fourth and fifth fingers. This sign allows the evaluation of the laxity grade of the phalangeal joints, in this regard, the relationship between the rigidity of the palmar joint and involvement of cervical and laryngeal tissues in diabetic patients, which themselves lead to difficult airway and laryngoscopy (5). In some cases, even compared to the standard grading using the Mallampati grading, the use of the prayer sign is less sensitive. However, it has a similar specificity in determining the degree of difficult laryngoscopy in patients (12).

Due to the fact that intubation in diabetic patients is more difficult than in non-diabetic individuals, the accurate determination of the diagnostic value of prayer or palm print sign tests in the grading of difficult laryngoscopy in these patients is far more important than in non-diabetics. The problem is that there is still limited information about the sensitivity and specificity of these signs in the face of routine tests, such as the Mallampati test.

2. Objectives

This study aimed to assess two methods of palm print sign and prayer sign in diabetic patients and compare their degree of compatibility with the degree of laryngoscopy of the Cormack test.

3. Methods

This cross-sectional analytical study was performed in Ghaem and Imam Reza hospitals in Mashhad, Iran, in 2020 on patients with type 2 diabetes who were candidates for various types of surgery under general anesthesia. The inclusion criteria included the age of over 18 years and diabetic patients undergoing surgery under general anesthesia. The exclusion criteria included noticeable anatomical changes (e.g., congenital and traumatic after surgery) of the face, neck, or hands and related factors, such as rheumatoid arthritis, large thyroid, the mass of mouth and neck, pharyngeal abscess or other possible causes of difficult intubation, and body mass index (BMI) > 35 kg/m². The patients who could not cooperate for any reason, including physical disability and poor mental state, were also excluded.

Before surgery, patients with a history of diabetes who were candidates for surgery under general anesthesia were assessed for airway conditions. The tests used in this study included the Mallampati classification, TMD, mouth-opening test, and upper lip biting test (ULBT). In addition

to demographic characteristics, neck circumference and BMI were also determined and recorded.

For the measurement of TMD, the patient was in the supine position, with a 10 cm pillow under his/her head, and his/her neck was in the extension position. The distance between the thyroid incision and the internal incision of the chin protrusion was measured using a plastic ruler called TMD. This distance is divided into two groups; in one group, the TMD < 6.5 cm indicates the probability of difficult laryngoscopy, and in the other group, the TMD > 6.5 cm indicates the probability of easy laryngoscopy.

The ULBT is divided into three classes. Class I is the lower incisors that can bite above the upper lip line (probability of easy laryngoscopy). Class II is the lower incisors that can bite the upper lip. Class III is that lower teeth cannot bite the upper lip (probability of difficult laryngoscopy).

This study also examined the palm print sign and prayer sign. To assess the prayer sign, the patient was asked to place his/her hands together as in a prayer position; if there was a gap between the palms and the fingers, the test was positive and otherwise negative. To evaluate the palm print sign, the patient, while sitting, pressed the palm and fingers of one hand on a pad soaked in blue ink measured 25 × 16 cm for about 5 seconds, and after the hand was completely soaked in ink for 5 seconds, pressed them on the white paper which is located on a hard surface. Then, based on the recorded fingerprint on the paper, it was divided into four categories, which included class 0 (the effect of all interphalangeal is visible), class I (the fourth and fifth interphalangeal effect disappears), class II (the second to the fifth interphalangeal effect disappears), and class III (only fingerprints are visible). Classes II to III were considered positive.

On the day of surgery, after an intravenous line and complete monitoring, induction of anesthesia with midazolam, fentanyl, propofol, or thiopental sodium was performed, and an atracurium muscle relaxant was used to facilitate intubation. Laryngoscopy was performed with a Macintosh metal laryngoscope blade by an anesthesia assistant who had more than 2 years of intubation experience. Difficult laryngoscopy was defined in the present study using the Cormack test as grade III and higher. All diagnostic tests were compared to Cormack test results for difficult airways in terms of their sensitivity and specificity in difficult laryngoscopy.

3.1. Sample Size Calculation and Statistical Analysis

The sample size was calculated based on the data obtained from Hashim and Thomas (13), which reported a palm sign sensitivity of 77%. Considering alpha = 0.05, beta

= 0.2, and $d = 0.15$ p, the initial sample size was calculated at 57 patients using the following formula:

$$n = (z_{1-\alpha/2})^2 p(1-p)/d^2$$

Then, it was divided by 0.3 (i.e., the estimated prevalence of diabetic patients who needed general anesthesia in the studied center), and the final sample size was calculated at 190 patients.

The data were analyzed using SPSS software (version 23). To determine the sensitivity, specificity, positive and negative predictive value, and diagnostic accuracy of the palm print sign and prayer sign in comparison to the usual tests, the cross-tabulation method was used to calculate these parameters. Diagnostic parameters were calculated based on the percentage.

3.2. Ethical Considerations

Ethical approval for this study (IR.MUMS.AC.REC.1398.674) was granted by the Ethics Committee of Mashhad University of Medical Sciences, Mashhad, Iran, on January 1, 2020.

4. Results

A total of 200 patients were included in this study. The mean age of the patients was 59.30 ± 11.90 years. Regarding gender distribution, 137 (68.5%) and 63 (31.5%) cases were male and female, respectively. The mean neck circumference was 43.13 ± 3.63 cm. Table 1 shows patients' mean weight, height, HbA1c, and blood sugar.

Table 1. Demographic, Clinical Information, and Laboratory Findings of Patients with Type 2 Diabetes in the Current Study

| Variables | Mean \pm SD |
|---------------------|--------------------|
| Age (y) | 59.30 \pm 11.90 |
| Weight (kg) | 93.09 \pm 16.17 |
| Height (cm) | 169.57 \pm 17.42 |
| Blood Sugar (mg/dL) | 144.17 \pm 32.32 |
| HbA1c | 5.92 \pm 0.78 |

Abbreviation: SD, standard deviation.

In terms of BMI, 21 (10.5%), 71 (35.5%), and 108 (54%) patients were normal (BMI: 18.5 - 24.9 kg/m²), overweight (BMI: 25 - 29.9 kg/m²), and obese (BMI: 30 - 34.9 kg/m²), respectively. Furthermore, the mean values of blood sugar and HbA1c were 144.17 ± 32.32 mg/dL and 5.92 ± 0.78 , respectively. In terms of the tests used to determine laryngoscopic difficulty, firstly, regarding grading based on the Cormack test, grades I, II, and III/IV were determined in 51 (25.5%), 73 (36.5%), and 76 (38%) patients, respectively. Table 2 shows the frequency of clinical indicators of the airways. In comparison to the results of the Cormack scoring

system, the diagnostic accuracy of each test in the present study was determined (Table 3). The highest sensitivity was related to the Mallampati test, prayer sign test, and palm print sign test (100%). The mouth-opening test had the highest specificity (100%).

Table 2. Frequency of Clinical Indicators of the Airways

| Variables | No. (%) |
|------------------------------|------------|
| Cormack test | |
| Grade 1 | 51 (25.5) |
| Grade 2 | 73 (36.5) |
| Grade 3 | 46 (23) |
| Grade 4 | 30 (15) |
| Mallampati test | |
| Grade 1 | 42 (21) |
| Grade 2 | 112 (56) |
| Grade 3 | 36 (18) |
| Grade 4 | 10 (5) |
| Thyromental distance | |
| < 6.5 cm | 97 (48.5) |
| > 6.5 cm | 103 (51.5) |
| Mouth opening test | |
| < 3 cm | 26 (13) |
| > 3 cm | 174 (87) |
| Upper lip biting test | |
| Grade 1 | 48 (24) |
| Grade 2 | 107 (53.5) |
| Grade 3 | 45 (22.5) |
| Palm print sign test | |
| Class 0 | 27 (13.5) |
| Class 1 | 42 (21) |
| Class 2 | 96 (48) |
| Class 3 | 35 (17.5) |
| Prayer sign test | |
| Positive | 146 (73) |
| Negative | 54 (27) |

5. Discussion

Laryngoscopy or endotracheal intubation in diabetic patients due to the impaired mobility of the atlanto-occipital joint due to non-enzymatic glycosylation in connective tissue has always been a problem. In this regard, the use of some anatomical criteria even before general

Table 3. Diagnostic Values of Tests ^a

| Tests | Cormack Test Result | | Sensitivity (%) | Specificity (%) | PPV (%) | NPV (%) | Accuracy (%) |
|------------------------------|---------------------|-----|-----------------|-----------------|---------|---------|--------------|
| | a | b | | | | | |
| Mallampati test | | | | | | | |
| Class 3, 4 | 3 | 119 | 100 | 96.2 | 97.5 | 100 | 98.5 |
| Class 1, 2 | 78 | 0 | | | | | |
| Thyromental distance | | | | | | | |
| < 6.5 | 3 | 94 | 79 | 96.2 | 96.9 | 75.7 | 86 |
| > 6.5 | 78 | 25 | | | | | |
| Upper lip biting test | | | | | | | |
| Grade 3 | 39 | 6 | 32.8 | 92.6 | 86.7 | 48.4 | 57 |
| Grades 1, 2 | 80 | 75 | | | | | |
| Mouth opening test | | | | | | | |
| < 3 | 81 | 93 | 21.8 | 100 | 100 | 46.5 | 53.5 |
| > 3 | 0 | 26 | | | | | |
| Palm print sign test | | | | | | | |
| Grades 2, 3 | 12 | 119 | 100 | 85.2 | 90.8 | 100 | 94 |
| Grades 0, 1 | 69 | 0 | | | | | |
| Prayer sign test | | | | | | | |
| Positive | 27 | 119 | 100 | 66.7 | 81.5 | 100 | 86.5 |
| Negative | 54 | 0 | | | | | |

Abbreviations: PPV, positive predictive value; NPV, negative predictive value.

^a a: Grade 3, 4 (difficult laryngoscopy); b: Grade 1, 2 (easy laryngoscopy).

anesthesia of the patient can be difficult to predict laryngoscopy and therefore disrupt the intubation process.

However, even some of the used methods, especially on their own, have not been able to predict such a feature in these patients and have not had enough sensitivity, specificity, and diagnostic accuracy for such a prediction. In such cases, the use of the Cormack scoring system as a standard method in determining the accuracy of laryngoscopy has always been helpful. The Cormack-Lehane four-part classification system, introduced in 1984 to describe landscape during direct laryngoscopy, is widely used in clinical trials to manage patients with a difficult airway.

According to the Cormack-Lehane classification, the glottal entrance view during direct laryngoscopy is divided into four subgroups. However, in cases with high grades of this system, the possibility of performing a laryngoscopy is difficult. Even in grade II of this system, difficulty in performing laryngoscopy is reported; therefore, the use of new grading systems for such a purpose is recommended. On the other hand, it is not possible to predict the difficulty of laryngoscopy before performing it in this method, and it is not possible to predict the performance of such a procedure until the laryngoscope enters

and observes or does not observe the glottis. Therefore, researchers have been looking for preferably anatomical indices that can predict the difficulty of this procedure even before the patient enters the operating room and undergoes anesthesia. In this regard, various tests, such as the Mallampati test, TMD (> 6.5 cm with probability of easy laryngoscopy), ULBT, mouth opening (> 3 cm with probability of easy laryngoscopy), and two palm print sign and prayer sign test are presented, each with its own diagnostic accuracy in predicting the ease of laryngoscopy.

What was discussed in the present study was the value and diagnostic value of these tests, especially the two recent tests in comparison to the Cormack grading system in predicting the difficulty of laryngoscopy. The results of the present study provided significant points in the use of these tests; however, it should be borne in mind that the test will be valuable if it has an acceptable sensitivity and specificity at the same time.

In this study, among the mentioned tests, two tests of biting the upper lip and opening the mouth due to very low sensitivity were not valuable in evaluating and predicting laryngoscopic difficulty. Among the other four tests, the highest sensitivity was related to the Mallampati test,

palm print sign test, and prayer sign test. Nevertheless, based on the available evidence, the Mallampati test also had some limitations. For example, how the patient is placed in bed and even the patient's sighing and breathing pattern can affect the diagnostic sensitivity and accuracy of this test. Considering that the palm print sign and prayer sign tests are not affected at all by the other mentioned factors, they can be the most practical and reliable tests in such evaluation. In different studies, the sensitivity and specificity of the palm print sign and prayer sign tests have been evaluated, and almost all studies emphasized the effectiveness of these two tests.

In the study of Hashim and Thomas (13), the sensitivity, specificity, positive predictive value, negative predictive value, and accuracy of the palm print test were 76.9%, 89.4%, 71.4%, 91.3%, and 86.7%, respectively, very similar to the results of the present study. In the study of Vani et al. (14), the palm print sign test has the highest sensitivity (equal to 75%). In the study of Nadal et al. (15), similar to the current study, the palm print sign test had the highest sensitivity of 100%. In the study by Reissell et al. (16), there was a linear correlation between laryngoscopic difficulty and the palm print sign test classification. In the study of Baig and Khan (12), contrary to the findings of the present study, the sensitivity of the prayer sign test and the Mallampati test was 29.6% and 79.3%, respectively. However, both tests had low specificity in his study.

The summary of the studies indicated the value of the palm print sign test in predicting the difficulty of laryngoscopy. Regarding the difficulty of laryngoscopy in diabetic patients, the frequency mentioned in different studies has been very different. In the present study, this frequency is equal to 59.5% and within the range mentioned in other studies.

In the study by Vani et al. (14), the incidence of laryngoscopy was 16%, which is much lower than in the present study. In the study by Reissell et al. (16), the overall incidence of laryngoscopy was 31%, which is lower than in the current study. In the study by Erden et al. (10), the incidence of laryngoscopy was 18.75% and 2.5% in diabetic and non-diabetic groups, respectively. In addition, in the study by Baig and Khan (12), a total of 35% of patients had difficult laryngoscopy. It seems that the higher degree of difficulty in laryngoscopy, in addition to underlying disorders in the patient (e.g., diabetes and laryngopharyngeal duct disorders), depends mainly on the experience of the relevant anesthesiologist. According to the limitations of the present study, further studies with more observations are needed for these purposes.

5.1. Conclusions

To conclude, among the tests studied to predict difficulty in laryngoscopy in diabetic patients, Mallampati and palm print sign tests have good sensitivity, specificity, and accuracy. Studies with a larger sample size are suggested to obtain more accurate results.

Acknowledgments

This article is the result of a dissertation in anesthesiology at Mashhad University of Medical Sciences. The authors would like to express their gratitude to the Clinical Research Development Unit of Ghaem hospital for cooperation in data analysis.

Footnotes

Authors' Contribution: Amirreza Vakilian: Design of the project, supervision of the study, writing the manuscript, and performing the study; Masoomeh Tabari: Design of the project, supervision of the study, editing the manuscript, and performing the study; Maryam Emadzadeh: Analyzing the data and editing the manuscript; Ghasem Soltani: Writing the manuscript and performing the study.

Conflict of Interests: Funding or research support: This article was extracted from Vakilian's dissertation, and no fund from any company was received. Employment: All the authors are faculty members of Mashhad University of Medical Sciences except Vakilian, who was an anesthesiology resident, and this article is extracted from his dissertation. Personal financial interests: This article was extracted from Vakilian's dissertation, and no financial support was received. Stocks or shares in companies: The authors did not have a contractual relationship with any company. Consultation fees: The authors did not receive consulting fees from any organization. Patents: The authors had no patent within the last 5 years. Personal or professional relations with organizations and individuals (e.g., parents and children, wife and husband, and family relationships): The authors had no personal or professional relationships with organizations and individuals. Unpaid membership in a governmental or non-governmental organization: None of the authors has unpaid membership in a governmental or non-governmental organization. Are you one of the editorial board members or a reviewer of this journal? The authors declared that one of the authors (Ghasem Soltani, reviewer) is one of the reviewers. The journal confirmed that the mentioned author with conflicts of interest (CoI) was completely excluded from all review processes. The authors also introduced this author with CoI as an opposed reviewer during the submission.

Ethical Approval: Ethical approval for this study (IR.MUMS.AC.REC.1398.674) was granted by the Ethics Committee of Mashhad University of Medical Sciences, Mashhad, Iran, on January 1, 2020.

Funding/Support: This article is the result of a dissertation in anesthesiology at Mashhad University of Medical Sciences. No grant was used for this study.

Informed Consent: All patients completed an informed consent form.

References

- Shiga T, Wajima Z, Inoue T, Sakamoto A. Predicting difficult intubation in apparently normal patients: a meta-analysis of bedside screening test performance. *Anesthesiology*. 2005;**103**(2):429-37. [PubMed ID: 16052126]. <https://doi.org/10.1097/00000542-200508000-00027>.
- Newacheck PW, Halfon N. Prevalence and impact of disabling chronic conditions in childhood. *Am J Public Health*. 1998;**88**(4):610-7. [PubMed ID: 9551003]. [PubMed Central ID: PMC1508436]. <https://doi.org/10.2105/ajph.88.4.610>.
- Bell RA, Mayer-Davis EJ, Beyer JW, D'Agostino Jr RB, Lawrence JM, Linder B, et al. Diabetes in non-Hispanic white youth: prevalence, incidence, and clinical characteristics: the SEARCH for Diabetes in Youth Study. *Diabetes Care*. 2009;**32** Suppl 2(S102-11). [PubMed ID: 19246575]. [PubMed Central ID: PMC2647694]. <https://doi.org/10.2337/dc09-S202>.
- Hogan K, Rusy D, Springman SR. Difficult laryngoscopy and diabetes mellitus. *Anesth Analg*. 1988;**67**(12):1162-5. [PubMed ID: 3057934].
- Nolan CJ, Damm P, Prentki M. Type 2 diabetes across generations: from pathophysiology to prevention and management. *Lancet*. 2011;**378**(9786):169-81. [PubMed ID: 21705072]. [https://doi.org/10.1016/S0140-6736\(11\)60614-4](https://doi.org/10.1016/S0140-6736(11)60614-4).
- Warner ME, Contreras MG, Warner MA, Schroeder DR, Munn SR, Maxson PM. Diabetes mellitus and difficult laryngoscopy in renal and pancreatic transplant patients. *Anesth Analg*. 1998;**86**(3):516-9. [PubMed ID: 9495404]. <https://doi.org/10.1097/00000539-199803000-00012>.
- Apfelbaum JL, Hagberg CA, Caplan RA, Blitt CD, Connis RT, Nickinovich DG, et al. Practice guidelines for management of the difficult airway: an updated report by the American Society of Anesthesiologists Task Force on Management of the Difficult Airway. *Anesthesiology*. 2013;**118**(2):251-70. [PubMed ID: 23364566]. <https://doi.org/10.1097/ALN.0b013e3182773b2>.
- Wild S, Roglic G, Green A, Sicree R, King H. Global prevalence of diabetes: estimates for the year 2000 and projections for 2030. *Diabetes Care*. 2004;**27**(5):1047-53. [PubMed ID: 15111519]. <https://doi.org/10.2337/diacare.27.5.1047>.
- Kim RP, Edelman SV, Kim DD. Musculoskeletal Complications of Diabetes Mellitus. *Clin Diabetes*. 2001;**19**(3):132-5. <https://doi.org/10.2337/diaclin.19.3.132>.
- Erden V, Basaranoglu G, Delatioglu H, Hamzaoglu NS. Relationship of difficult laryngoscopy to long-term non-insulin-dependent diabetes and hand abnormality detected using the 'prayer sign'. *Br J Anaesth*. 2003;**91**(1):159-60. [PubMed ID: 12821580]. <https://doi.org/10.1093/bja/aeg583>.
- Cormack RS, Lehane J. Difficult tracheal intubation in obstetrics. *Anaesthesia*. 1984;**39**(11):1105-11. [PubMed ID: 6507827].
- Baig MM, Khan FH. To compare the accuracy of prayer's sign and Mallampatti test in predicting difficult intubation in diabetic patients. *J Pak Med Assoc*. 2014;**64**(8):879-83. [PubMed ID: 25252511].
- Hashim K, Thomas M. Sensitivity of palm print sign in prediction of difficult laryngoscopy in diabetes: A comparison with other airway indices. *Indian J Anaesth*. 2014;**58**(3):298-302. [PubMed ID: 25024473]. [PubMed Central ID: PMC4090996]. <https://doi.org/10.4103/0019-5049.135042>.
- Vani V, Kamath SK, Naik LD. The palm print as a sensitive predictor of difficult laryngoscopy in diabetics: a comparison with other airway evaluation indices. *J Postgrad Med*. 2000;**46**(2):75-9. [PubMed ID: 11013469].
- Nadal JL, Fernandez BG, Escobar IC, Black M, Rosenblatt WH. The palm print as a sensitive predictor of difficult laryngoscopy in diabetics. *Acta Anaesthesiol Scand*. 1998;**42**(2):199-203. [PubMed ID: 9509203]. <https://doi.org/10.1111/j.1399-6576.1998.tb05109.x>.
- Reissell E, Orko R, Maunukela EL, Lindgren L. Predictability of difficult laryngoscopy in patients with long-term diabetes mellitus. *Anaesthesia*. 1990;**45**(12):1024-7. [PubMed ID: 1980577]. <https://doi.org/10.1111/j.1365-2044.1990.tb14879.x>.