Published online 2022 November 20.

### **Research Article**

# Intravenous Versus Wetting Solution Magnesium Sulphate to Counteract Epinephrine Cardiac Adverse Events in Abdominal Liposuction: A Randomized Controlled Trial

Motaz Amr Abu Sabaa 💿<sup>1,\*</sup>, Amr Arafa Elbadry 💿<sup>1</sup>, Safwat Hegazy<sup>2</sup> and Dina Ahmed El Malla<sup>1</sup>

<sup>1</sup>Surgical Intensive Care and Pain Medicine, Faculty of Medicine, Tanta University, Tanta, Egypt
<sup>2</sup>Faculty of Medicine, Kafr Elsheikh University, Kafr Elsheikh, Egypt

Corresponding author: Lecturer of Anesthesiology, Surgical Intensive Care and Pain Medicine, Faculty of Medicine, Tanta University, Tanta, Egypt. Email: motazabusabaa@gmail.com

Received 2022 July 12; Accepted 2022 November 02.

# Abstract

**Background:** The increased frequency of liposuction has resulted in more complications being reported. Adding epinephrine to the wetting solution could induce some cardiac adverse effects, some of which may be fatal. For instance, magnesium sulfate (MgSO<sub>4</sub>) is known for its cardioprotective effects.

**Objectives:** This study aimed to evaluate the efficacy of intravenous (IV) versus wetting solution containing MgSO<sub>4</sub> in decreasing such cardiac adverse events during abdominal liposuction.

**Methods:** This randomized controlled study included 129 adult cases undergoing abdominal liposuction under general anesthesia. The participants were divided into 3 groups: Group I (control group) was only subjected to the injection of the wetting fluid (1 mL 1/1000 epinephrine added to every 1000 mL of normal saline), group II was subjected to IV MgSO<sub>4</sub> (40 mg/kg over 1 minute) at the same time of installing the wetting solution, and group III was subjected to MgSO<sub>4</sub> (40 mg/kg) added to the wetting solution. **Results:** Intraoperative isoflurane consumption, intraoperative heart rate (HR), mean arterial pressure (MAP), and postoperative pain scores were significantly lower in the MgSO<sub>4</sub> groups (groups II and III) than group I. Cardiac adverse events (sinus tachycardia and premature ventricular contractions (PVCs)) were also significantly less frequent in groups II and III compared to group I. **Conclusions:** Adding MgSO<sub>4</sub>, either through IV or subcutaneous routes, is associated with lower intraoperative HR, MAP, and postoperative pain scores and a remarkable decrease in epinephrine-induced cardiac adverse events during liposuction.

Keywords: MgSO<sub>4</sub>, Epinephrine, Liposuction, Cardiac Adverse Events

# 1. Background

Liposuction is one of the most commonly performed surgical procedures worldwide. According to the American Society of Plastic Surgeons, this surgery is ranked second, following breast augmentation surgery (1, 2). With the introduction of wetting solutions, liposuction has become much safer. Before liposuction, a wetting solution is injected into the subcutaneous tissue, decreasing the bleeding caused by the early "dry" liposuction procedure (3). Despite its increased safety, the increased rates of the operations have enhanced relevant complications, some of which are rare but serious (2).

Epinephrine is an essential component of liposuctionwetting solutions. It is often used in 1/1000 mL because of its vasoconstrictor effect, which decreases intraoperative bleeding. Moreover, it prolongs the duration of local anesthesia if used with the injectate (4, 5). However, over time, large volumes of wetting solution can be installed with procedure improvement, exposing patients to doses > 5 - 10 mg during large-volume cases (4, 6).

Several studies have attributed the mortality of liposuction procedures to cardiac dysrhythmias, myocardial infarction, fluid overload, pulmonary edema, and fatal asystole. However, no clear etiology has been determined (7-9).

Although epinephrine toxicity could be incriminated in most of such cardiac side effects, this is difficult to prove on the clinical aspect for many reasons. First, epinephrine has a very short half-life (2 minutes). Second, post-mortem metabolism can also occur in humans. Moreover, the patient usually receives exogenous epinephrine during resuscitation attempts (9).

Magnesium has a documented beneficial effect on the cardiovascular system and can stabilize membrane po-

Copyright © 2022, Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (http://creativecommons.org/licenses/by-nc/4.0/) which permits copy and redistribute the material just in noncommercial usages, provided the original work is properly cited.

tential, activate antioxidant pathways, and enhance mitochondrial function (10). Moreover, its deficiency has been incriminated in multiple cardiovascular pathologies, including cardiomyopathy, atherosclerosis, hypertension, and dyslipidemia (11, 12).

# 2. Objectives

We hypothesized that adding MgSO<sub>4</sub> during the liposuction procedure could counteract the cardiac epinephrine-related side effects. Few studies have addressed this issue; hence, we conducted this study to evaluate the effectiveness of intravenous (IV) versus wetting solution MgSO<sub>4</sub> in counteracting epinephrine-induced cardiac adverse events in patients undergoing abdominal liposuction procedures.

#### 3. Methods

This randomized controlled single-blind study included 129 patients aged 20 - 60 years, with body mass index (BMI) of 25 - 35 kg/m<sup>2</sup>, and American Society of Anesthesiologists (ASA) class I or II, who underwent abdominal liposuction. The study was conducted in the Plastic Surgery Department, Tanta University Hospitals, Egypt, from March 2021 to March 2022 after gaining approval from the Ethics Committee of the Tanta University (code: 35029/11/21) and then registration in the Pan African Clinical Trial Registry (code: PACTR202105734020705). Informed written consent was obtained from all participants' guardians.

Exclusion criteria were hepatic or renal insufficiency, known allergy to the study medications, and non-abdominal liposuction.

#### 3.1. Randomization and Blindness

The participants were randomly assigned into 3 equal groups. Randomization was done using the sealed envelope method by a nurse not participating in the study. The research groups were as follows: Group I (control group) was only subjected to the wetting solution along with wetting fluid with 1 mL 1/1000 epinephrine added to 1000 mL of normal saline with no further adjuvants; group II was subjected to the IV administration of MgSO<sub>4</sub> (40 mg/kg) for 1 minute at the same time of wetting solution injection, and group III was subjected to MgSO<sub>4</sub> (40 mg/kg) added to the wetting solution (the total dose was divided equally according to the expected number of saline bottles injected).

Before surgery, all cases were subjected to full history recording, clinical examination, and routine laboratory investigations. All operations were performed under general anesthesia (GA) using a wide bone IV cannula after establishing routine hemodynamic monitoring (pulse oximetry, blood pressure, ECG, capnogram, and temperature probe). Moreover, GA was induced by fentanyl 1  $\mu$ g/kg, propofol 2 mg/kg, and atracurium 0.5 mg/kg. The maintenance of anesthesia was achieved by isoflurane (1-1.5%) in an oxygen and air mixture, along with cisatracurium (0.03 mg/kg) (MV > ETCO2 35 - 90).

Following anesthesia and skin sterilization, wetting fluid was injected. For all the included cases, 1 mL 1/1000 epinephrine was added to 1000 mL of normal saline. The tumescent technique was used for all cases, as the solution was administered until the targeted tissues (fatty deposits) were sufficiently filled or engorged. The dose of MgSO<sub>4</sub> injected was set based on the previous study (13).

Mean arterial blood pressure (MAP) and heart rate (HR) were recorded at the baseline, induction, installation of wetting solution, and then every other 15 minutes by the end of the procedure. Intraoperative cardiac adverse effects, including sinus tachycardia, premature ventricular contractions, and cardiac arrest, were recorded. Total isoflurane consumption was measured using MVCGE after surgery, and all patients received IV paracetamol (1 mg/8 hours) in addition to ketorolac (30 mg per 12 hours). Postoperative pain was assessed by the visual analogue scale (VAS) at PACU and then 1, 3, 6, 12, and 24 hours after surgery. If the patient reported VAS > 3, IV morphine (2 mg) was administered.

Our primary outcome was the mean HR following the installation of the wetting solution during liposuction. In contrast, the secondary outcomes were intraoperative isoflurane consumption, postoperative pain, morphine consumption, and the incidence of cardiac side effects.

Sinus tachyarrhythmia is a heart rhythm with a ventricular rate of > 100 beats/min. Tachyarrhythmias are broadly categorized as narrow complex tachycardia (NCT; < 120 ms) or wide complex tachycardia ( $\geq$  120 ms) (14). PVCS is a premature ventricular contraction, expected when the heartbeat is initiated by Purkinje fibers in the ventricles rather than by the sinoatrial node (15).

#### 3.2. Sample Size Estimation

The sample size was calculated using G\*Power 3.1.9.2 (Universitat Kiel, Germany). A pilot study (10 cases per group) was conducted. It was revealed that the mean  $\pm$  standard deviation (SD) of HR following the installation of the wetting solution during liposuction (i.e., primary outcome) was 92.02  $\pm$  12.73 in the epinephrine group, 80.88  $\pm$  15.20 in the IV MG group, and 86.71  $\pm$  18.76 in the SC MG group. The sample size was determined considering the following points: Effect size = 0.29, 95% confidence limit, and 80% study power. Four cases were added to each group

to eliminate the dropout effect; hence, 43 patients were assigned to each group.

#### 3.3. Statistical Analysis

Statistical analysis was performed with SPSS software version 26 (IBM Inc., Chicago, IL, USA). Shapiro-Wilks normality test and histograms were used to test the distribution of quantitative variables. Parametric variables were expressed as mean and SD and compared using ANOVA. Categorial variables were expressed as frequency and percentage and then statistically analyzed using chi-square tests. A two-tailed P-value  $\leq 0.05$  was considered the significance level in this study.

# 4. Results

In this study, 152 patients were assessed for eligibility, among whom 14 did not meet the inclusion criteria, and 9 refused to participate. Accordingly, 129 patients were randomly assigned into 3 equal groups (n = 43 per group) and then followed up and analyzed statistically (Figure 1).

The patients' demographics and duration of surgery showed no significant difference among the three groups (P > 0.05) (Table 1).

Basal and induction HRs were not significantly different among the three studied groups, and intraoperative HR was not significantly different regarding all measurements between the  $MgSO_4$  groups.

Intraoperative HR was significantly lower in group II compared to group I from wetting to 120 min, and it was significantly lower in group III compared to group I at wetting, 30, 45, and 105 min; however, there was no significant difference between group III and group I regarding these measurements (Table 2).

At baseline and induction, MAPs were insignificant among the three studied groups. However, MAP was significantly lower in the MgSO<sub>4</sub> groups (groups II and III) compared to group I, and it was significantly lower in group II compared to group III (Table 3).

Intraoperative isoflurane consumption was significantly lower in the MgSO<sub>4</sub> groups (groups II and III) compared to group I, with no significant difference between the MgSO<sub>4</sub> groups (P < 0.001). Postoperative morphine consumption revealed no significant difference among the research groups.

Regarding epinephrine-related side effects, sinus tachycardia occurred in 51.2% of the patients in group I, 25.6% in group II, and 27.9% in group III which was significantly lower in group I and in group III compared to group I (P = 0.014 and 0.027 respectively) and insignificantly different between group II and group III. PVCs occurred in 11.6% of the patients in group I, 0.0% of the

patients in group II, and no patients in group III. PVCs were significantly lower in the  $MgSO_4$  groups compared to group I (Table 4).

The VAS scores were significantly lower in the MgSO<sub>4</sub> groups than in group I. Nevertheless, the three groups showed statistically significant VAS scores after 12 and 24 hours (Table 5).

# 5. Discussion

The wetting solution injected during the liposuction procedure should inform anesthesiologists of some important physiological considerations (3, 16). Most injected fluids are transferred into the general circulation; however, only 22 - 29% of the fluids are recovered during the liposuction procedure (17, 18). This implies that about 2/3 of the injectate with its added drugs (including epinephrine) transfers to the intravascular compartment (19). A previous study confirmed that 70% of injected epinephrine is reabsorbed into the systemic circulation, with a maximum peak after 5 hours (20).

Some studies have reported epinephrine toxicity following the accidental subcutaneous infiltration of 3 mg 1: 1000 (21-23). Toxicity may be associated with hypertension, tachycardia, arrhythmia, chest pain, diaphoresis, and limb paresthesia, leading to pulmonary edema, asystole, and mortality (9).

Previous studies dealing with liposuction-induced complications have stated that cardiac side effects requiring therapy were observed in < 1% of the cases undergoing surgery (24, 25). Although their prevalence is not high, they may be fatal. Moreover, previous studies are not conducted recently and used wetting solutions with lower doses of epinephrine.

Few studies have discussed preventing such cardiac complications, and there is no rich literature on this issue.

Honarmand et al. (13) reported that  $MgSO_4$  in doses < 50 mg/kg could effectively reduce the cardiovascular instability associated with intubation. They tested 3 different doses of this drug (30, 40, and 50 mg/kg), according to which we decided to use the intermediate dose proposed by these researchers in the present study.

In this study, although basal and induction HR were not significantly different among the research groups, intraoperative HR was significantly lower in the MgSO<sub>4</sub> groups compared to group I in most intraoperative readings. In the same context, Elsharnouby and Elsharnouby reported a significant decline in intraoperative HR readings following the administration of IV MgSO<sub>4</sub> (26). Stanbury reported that MgSO<sub>4</sub> prolongs nodal recovery time using both direct and indirect mechanisms (27). It also slows HR at rest by blocking the nictitating membrane of



the sympathetic ganglia (13). Furthermore, it has a negative chronotropic effect following the administration of atropine (28).

According to the findings, intraoperative MAP was significantly lower in  $MgSO_4$  readings compared to group I. As mentioned in previous reports,  $MgSO_4$  does not affect cardiac output; hence, its effect on blood pressure is mediated by decreasing the peripheral arterial resistance using its vasodilator action (29, 30). This finding has been documented in vitro (31, 32), animal (33), and human studies (34, 35). Our findings showed a significant decrease in isoflurane consumption following the MgSO<sub>4</sub> administration compared to the controls. Likewise, another study reported lower doses of volatile anesthetic requirement following the administration of MgSO<sub>4</sub> (26). Moreover, James reported that MgSO<sub>4</sub> was used to produce general anesthesia and enhance the action of anesthetic agents (36). This effect could be explained by the antagonistic effect of MgSO<sub>4</sub> on N-methyl-d-aspartate (NMDA) receptors, decreased catecholamine production by decreasing sympathetic stimulation, and decreased nociceptor sensitization

<b>Fable 1.</b> Demographic Characteristics and Duration of Surgery of 3 Research Groups <sup>a</sup>							
Variables	Group I (n = 43)	Group II (n = 43)	Group III (n = 43)	Р			
Age (y)	$35.16 \pm 8.748$	$37.33 \pm 10.200$	$36.84 \pm 8.174$	0.513			
Gender				0.215			
Male	18 (41.9)	11% (25.6)	12% (27.9)				
Female	25 (58.1)	32% (74.4)	31% (72.1)				
BMI (kg/m <sup>2</sup> )	$31.59 \pm 2.013$	$31.85 \pm 1.798$	$31.91 \pm 1.983$	0.724			
ASA physical status				0.376			
Ι	20 (46.5)	15% (34.9)	21% (48.8)				
II	23 (53.5)	28% (65.1)	22% (51.2)				
Duration of surgery (h)	$146.70\pm18.55$	$150.62 \pm 16.97$	$153.37 \pm 19.68$	0.247			

Abbreviations: BMI, body mass index; ASA, American Society of Anesthesiologists.

<sup>a</sup> Data presented as mean  $\pm$  SD or No. (%).

Fable 2. Intra-operative Heart Rate Follow-up in 3 Groups <sup>a</sup>								
Heart rate (bpm)	Group I (n = 43)	Group II (n = 43)	Group III (n = 43)	Р	P1 <sup>b</sup>	P2 <sup>c</sup>	P3 <sup>d</sup>	
Basal	$80.49\pm9.468$	80.81±11.076	79.00±13.765	0.741	_	_	_	
Induction	$85.95 \pm 9.388$	86.21±12.007	83.93 ± 15.363	0.652	-	-	-	
Wetting	$98.02\pm12.805$	$87.49 \pm 13.754$	$89.86 \pm 15.824$	0.002	0.002	0.026	1	
15 min	$100.12 \pm 12.004$	88.91±13.151	$95.02\pm18.089$	0.002	0.002	0.329	0.165	
30 min	97.60 ± 12.600	$85.95 \pm 12.928$	$89.09 \pm 20.892$	0.003	0.003	0.044	1	
45 min	92.67± 11.704	83.00 ± 12.786	$85.51 \pm 15.182$	0.003	0.003	0.041	1	
60 min	$88.70 \pm 11.281$	$79.21 \pm 12.779$	83.37±16.311	0.007	0.005	0.217	0.482	
75 min	88.65±11.080	$78.98 \pm 12.540$	$82.60 \pm 20.096$	0.013	0.011	0.197	0.814	
90 min	$88.70\pm12.474$	$79.28\pm12.374$	$81.12 \pm 18.931$	0.010	0.012	0.060	1	
105 min	88.70 ± 12.260	$79.37 \pm 12.458$	80.98±17.805	0.007	0.010	0.043	1	
120 min	88.81± 11.576	$79.56 \pm 12.655$	$83.40\pm18.564$	0.015	0.012	0.263	0.677	

<sup>a</sup> Data presented as mean  $\pm$  SD.

<sup>b</sup> Group I vs. group II

Group I vs. group III

<sup>d</sup> Group II vs. group III

(37). Moreover, it decreases acetylcholine release at the neuromuscular junction (38, 39).

In the present study, the incidence of cardiac arrhythmia showed a significant decrease in MgSO<sub>4</sub> compared to group I. Sinus tachycardia was detected in 51.2, 25.6, and 27.9% of the cases. At the same time, PVCs were noticed in 11.6, 0, and 0 of the participants in groups I, II, and III, respectively. Magnesium sulfate is often used as an antiarrhythmic agent by many anesthesiologists in critical intraoperative care settings (40). Moreover, it is usually administered for prophylaxis and the management of cardiac arrhythmia following cardiac operations (41). It decreases the sinoatrial node (SA node) impulse rate; however, it induces refractory period prolongation in the atrioventricular node (AV node)(42, 43). In their study, Saran et al. confirmed the decreased incidence of cardiac arrhythmia following the  $MgSO_4$  administration for patients undergoing thoracotomy (43).

Our findings showed that group I significantly expressed higher VAS scores than the other two groups (P < 0.05). Several studies have confirmed the analgesic effects of MgSO<sub>4</sub> (44), which can induce the suppression of neuropathic pain, potentiate morphine-induced analgesia, and decrease the level of tolerance to opioids (45, 46). The mechanism of this analgesic effect has yet to be completely understood. However, it is assumed to be mediated by regulating calcium influx in the cells and antagonizing NMDA receptors in the central nervous system (47, 48).

The previously reported pain outcomes exhibited their effects on postoperative morphine consumption, which

Table 3. Intra-operative Mean Arterial Blood Pressure Follow-up in 3 Groups <sup>a</sup>								
MAP(mmHg)	Group I (n = 43)	Group II (n = 43)	Group III (n = 43)	Р	P1 <sup>b</sup>	P2 <sup>c</sup>	P3 <sup>d</sup>	
Basal	97.33± 9.286	96.63± 6.786	$94.88\pm9.027$	0.388	_	_	-	
Induction	$103.53 \pm 9.876$	$100.30\pm7.418$	99.60 ± 11.681	0.145	-	-	-	
Wetting	$112.70 \pm 13.534$	$94.02 \pm 9.595$	$100.58 \pm 13.548$	< 0.001	< 0.001	< 0.001	0.035	
15 min	$114.23 \pm 13.325$	$88.60 \pm 9.725$	$100.49 \pm 13.188$	< 0.001	< 0.001	< 0.001	< 0.001	
30 min	$107.23 \pm 14.410$	$80.93 \pm 10.366$	97.93±12.747	< 0.001	< 0.001	0.003	< 0.001	
45 min	$99.72\pm14.413$	$75.40\pm10.347$	$90.40 \pm 13.968$	< 0.001	< 0.001	0.004	< 0.001	
60 min	$94.40 \pm 14.721$	$71.95 \pm 10.021$	86.30 ± 13.562	< 0.001	< 0.001	0.013	< 0.001	
75 min	$95.26 \pm 14.206$	$71.86 \pm 10.218$	85.21±13.731	< 0.001	< 0.001	0.001	< 0.001	
90 min	$95.93 \pm 15.890$	$72.09\pm10.768$	85.95±13.856	< 0.001	< 0.001	0.003	< 0.001	
105 min	$95.72 \pm 15.443$	$72.14 \pm 10.255$	86.07±14.757	< 0.001	< 0.001	0.004	< 0.001	
120 min	$95.05 \pm 14.705$	$72.42\pm10.658$	87.77±13.082	< 0.001	< 0.001	0.030	< 0.001	

Abbreviation: MAP, mean arterial blood pressure. <sup>a</sup> Data presented as mean ± SD. <sup>b</sup> Group I vs. group II <sup>c</sup> Group I vs. group III <sup>d</sup> Group II vs. group III

Variables	Group I (n = 43)	Group II (n = 43)	Group III (n = 43)	Р	P1 <sup>b</sup>	P2 <sup>c</sup>	P3 <sup>d</sup>
Intraoperative isoflurane consumption (mL)	100.23 ± 24.832	83.49 ± 19.009	$76.05 \pm 15.452$	< 0.001	0.001	< 0.001	0.240
Postoperative morphine consumption (mg)	$2.23 \pm 1.324$	$1.79\pm0.965$	$1.77\pm1.130$	0.111			
Arrhythmia							
Sinus tachycardia	22 (51.2)	11 (25.6)	12 (27.9)	0.023	0.014	0.027	0.807
PVCs	5 (11.6)	0(0.0)	0(0.0)	0.005	0.039	0.021	_

Abbreviation: PVCs, premature ventricular contractions. <sup>a</sup> Data presented as mean ± SD or No. (%). <sup>b</sup> Group I vs. group II <sup>c</sup> Group I vs. group III <sup>d</sup> Group II vs. group III

<b>Fable 5.</b> Visual Analog Scale Follow-up Score in Research Groups <sup>a</sup>								
VAS	Group I (n = 43)	Group II (n = 43)	Group III (n = 43)	Р	P1 <sup>b</sup>	P2 <sup>c</sup>	P3 <sup>d</sup>	
PACU	$4.60 \pm 1.198$	$2.84 \pm 1.132$	3.35 ± 1.412	< 0.001	< 0.001	< 0.001	0.182	
1 hr	$4.12\pm1.401$	$2.95\pm1.253$	$3.02\pm1.389$	< 0.001	< 0.001	0.001	1	
3 hr	$2.91 \pm 0.895$	$2.60 \pm 0.877$	$2.26 \pm 0.875$	0.004	0.344	0.003	0.207	
6 hr	$2.88 \pm 1.418$	2.21± 0.965	$2.44 \pm 1.098$	0.029	0.026	0.251	1	
12 hr	$2.58 \pm 1.006$	$2.16\pm1.022$	$2.35\pm0.948$	0.151				
24 hr	$2.16\pm1.045$	$2.12\pm0.851$	$1.91 \pm 0.996$	0.428				

Abbreviations: VAS, visual analogue scale; PACU, post anesthesia care unit. <sup>a</sup> Data presented as mean ± SD. <sup>b</sup> Group I vs. group II <sup>c</sup> Group I vs. group III <sup>d</sup> Group I vs. group III

was increased in Group I; however, it was not significant (P = 0.111). Another study confirmed the slight but nonsignificant decrease in morphine need during the postoperative period after the MgSO<sub>4</sub> administration. The insignificance of the statistical analysis results was probably due to the included small sample size (49).

To sum up, one should notice the significant positive effects of the IV or subcutaneous administration of  $MgSO_4$ . The effects of its subcutaneous administration could be explained by its systemic absorption, as reported earlier in this discussion. Its administration should be encouraged by surgeons, especially for patients at high risk of epinephrine-related cardiac events during the liposuction procedure.

This study had some limitations; first, it was a singlecenter study, including relatively small sample size. Moreover, we should have evaluated serum magnesium levels in the receiving groups to detect existing correlations between its levels and hemodynamic variation. This would have also helped to estimate the level of  $MgSO_4$  absorbed into the systemic circulation in group III. Accordingly, further studies on more patients from different plastic centers are recommended.

## 5.1. Conclusions

The IV or subcutaneous administration of  $MgSO_4$  is associated with lower intraoperative HR, MAP, postoperative pain score, and a significant decrease in epinephrineinduced cardiac adverse effects during liposuction. Further positive effects were observed for the IV administration.

#### Footnotes

**Authors' Contribution:** M. A. A. developed the original idea and the research protocol, abstracted and analyzed data, wrote the manuscript, and is a guarantor. A. E., S. H., and D. E. developed the protocol, abstracted data, and prepared the manuscript. All authors revised the manuscript.

ClinicalTrialRegistrationCode:PACTR202105734020705.

Conflict of Interests: There is none to be declared.

**Data Reproducibility:** The data presented in this study are openly available for readers upon request.

**Ethical Approval:** The study was approved by the Ethics Committee of Tanta University (Code: 35029/11/21).

Funding/Support: There is none to be declared.

**Informed Consent:** Informed written consent was obtained from all participants.

## References

- 1. Taha AA, Tahseen H. Liposuction: Keeping It Safe!. *Plast Reconstr Surg Glob Open*. 2020;**8**(4). e2783. [PubMed ID: 32440445]. [PubMed Central ID: PMC7209863]. https://doi.org/10.1097/GOX.00000000002783.
- Taha AA, Tahseen H. Pneumothorax with Liposuction: Spreading Awareness. *Plast Reconstr Surg Glob Open*. 2020;8(3). e2711. [PubMed ID: 32537362]. [PubMed Central ID: PMC7253280]. https://doi.org/10.1097/GOX.00000000002711.
- Mendez BM, Coleman JE, Kenkel JM. Optimizing Patient Outcomes and Safety With Liposuction. *Aesthet Surg J.* 2019;**39**(1):66–82. [PubMed ID: 29947738]. https://doi.org/10.1093/asj/sjy151.
- Garcia O. Anesthesia and Wetting Solutions. In: Garcia O, editor. Ultrasound-Assisted Liposuction. New York City, USA: Springer Cham; 2020. p. 37-45. https://doi.org/10.1007/978-3-030-26875-6\_4.
- Sood J, Jayaraman L, Sethi N. Liposuction: Anaesthesia challenges. Indian J Anaesth. 2011;55(3):220–7. [PubMed ID: 21808392]. [PubMed Central ID: PMC3141144]. https://doi.org/10.4103/0019-5049.82652.
- Bellini E, Grieco MP, Raposio E. A journey through liposuction and liposculture: Review. Ann Med Surg (Lond). 2017;24:53-60. [PubMed ID: 29158895]. [PubMed Central ID: PMC5681335]. https://doi.org/10.1016/j.amsu.2017.10.024.
- Platt MS, Kohler LJ, Ruiz R, Cohle SD, Ravichandran P. Deaths associated with liposuction: case reports and review of the literature. J Forensic Sci. 2002;47(1):205-7. [PubMed ID: 12064654].
- Grazer FM, de Jong RH. Fatal outcomes from liposuction: census survey of cosmetic surgeons. *Plast Reconstr Surg.* 2000;**105**(1):436–46. discussion 447-8. [PubMed ID: 10627013]. https://doi.org/10.1097/00006534-200001000-00070.
- Brown SA, Lipschitz AH, Kenkel JM, Sorokin E, Shepherd G, Grebe S, et al. Pharmacokinetics and safety of epinephrine use in liposuction. *Plast Reconstr Surg.* 2004;**114**(3):756–63. discussion 764-5. [PubMed ID: 15318058]. https://doi.org/10.1097/01.prs.0000131021.17800.be.
- Marzetti E, Csiszar A, Dutta D, Balagopal G, Calvani R, Leeuwenburgh C. Role of mitochondrial dysfunction and altered autophagy in cardiovascular aging and disease: from mechanisms to therapeutics. *Am J Physiol Heart Circ Physiol*. 2013;**305**(4):H459– 76. [PubMed ID: 23748424]. [PubMed Central ID: PMC3891249]. https://doi.org/10.1152/ajpheart.00936.2012.
- Kostov K, Halacheva L. Role of Magnesium Deficiency in Promoting Atherosclerosis, Endothelial Dysfunction, and Arterial Stiffening as Risk Factors for Hypertension. *Int J Mol Sci.* 2018;19(6). [PubMed ID: 29891771]. [PubMed Central ID: PMC6032400]. https://doi.org/10.3390/ijms19061724.
- Severino P, Netti L, Mariani MV, Maraone A, D'Amato A, Scarpati R, et al. Prevention of Cardiovascular Disease: Screening for Magnesium Deficiency. *Cardiol Res Pract.* 2019;**2019**:4874921. [PubMed ID: 31192005]. [PubMed Central ID: PMC6525869]. https://doi.org/10.1155/2019/4874921.
- Honarmand A, Safavi M, Badiei S, Daftari-Fard N. Different doses of intravenous Magnesium sulfate on cardiovascular changes following the laryngoscopy and tracheal intubation: A doubleblind randomized controlled trial. *J Res Pharm Pract.* 2015;4(2):79– 84. [PubMed ID: 25984545]. [PubMed Central ID: PMC4418140]. https://doi.org/10.4103/2279-042X.154365.
- Waxman HL, Myerburg RJ, Appel R, Sung RJ. Verapamil for control of ventricular rate in paroxysmal supraventricular tachycardia and atrial fibrillation or flutter: a double-blind randomized cross-over study. Ann Intern Med. 1981;94(1):1–6. [PubMed ID: 7447203]. https://doi.org/10.7326/0003-4819-94-1-1.
- Adhikary SD, Pruett A, Forero M, Thiruvenkatarajan V. Erector spinae plane block as an alternative to epidural analgesia for postoperative analgesia following video-assisted thoracoscopic surgery: A case study and a literature review on the spread of local anaesthetic in the erector spinae plane. *Indian J Anaesth.* 2018;62(1):75-8. [PubMed ID: 29416155]. [PubMed Central ID: PMC5787896]. https://doi.org/10.4103/ija.IJA\_693\_17.

- Halk AB, Habbema L, Genders RE, Hanke CW. Safety Studies in the Field of Liposuction: A Systematic Review. *Dermatol Surg.* 2019;45(2):171-82. [PubMed ID: 30702445]. https://doi.org/10.1097/DSS.00000000001707.
- Samdal F, Amland PF, Bugge JF. Blood loss during liposuction using the tumescent technique. *Aesthetic Plast Surg.* 1994;18(2):157–60. [PubMed ID: 8017219]. https://doi.org/10.1007/BF00454476.
- Pitman GH, Aker JS, Tripp ZD. Tumescent liposuction. A surgeon's perspective. *Clin Plast Surg.* 1996;23(4):633–41. discussion 642-5. [PubMed ID: 8906394].
- Marin JEB. Liquid handling, lidocaine and epinephrine in liposuction. The properly form. *Colomb J Anesthesiol*. 2015;43(1):95–100. https://doi.org/10.1016/j.rcae.2014.10.003.
- Trott SA, Beran SJ, Rohrich RJ, Kenkel JM, Adams WP, Klein KW. Safety considerations and fluid resuscitation in liposuction: an analysis of 53 consecutive patients. *Plast Reconstr Surg.* 1998;**102**(6):2220– 9. [PubMed ID: 9811024]. https://doi.org/10.1097/00006534-199811000-00063.
- Fyfe AI, Daly PA, Dorian P, Tough J. Reversible "cardiomyopathy" after accidental adrenaline overdose. *Am J Cardiol.* 1991;67(4):318–9. [PubMed ID: 1990800]. https://doi.org/10.1016/0002-9149(91)90570-b.
- 22. Karch SB. Coronary artery spasm induced by intravenous epinephrine overdose. *Am J Emerg Med.* 1989;7(5):485–8. [PubMed ID: 2757714]. https://doi.org/10.1016/0735-6757(89)90250-7.
- Larsen LS, Larsen A. Labetalol in the treatment of epinephrine overdose. Ann Emerg Med. 1990;19(6):680-2. [PubMed ID: 2344087]. https://doi.org/10.1016/s0196-0644(05)82478-3.
- Dillerud E. Suction lipoplasty: a report on complications, undesired results, and patient satisfaction based on 3511 procedures. *Plast Recon*str Surg. 1991;88(2):239–46. discussion 247-9. [PubMed ID: 1852816].
- Hanke CW, Bernstein G, Bullock S. Safety of tumescent liposuction in 15,336 patients. National survey results. *Dermatol Surg.* 1995;21(5):459–62. [PubMed ID: 7743109]. https://doi.org/10.1111/j.1524-4725.1995.tb00213.x.
- Elsharnouby NM, Elsharnouby MM. Magnesium sulphate as a technique of hypotensive anaesthesia. *Br J Anaesth*. 2006;96(6):727-31. [PubMed ID: 16670112]. https://doi.org/10.1093/bja/ael085.
- Stanbury JB. The blocking action of magnesium ion on sympathetic ganglia. J Pharmacol Exp Ther. 1948;93(1):52–62. [PubMed ID: 18865186].
- Somjen GG, Baskerville EN. Effect of excess magnesium on vagal inhibition and acetylcholine sensitivity of the mammalian heart in situ and in vitro. *Nature*. 1968;217(5129):679–80. [PubMed ID: 5637751]. https://doi.org/10.1038/217679a0.
- Nakaigawa Y, Akazawa S, Shimizu R, Ishii R, Ikeno S, Inoue S, et al. Effects of magnesium sulphate on the cardiovascular system, coronary circulation and myocardial metabolism in anaesthetized dogs. Br J Anaesth. 1997;**79**(3):363-8. [PubMed ID: 9389857]. https://doi.org/10.1093/bja/79.3.363.
- Kamble SP, Bevinaguddaiah Y, Nagaraja DC, Pujar VS, Anandaswamy TC. Effect of Magnesium Sulfate and Clonidine in Attenuating Hemodynamic Response to Pneumoperitoneum in Laparoscopic Cholecystectomy. *Anesth Essays Res.* 2017;**11**(1):67–71. [PubMed ID: 28298759]. [PubMed Central ID: PMC5341635]. https://doi.org/10.4103/0259-1162.200228.
- Turlapaty PD, Altura BM. Extracellular magnesium ions control calcium exchange and content of vascular smooth muscle. Eur J Pharmacol. 1978;52(3-4):421-3. [PubMed ID: 729651]. https://doi.org/10.1016/0014-2999(78)90303-5.
- Altura BM, Altura BT. Magnesium and contraction of arterial smooth muscle. *Microvasc Res.* 1974;7(2):145–55. [PubMed ID: 4363012]. https://doi.org/10.1016/0026-2862(74)90001-6.
- James MF, Cork RC, Harlen GM, White JF. Interactions of adrenaline and magnesium on the cardiovascular system of the baboon. *Magnesium*. 1988;7(1):37-43. [PubMed ID: 3379980].

- Zhang J, Wang Y, Xu H, Yang J. Influence of magnesium sulfate on hemodynamic responses during laparoscopic cholecystectomy: A meta-analysis of randomized controlled studies. *Medicine (Baltimore)*. 2018;97(45). e12747. [PubMed ID: 30407279]. [PubMed Central ID: PMC6250549]. https://doi.org/10.1097/MD.000000000012747.
- Ray M, Bhattacharjee DP, Hajra B, Pal R, Chatterjee N. Effect of clonidine and magnesium sulphate on anaesthetic consumption, haemodynamics and postoperative recovery: A comparative study. *Indian J Anaesth*. 2010;54(2):137–41. [PubMed ID: 20661352]. [PubMed Central ID: PMC2900737]. https://doi.org/10.4103/0019-5049.63659.
- James MF. Clinical use of magnesium infusions in anesthesia. Anesth Analg. 1992;74(1):129–36. [PubMed ID: 1734773]. https://doi.org/10.1213/00000539-199201000-00021.
- Dube L, Granry JC. The therapeutic use of magnesium in anesthesiology, intensive care and emergency medicine: a review. Can J Anaesth. 2003;50(7):732–46. [PubMed ID: 12944451]. https://doi.org/10.1007/BF03018719.
- Fuchs-Buder T, Tassonyi E. Magnesium sulphate enhances residual neuromuscular block induced by vecuronium. Br J Anaesth. 1996;76(4):565–6. [PubMed ID: 8652332]. https://doi.org/10.1093/bja/76.4.565.
- Frakes MA, Richardson LE. Magnesium sulfate therapy in certain emergency conditions. *Am J Emerg Med*. 1997;15(2):182-7. [PubMed ID: 9115526]. https://doi.org/10.1016/s0735-6757(97)90098-x.
- Soave PM, Conti G, Costa R, Arcangeli A. Magnesium and anaesthesia. *Curr Drug Targets*. 2009;**10**(8):734–43. [PubMed ID: 19702521]. https://doi.org/10.2174/138945009788982487.
- Sirvinskas E, Laurinaitis R. [Use of magnesium sulfate in anesthesiology]. *Medicina (Kaunas)*. 2002;**38**(7):695–8. Lithuanian. [PubMed ID: 12474653].
- James MF. Magnesium: an emerging drug in anaesthesia. Br J Anaesth. 2009;103(4):465–7. [PubMed ID: 19749114]. https://doi.org/10.1093/bja/aep242.
- 43. Saran T, Perkins GD, Javed MA, Annam V, Leong L, Gao F, et al. Does the prophylactic administration of magnesium sulphate to patients undergoing thoracotomy prevent postoperative supraventricular arrhythmias? A randomized controlled trial. Br J Anaesth. 2011;106(6):785–91. [PubMed ID: 21558066]. https://doi.org/10.1093/bja/aer096.
- 44. Albrecht E, Kirkham KR, Liu SS, Brull R. Peri-operative intravenous administration of magnesium sulphate and postoperative pain: a meta-analysis. *Anaesthesia*. 2013;68(1):79–90. [PubMed ID: 23121612]. https://doi.org/10.1111/j.1365-2044.2012.07335.x.
- Feria M, Abad F, Sanchez A, Abreu P. Magnesium sulphate injected subcutaneously suppresses autotomy in peripherally deafferented rats. *Pain.* 1993;**53**(3):287-93. [PubMed ID: 8351158]. https://doi.org/10.1016/0304-3959(93)90225-E.
- 46. McCarthy RJ, Kroin JS, Tuman KJ, Penn RD, Ivankovich AD. Antinociceptive potentiation and attenuation of tolerance by intrathecal co-infusion of magnesium sulfate and morphine in rats. Anesth Analg. 1998;86(4):830-6. [PubMed ID: 9539610]. https://doi.org/10.1097/00000539-199804000-00028.
- Mostafa MF, Herdan R, Fathy GM, Hassan Z, Galal H, Talaat A, et al. Intrathecal dexmedetomidine versus magnesium sulphate for postoperative analgesia and stress response after caesarean delivery; randomized controlled double-blind study. *Eur J Pain*. 2020;24(1):182–91. [PubMed ID: 31461801]. https://doi.org/10.1002/ejp.1476.
- McKeown A, Seppi V, Hodgson R. Intravenous Magnesium Sulphate for Analgesia after Caesarean Section: A Systematic Review. Anesthesiol Res Pract. 2017;2017:9186374. [PubMed ID: 29333156]. [PubMed Central ID: PMC5733151]. https://doi.org/10.1155/2017/9186374.
- Wilder-Smith CH, Knopfli R, Wilder-Smith OH. Perioperative magnesium infusion and postoperative pain. Acta Anaesthesiol Scand. 1997;41(8):1023-7. [PubMed ID: 9311401]. https://doi.org/10.1111/j.1399-6576.1997.tb04830.x.