



Relationship Between Pain and Disability Among Stroke Patients

Aryoobarzan Rahmatian¹, Ali Karbasfrushan² and Farhad Modara^{1,*}

¹School of Medicine, Shahid Mostafa Khomaeini Hospital, Ilam University of Medical Sciences, Ilam, Iran

²Department of Anesthesiology, School of Medicine, Imam Khomeini Hospital, Imam Reza Hospital, Kermanshah University of Medical Sciences, Kermanshah, Iran

*Corresponding author: School of Medicine, Shahid Mostafa Khomaeini Hospital, Ilam University of Medical Sciences, Ilam, Iran. Email: farhadmodara2001@gmail.com

Received 2023 March 14; Revised 2023 April 08; Accepted 2023 April 12.

Abstract

Background: Stroke is prevalent in Iran, and its complications can decline patients' quality of life and psychological state.

Objectives: This study aimed to investigate the relationship between pain following stroke (PFS) and disability in stroke patients.

Methods: The current case-control study was conducted on 184 stroke patients (92 cases and 92 controls). Data collection tools included a demographic questionnaire, the Visual Analog Scale (VAS), and the Disabilities of the Arm Shoulder and Hand (DASH) questionnaire. Based on the medical files of all patients with stroke, eligible ones were invited to participate in the study via phone calls. The collected data were then analyzed.

Results: The disability rate was higher in the case group of patients with a history of hemorrhagic stroke, smoking, stroke, and diabetes ($P < 0.05$). The mean \pm SD age was 72.45 (11.54). Also, the level of disability in patients increased with age ($P = 0.000$). The mean \pm SD disability score was 78.63 (6.92) in patients with PFS and 54.9 (10.87) in patients without PFS. Also, $R = 0.795$ and $R^2 = 0.631$ indicated the significance of disability severity in patients with PFS.

Conclusions: The prevalence of disability was higher in patients with higher PFS levels. Hence, drug interventions or rehabilitation programs can be used to reduce the disability of stroke patients.

Keywords: Shoulder Pain, Stroke, Disability

1. Background

The stroke incidence rate increases in men and women with age, affecting almost 50% and 30% of them over 75 and 85 years old, respectively (1-3). Cerebrovascular accident (CVA) is a neurological defect with a local vascular origin that causes death and disability worldwide and affects the lives of millions of people (4, 5). After heart disease and cancer, stroke is the third cause of death worldwide, affecting about 795,000 people in the United States yearly, of whom 185,000 experience a recurrent stroke (6, 7).

Ischemic stroke, hemorrhagic stroke, and transient ischemic attacks are caused by different types of strokes, and symptoms and location of CVA are different in each type. They can last a few hours or more or sometimes lead to the patient's death (8, 9). In these patients, symptoms include a general inability to move or decreased ability and strength of muscles to move the limbs on one side of the body, difficulty speaking, and dizziness. Also, stroke patients have difficulty regulating limb movement due to abnormal reflex activities and muscle tone. The post-stroke mechanical changes are considered a limiting factor for the patient's voluntary activities (10, 11).

In Nikbakht et al.'s meta-analysis study of 17 articles, the hospital mortality rate was 18.71%, and the one-month mortality rate was 23.43% (12). Also, in the study of Krishnamurthi et al., the global prevalence of hemorrhagic stroke was reported as 3,725,085, and the prevalence of ischemic stroke was 7,258,216 (13). Stroke complications include pain and disability, which lead to a decline in patients' quality of life and psychological state (14, 15). Reducing or relieving patients' pain is important (16-18).

Pain following stroke (PFS) affects different parts of the body, such as the shoulder, knee, and head, and negatively affects the daily activities of these patients and their return to work (19, 20). Also, depression, stress, and fatigue are other complications of this disease, which can cause short-term and long-term psychological complications and ultimately reduce the quality of life (21, 22). These symptoms and complications have caused disability in these patients. According to previous studies, about 75% of stroke survivors have reported different degrees of disability, which requires attention (23). Another complication of CVA is pain which causes various complications; the patient faces various challenges, including disruption in daily activities

and psychological problems (3, 24).

2. Objectives

The present study aimed to determine the relationship between PFS and disability in stroke patients.

3. Methods

3.1. Study Design and Population

The current study was conducted on CVA patients from Ilam City in 2022. Consecutive purposive sampling was used. The study was conducted in a Hospital in Ilam City on 184 patients (92 cases and 92 controls).

3.2. Inclusion and Exclusion Criteria

3.2.1. Inclusion Criteria

The inclusion criteria included a diagnosis of CVA by a neurologist, suffering from CVA for the first time, suffering from CVA for at least six months, informed consent to participate in the study, 30 - 80 years of age, speaking ability, being hospitalized in Ilam city in the past, the physical and mental ability of the patient to complete the research instrument, and having access to make a phone call.

3.2.2. Exclusion Criteria

They included suffering from a chronic disease affecting the pain condition, psychological stress during the last three months, confirmed mental disorders, rheumatic disorders, history of trauma or shoulder involvement, bilateral brain involvement, and tumor or brain trauma.

3.3. Data Gathering

Data collection tools included a demographic questionnaire, the Visual Analog Scale (VAS) (25), and the Disabilities of the Arm Shoulder and Hand (DASH) Questionnaire (26).

3.3.1. Demographic Characteristics

3.3.1.1. Demographic Characteristics Questionnaire

This instrument included questions on age, gender, affected side, dominant side, duration of illness in months, stroke type (hemorrhagic or thrombolytic), smoking history, family history of CVA, history of diabetes, and hypertension.

3.3.2. Visual Analog Scale

The presence of shoulder pain for at least two days was rated using the VAS (27). All patients were evaluated for the presence or absence of shoulder muscle spasticity and shoulder-hand syndrome. The evaluation was stopped if the participants felt pain or discomfort during any of the evaluation stages (25). If there was pain, the patients were placed in the group with PFS; if there was no pain, they were placed in the group without PFS.

3.3.3. Disabilities of the Arm Shoulder and Hand Scale

This instrument consists of 30 questions to evaluate the function of the person's upper limb in the last week. Each question is assigned a score from 1 to 5, and the overall score is 100. If the score is closer to 100, it indicates a more severe disability (26).

3.4. Research Method

This case-control study used a case group of patients with PFS and a control group of patients without PFS. A total of 184 people were examined, 92 in the PFS group and 92 in the non-PFS group.

Based on the medical file of all patients with CVA referred to a hospital in Ilam City, eligible ones were invited to participate in the study via phone calls. The participants entered the study after obtaining their informed consent, explaining the study's objectives, and being assured about the confidentiality of their answers. It was also possible for patients to withdraw from the study at any stage. Since most patients were illiterate, the interview method was used to complete the questionnaires. The questionnaires were supposed to complete within three months after CVA. Ethical considerations in the research included obtaining informed consent and not imposing costs on the patient. Also, all ethical guidelines were followed per the Ilam University of Medical Sciences.

3.5. Data Analysis

The frequency and percentage of demographic variables were reported as descriptive indicators. Then, the ANOVA test and *t*-test were used to compare the disability scores between the groups. A regression test was also used to associate the patients' disability status and pain. In all analyses, the significance level was < 0.05 , and SPSS 16 software was used for data analysis.

4. Results

Based on Table 1, the disability rate was higher in the case group of patients with a history of hemorrhagic

stroke, smoking, stroke, and diabetes ($P < 0.05$). The mean \pm SD age was 72.45 (11.54). Also, the level of disability in patients increased with age ($P = 0.000$).

The mean \pm SD disability score was 78.63 (6.92) in patients with PFS and 54.9 (10.87) in patients without PFS. Also, $R = 0.795$ and $R^2 = 0.631$ indicated the significance of disability severity in patients with PFS (Table 2).

5. Discussion

Chronic patients experience many complications, including pain, reduced quality of life, and psychological problems (28-30). Pain following stroke is also a complication (31, 32). In this study, most patients were elderly (72.45 \pm 11.54 years) and male. Stroke patients were in their old age and were male in various studies. For example, most patients were over 63 years of age and male in Diener et al.'s study (33), 66 years and male in Hart et al.'s study (over 60% of patients) (34), 63.6 \pm 11.2 years and male ($n = 85$, 72%) in Stiekema et al.'s study (35), and 70.9 \pm 17.3 and male (51.3%) in Al-Gibbawi et al.'s study (36).

According to the results, the prevalence of diabetes in patients was 5.4%, and the mean disability score was higher in diabetic patients. Xing et al. reported that the prevalence of diabetes in stroke patients was 24.5% (37). The prevalence of diabetes was also 18.8% in Regenhardt et al.'s study (38). Tu et al. also reported that diabetes affected 20.9% of patients (39), contrary to this study's results.

The present study also showed that the mean \pm SD disability scores in PFS and non-PFS patients were 78.63 (6.92) and 54.9 (10.87), respectively. In the study by Regenhardt et al., the prevalence of disability was 49% and 56% in patients with hemorrhagic CVA and patients with ischemic CVA, respectively (38). Tu et al. also stated that the prevalence of 12-month disability was 16.6% (39). Cucchiara et al. also reported that the disability rate varied from 9.6% to 18.2% (40). In Kelly et al.'s study, the mean \pm SD Rehabilitation Length of Stay (LOS) score was 33.7 \pm 25.6 (41), which indicates the high level of disability in stroke patients.

The present study also revealed more disability in patients with higher pain levels. In the meta-analysis of patients with low-back pain, Alamam et al. showed a relationship between disability and pain self-efficacy and catastrophizing (42), which indicates the effect of pain on the disability of patients, consistent with the results of the present study.

A few studies have been conducted on PFS status in stroke patients, and this study was conducted for the first time in Ilam City. One of the limitations of this study is

the lack of examination of other variables. Hence, it is suggested that future researchers investigate other variables related to disability in stroke patients. It is also suggested that meta-analysis studies be conducted on the difference in disability status in patients with and without PFS.

5.1. Conclusions

The prevalence of disability was higher in patients with higher pain levels in the present study. For this reason, we can use drug interventions or rehabilitation programs to reduce the disability of CVA patients.

Acknowledgments

We thank the Ilam University of Medical Sciences.

Footnotes

Authors' Contribution: Study concept and design: FM, AK, and AR; Acquisition of data: FM, AK, AR; Analysis and interpretation of data: FM, AK, and AR; Drafting of the manuscript: FM, AK, and AR; Critical revision of the manuscript for important intellectual content: FM, AK, and AR; Statistical analysis: FM, AK, and AR; Administrative, technical, and material support: FM, AK, and AR; Study supervision: FM, AK, and AR.

Conflict of Interests: Funding or research support: None; Employment: Ilam University of Medical Sciences; Personal financial interests: None; Stocks or shares in companies: None; Consultation fees: None; Patents: None; Personal or professional relations with organizations and individuals (parents and children, wife and husband, family relationships, etc.): None; Unpaid membership in a government or non-governmental organization: None; Are you one of the editorial board members or a reviewer of this journal? None.

Data Reproducibility: The dataset presented in the study is available on request from the corresponding author during submission or after publication.

Ethical Approval: The current study was conducted after approval by the Ethics Committee (IR.MEDILAM.REC.1401.267).

Funding/Support: Ilam University of Medical Sciences supported this research project with A-10-3673-1.

Informed Consent: Written informed consent was obtained from the patient or their guardians.

Table 1. Disability of Patients in the Case and Control Groups

Variables	No. (%)	Disability	
		Case, Mean \pm SD	Control, Mean \pm SD
Gender			
Male	56 (60.9)	77.53 \pm 7.19	52.98 \pm 8.05
Female	36 (39.1)	79.07 \pm 6.82	57.88 \pm 13.8
P-value		0.53	0.000
F		0.32	32.99
Type of CVA			
Ischemic stroke	63 (68.5)	77.88 \pm 6.18	55.72 \pm 11.51
Hemorrhagic stroke	29 (31.5)	79.70 \pm 7.82	53.12 \pm 9.26
P-value		0.002	0.22
F			1.51
Smoking			
Yes	36 (39.1)	81.02 \pm 8.01	54.79 \pm 9.70
No	56 (60.9)	76.88 \pm 5.43	54.97 \pm 11.65
P-value		0.000	0.24
F		29.05	1.38
History of stroke			
Yes	43 (46.7)	79.19 \pm 7.17	60.86 \pm 11.68
No	49 (53.3)	75.77 \pm 4.65	49.67 \pm 6.7
P-value		0.000	0.001
F		13.63	11.87
History of diabetes			
Yes	5 (5.4)	80.25 \pm 7.03	79.73 \pm 3.93
No	87 (94.6)	74.74 \pm 4.85	53.47 \pm 9.3
P-value		0.000	0.03
F		16.14	4.75

Table 2. Relationship Between Disability and Pain Following Stroke in Stroke Patients

Model	Sum of Squares	Mean Square	F	P-Value
Regression	25907.307	25907.307	311.765	0.000
Residual	15123.995	83.099		
Total	41031.302			

References

- Mozaffarian D, Benjamin EJ, Go AS, Arnett DK, Blaha MJ, Cushman M, et al. Heart disease and stroke statistics–2015 update: a report from the American Heart Association. *Circulation*. 2015;**131**(4):e29–322. [PubMed ID: 25520374]. <https://doi.org/10.1161/CIR.000000000000152>.
- Pang P, Zheng K, Wu S, Xu H, Deng L, Shi Y, et al. Baicalin Down-regulates RLRs Signaling Pathway to Control Influenza A Virus Infection and Improve the Prognosis. *Evid Based Complement Alternat Med*. 2018;**2018**:4923062. [PubMed ID: 29681974]. [PubMed Central ID: PMC5846362]. <https://doi.org/10.1155/2018/4923062>.
- de Sire A, Moggio L, Demeco A, Fortunato F, Spano R, Aiello V, et al. Efficacy of rehabilitative techniques in reducing hemiplegic shoulder pain in stroke: Systematic review and meta-analysis. *Ann Phys Rehabil Med*. 2022;**65**(5):101602. [PubMed ID: 34757009]. <https://doi.org/10.1016/j.rehab.2021.101602>.
- Schnakers C, Vanhauzenhuysse A, Giacino J, Ventura M, Boly M, Majerus S, et al. Diagnostic accuracy of the vegetative and minimally conscious state: clinical consensus versus standardized neurobehavioral assessment. *BMC Neurol*. 2009;**9**:35. [PubMed ID: 19622138]. [PubMed Central ID: PMC2718857]. <https://doi.org/10.1186/1471-2377-9-35>.
- Tahmasbi F, Madani Neishaboori A, Mardani M, Toloui A, Komlakh K, Azizi Y, et al. Efficacy of polyarginine peptides in the treatment of

- stroke: A systematic review and meta-analysis. *Brain Behav.* 2023;**13**(1):e2858. [PubMed ID: 36542540]. [PubMed Central ID: PMC9847609]. <https://doi.org/10.1002/brb3.2858>.
6. Alipour L, Moosazadeh M, Esmaili R, Akbari H, Shafipour V. Effects of early family-centered sensory stimulation on disability and rehabilitation of stroke patients. *J Mazandaran Univ Med Sci.* 2020;**30**(185):61-73.
 7. Surrena H. *Handbook for Brunner and Suddarth's textbook of medical-surgical nursing.* Lippincott Williams & Wilkins; 2010.
 8. Bodin J, Ha C, Chastang JF, Descatha A, Leclerc A, Goldberg M, et al. Comparison of risk factors for shoulder pain and rotator cuff syndrome in the working population. *Am J Ind Med.* 2012;**55**(7):605-15. [PubMed ID: 22213435]. <https://doi.org/10.1002/ajim.22002>.
 9. Song SY, Zhao XX, Rajah G, Hua C, Kang RJ, Han YP, et al. Clinical Significance of Baseline Neutrophil-to-Lymphocyte Ratio in Patients With Ischemic Stroke or Hemorrhagic Stroke: An Updated Meta-Analysis. *Front Neurol.* 2019;**10**:1032. [PubMed ID: 31636598]. [PubMed Central ID: PMC6787274]. <https://doi.org/10.3389/fneur.2019.01032>.
 10. Kirdar NN, Şenol H. [The Frequency of Neuropathic Pain in Stroke Patients and Medical Treatment Approaches]. *Hipokrat Tıp Dergisi.* 2022;**2**(2):24-8. Turkish. <https://doi.org/10.29228/HMj.16>.
 11. Salsano G, Pracucci G, Mavilio N, Saia V, Bandettini di Poggio M, Malfatto L, et al. Complications of mechanical thrombectomy for acute ischemic stroke: Incidence, risk factors, and clinical relevance in the Italian Registry of Endovascular Treatment in acute stroke. *Int J Stroke.* 2021;**16**(7):818-27. [PubMed ID: 33283685]. <https://doi.org/10.1177/1747493020976681>.
 12. Nikbakht H, Shojaiie L, Niknejad N, Hassani-pour S, Soleimanpour H, Heidari S, et al. Mortality Rate of Acute Stroke in Iran: A Systematic Review and Meta-Analysis. *Casp J Environ Sci.* 2022;**8**(4):252-67. <https://doi.org/10.32598/cjns.4.31.338.1>.
 13. Krishnamurthi RV, Moran AE, Feigin VL, Barker-Collo S, Norrving B, Mensah GA, et al. Stroke Prevalence, Mortality and Disability-Adjusted Life Years in Adults Aged 20-64 Years in 1990-2013: Data from the Global Burden of Disease 2013 Study. *Neuroepidemiology.* 2015;**45**(3):190-202. [PubMed ID: 26505983]. <https://doi.org/10.1159/000441098>.
 14. Caro CC, Costa JD, Da Cruz DMC. Burden and Quality of Life of Family Caregivers of Stroke Patients. *Occup Ther Health Care.* 2018;**32**(2):154-71. [PubMed ID: 29578827]. <https://doi.org/10.1080/07380577.2018.1449046>.
 15. Simmonds KP, Atem F, Welch BG, Ifejika NL. Abstract 146: Racial Disparities In The Treatment Of Post-stroke Complications Among Acute Stroke Patients. *Stroke.* 2023;**54**(Suppl_1). https://doi.org/10.1161/str.54.suppl_1.146.
 16. Hatefi M, Komlakh K, Nouri L. Investigating the effect of methylprednisolone pulse on the treatment of back pain. *Rom J Mil Med.* 2022;**125**(2):264-8. <https://doi.org/10.22034/ecc.2022.323514.1293>.
 17. Hatefi M, Tarjoman A, Borji M. Do Religious Coping and Attachment to God Affect Perceived Pain? Study of the Elderly with Chronic Back Pain in Iran. *J Relig Health.* 2019;**58**(2):465-75. [PubMed ID: 30610514]. <https://doi.org/10.1007/s10943-018-00756-9>.
 18. Vasigh A, Tarjoman A, Borji M. Relationship Between Spiritual Health and Pain Self-Efficacy in patients with Chronic Pain: A Cross-Sectional Study in West of Iran. *J Relig Health.* 2020;**59**(2):115-25. [PubMed ID: 31087227]. <https://doi.org/10.1007/s10943-019-00833-7>.
 19. Broussy S, Saillour-Glenisson F, Garcia-Lorenzo B, Rouanet F, Lesaine E, Maugeais M, et al. Sequelae and Quality of Life in Patients Living at Home 1 Year After a Stroke Managed in Stroke Units. *Front Neurol.* 2019;**10**:907. [PubMed ID: 31496987]. [PubMed Central ID: PMC6712081]. <https://doi.org/10.3389/fneur.2019.00907>.
 20. Mendigutia-Gomez A, Quintana-Garcia MT, Martin-Sevilla M, de Lorenzo-Barrientos D, Rodriguez-Jimenez J, Fernandez-de-Las-Penas C, et al. Post-needleling soreness and trigger point dry needling for hemiplegic shoulder pain following stroke. *Acupunct Med.* 2020;**38**(3):150-7. [PubMed ID: 32009422]. <https://doi.org/10.1177/0964528419882941>.
 21. Almhdawi KA, Alazrai A, Kanaan S, Shyyab AA, Oteir AO, Mansour ZM, et al. Post-stroke depression, anxiety, and stress symptoms and their associated factors: A cross-sectional study. *Neuropsychol Rehabil.* 2021;**31**(7):1091-104. [PubMed ID: 32419606]. <https://doi.org/10.1080/09602011.2020.1760893>.
 22. Sasaki K, Halder SK, Matsunaga H, Ueda H. Beneficial actions of prothymosin alpha-mimetic hexapeptide on central post-stroke pain, reduced social activity, learning-deficit and depression following cerebral ischemia in mice. *Peptides.* 2020;**126**:170265. [PubMed ID: 31982448]. <https://doi.org/10.1016/j.peptides.2020.170265>.
 23. Wu X, Liang Y, Zheng B, Wang H, Ning M, Zheng H, et al. Care stress in caregivers of disabled stroke patients: a cross-sectional survey. *Ann Palliat Med.* 2020;**9**(4):2211-20. [PubMed ID: 32692235]. <https://doi.org/10.21037/apm-20-1369>.
 24. Valencia-Chulian R, Heredia-Rizo AM, Moral-Munoz JA, Lucena-Anton D, Luque-Moreno C. Dry needling for the management of spasticity, pain, and range of movement in adults after stroke: A systematic review. *Complement Ther Med.* 2020;**52**:102515. [PubMed ID: 32951759]. <https://doi.org/10.1016/j.ctim.2020.102515>.
 25. Ghahremani F, Dadgoo M, Noorizadeh S, Mazdeh M, Pourahmadi MR. [The relationship between shoulder pain with shoulder neuromusculoskeletal disorders in post stroke hemiparesis]. *Mod Rehabil.* 2014;**8**(4). Persian.
 26. Mousavi SJ, Parnianpour M, Abedi M, Askary-Ashtiani A, Karimi A, Khorsandi A, et al. Cultural adaptation and validation of the Persian version of the Disabilities of the Arm, Shoulder and Hand (DASH) outcome measure. *Clin Rehabil.* 2008;**22**(8):749-57. [PubMed ID: 18678575]. <https://doi.org/10.1177/0269215508085821>.
 27. English LD. STEM education K-12: perspectives on integration. *Int J STEM Educ.* 2016;**3**:1-8. <https://doi.org/10.1186/s40594-016-0036-1>.
 28. Adhami-Moghadam F, Iran-Pour E. Psychological disorders in patients with retinitis pigmentosa in iran. *Iran J Public Health.* 2014;**43**(4):523-8. [PubMed ID: 26005663]. [PubMed Central ID: PMC4433734].
 29. Mazlomezhad A, Moghadam FA. Evaluation of quality of life and self-efficacy in adolescents with amblyopia. *J Med Life.* 2022;**15**(4):499-503. [PubMed ID: 35646181]. [PubMed Central ID: PMC9126460]. <https://doi.org/10.25122/jml-2020-0035>.
 30. Karimiyarandi H, Karbasfrushan A. Effect of Duloxetine on Pain Relief in Patients Undergoing Arthroplasty Surgery: A Systematic Review. *Gomal J Med Sci.* 2022;**20**(4):213-20. <https://doi.org/10.46903/gjms/20.04.1191>.
 31. Haslam BS, Butler DS, Kim AS, Carey LM. Somatosensory Impairment and Chronic Pain Following Stroke: An Observational Study. *Int J Environ Res Public Health.* 2023;**20**(2). [PubMed ID: 36673661]. [PubMed Central ID: PMC9859194]. <https://doi.org/10.3390/ijerph20020906>.
 32. Hassan AB, Danazumi MS, Abdullahi A, Yakasai AM. Effect of transcranial direct current stimulation (tDCS) delivered via dorsolateral prefrontal cortex on central post-stroke pain and depression: a case report. *Physiother Theory Pract.* 2022;**38**(11):1799-806. [PubMed ID: 33648417]. <https://doi.org/10.1080/09593985.2021.1891591>.
 33. Diener HC, Sacco RL, Easton JD, Granger CB, Bernstein RA, Uchiyama S, et al. Dabigatran for Prevention of Stroke after Embolic Stroke of Undetermined Source. *N Engl J Med.* 2019;**380**(20):1906-17. [PubMed ID: 31091372]. <https://doi.org/10.1056/NEJMoa1813959>.
 34. Hart RG, Sharma M, Mundl H, Kasner SE, Bangdiwala SI, Berkowitz SD, et al. Rivaroxaban for Stroke Prevention after Embolic Stroke of Undetermined Source. *N Engl J Med.* 2018;**378**(23):2191-201. [PubMed ID: 29766772]. <https://doi.org/10.1056/NEJMoa1802686>.
 35. Stiekema APM, Nijse B, de Kort PLM, Spikman JM, Visser-Meily

- JMA, van Heugten CM. The relationship between social cognition and participation in the long term after stroke. *Neuropsychol Rehabil.* 2021;**31**(2):278–92. [PubMed ID: 31854264]. <https://doi.org/10.1080/09602011.2019.1692670>.
36. Al-Gibbawi M, Ayinde HO, Bhatia NK, El-Chami MF, Westerman SB, Leon AR, et al. Relationship between device-detected burden and duration of atrial fibrillation and risk of ischemic stroke. *Heart Rhythm.* 2021;**18**(3):338–46. [PubMed ID: 33250442]. <https://doi.org/10.1016/j.hrthm.2020.10.017>.
37. Xing L, Jing L, Tian Y, Liu S, Lin M, Du Z, et al. High prevalence of stroke and uncontrolled associated risk factors are major public health challenges in rural northeast China: A population-based study. *Int J Stroke.* 2020;**15**(4):399–411. [PubMed ID: 31092151]. <https://doi.org/10.1177/1747493019851280>.
38. Regenhardt RW, Biseko MR, Shayo AF, Mmbando TN, Grundy SJ, Xu A, et al. Opportunities for intervention: stroke treatments, disability and mortality in urban Tanzania. *Int J Qual Health Care.* 2019;**31**(5):385–92. [PubMed ID: 30165650]. [PubMed Central ID: PMC6528701]. <https://doi.org/10.1093/intqhc/mzy188>.
39. Tu WJ, Chao BH, Ma L, Yan F, Cao L, Qiu H, et al. Case-fatality, disability and recurrence rates after first-ever stroke: A study from bigdata observatory platform for stroke of China. *Brain Res Bull.* 2021;**175**:130–5. [PubMed ID: 34329730]. <https://doi.org/10.1016/j.brainresbull.2021.07.020>.
40. Cucchiara B, Elm J, Easton JD, Coutts SB, Willey JZ, Biros MH, et al. Disability After Minor Stroke and Transient Ischemic Attack in the POINT Trial. *Stroke.* 2020;**51**(3):792–9. [PubMed ID: 32078486]. [PubMed Central ID: PMC7313376]. <https://doi.org/10.1161/STROKEAHA.119.027465>.
41. Kelly PJ, Furie KL, Shafiqat S, Rallis N, Chang Y, Stein J. Functional recovery following rehabilitation after hemorrhagic and ischemic stroke. *Arch Phys Med Rehabil.* 2003;**84**(7):968–72. [PubMed ID: 12881818]. [https://doi.org/10.1016/s0003-9993\(03\)00040-6](https://doi.org/10.1016/s0003-9993(03)00040-6).
42. Alamam DM, Leaver A, Alsobayel HI, Moloney N, Lin J, Mackey MG. Low Back Pain-Related Disability Is Associated with Pain-Related Beliefs Across Divergent Non-English-Speaking Populations: Systematic Review and Meta-Analysis. *Pain Med.* 2021;**22**(12):2974–89. [PubMed ID: 33624814]. <https://doi.org/10.1093/pm/pnaa430>.