



Musculoskeletal Evaluation of Preschool Children Referring to the School Screening Center

Mohaddeseh Azadvari ¹, Seyede Zahra Emami Razavi ^{1,*} and Fatemeh Behyad²

¹Physical Medicine & Rehabilitation Department, Imam Khomeini Hospital Complex, Tehran University of Medical Sciences, Tehran, Iran

²Faculty of Medicine, Tehran University of Medical Sciences, Tehran, Iran

*Corresponding author: Physical Medicine & Rehabilitation Department, Imam Khomeini Hospital Complex, Tehran University of Medical Sciences, Tehran, Iran. Email: zemamirazavi@gmail.com

Received 2021 December 20; Revised 2022 March 20; Accepted 2022 April 05.

Abstract

Background: One of the biggest global challenges of the 21st century is the increasing burden of non-communicable (chronic) diseases.

Objectives: Considering the high prevalence of systemic and musculoskeletal diseases and the need for further epidemiological studies to provide health programs, we conducted this research.

Methods: In this cross-sectional study, a total of 344 primary school children referred to the East Tehran Health Center and examined during a certain period of the autumn of 2019 were enrolled. Demographic information and history of systemic diseases were collected using a questionnaire. Height and weight measurements and feet and spine examinations were performed by the researcher.

Results: In summary, 50.3% were girls, and 49.7% were boys. In terms of body mass index (BMI), 10.2% were underweight, 70.3% were normal, 10.8% were overweight, and 8.7% were obese. The highest prevalence of diseases was obesity (8.7%), followed by musculoskeletal diseases (8.1%), which included only the flatfoot.

Conclusions: Screening programs are recommended from an early age to prevent diseases. To reduce the rate of musculoskeletal disorders and obesity in children, identifying and following them in the early stages of the disorders is recommended for early diagnosis and appropriate treatment.

Keywords: Musculoskeletal Diseases, Obesity, Preschool Child

1. Background

The increasing burden of non-communicable (chronic) diseases is one of the biggest global challenges of the 21st century (1). About two-thirds of deaths globally occur due to such diseases, more than 80% of which occur in low- and middle-income countries (2).

Obesity is the result of a long-term imbalance between energy intake and output in the body, during which the energy entering the body is greater than the energy consumed. In addition to long-term medical complications, childhood obesity may cause psychosocial problems, such as low self-esteem, depression, and poor quality of life (3).

One of the concerns of the parents of obese children is "micropenis," which should be considered (4, 5).

Obese children are at higher risk for other chronic conditions such as asthma, sleep apnea, bone, and joint problems, type 2 diabetes, and risk factors for cardiovascular diseases. Generally, a child is more likely to be obese if

he/she has bigger arms, thigh, and abdomen, becomes short of breath after the slightest physical activity, or is unable to physically compete with other children of the same age (3).

Musculoskeletal disorders are very common, and the risk of their development increases with age. Early diagnosis of musculoskeletal disorders is the most important factor in treating and preventing further damage to the body (5).

A review of the literature shows the high prevalence of postural abnormalities in Iran, which is higher among middle- and high-school students due to the lack of adequate knowledge and training, as well as a sedentary lifestyle. However, this issue has been less evaluated in children (6).

Flatfeet are one of the most common musculoskeletal disorders, in which the longitudinal arch of the foot is reduced, and in severe cases, the inner side of the sole comes into contact with the ground. Flatfeet are generally divided

into 2 types: (1) flexible flatfoot, and (2) rigid flatfoot (6).

Obviously, the earlier a diagnosis and therapeutic measures take place, the more short- and long-term benefits they will have. To manage this group of disorders, it is necessary to study its prevalence among children to estimate their current level of health, discover the critical points of disease in childhood, and provide an opportunity to estimate appropriate preventive and therapeutic measures at the beginning of life.

Some studies have examined these complications. For example, Karimian et al in a descriptive-correlational study on 148 children in Fasa City showed that non-standard facilities and equipment, such as backpacks, school desks, and the lack of specialists in prevention and treatment, have led to the prevalence of these disorders (7).

Flatfeet were present in 54% of 3-year-old and 26% of 6-year-old children. Boys were more likely to be flatfooted than girls (52% vs 36%). Obesity was associated with flatfeet. Flatfeet were present in 62% of obese, 51% of overweight, and 42% of young children with normal body weight (7).

Ogden et al mentioned that 8.4% of preschool children (age, 2 - 5 years), 17.7% of school-aged children (age, 6 - 11 years), and 20.5% of adolescents (age, 12 - 19 years) have obesity (8).

In a cross-sectional study conducted by Zakeri et al (2014) in Abadan, Iran, the prevalence of musculoskeletal disorders in primary school students was evaluated, in which 383 primary school students were randomly selected and studied. The most prevalent skeletal disorder observed in both females and males was scoliosis (85.4%), followed by drooping shoulders (81.7%). Flat back (1.6%) was the less common skeletal disorder. Also, head forward, genu valgus, and hallux valgus were significantly higher in girls than boys. The prevalence of flatfeet in the total population was 22.5% (9).

In the cross-sectional study by Gaeini et al, the prevalence of underweight, overweight, and obesity in preschool children of Tehran in 2009 - 2010 was studied. A total of 756 children aged 3 - 6 years were randomly selected from 5 different geographical regions of Tehran, and the findings showed that the prevalence of underweight, overweight, and obesity was 4.77%, 9.81%, and 4.77% in boys and 4.77%, 10.31%, and 4.49% in girls, respectively (10).

2. Objectives

Due to the importance of this subject for further prevention and comprehensive evaluation of musculoskeletal disorders, we conducted this study.

3. Methods

In this cross-sectional study, a total of 344 preschool children referred to the East Tehran Health Screening Center in 2020 were selected by a simple sampling method.

Inclusion criteria were all children between the ages of 5 to 7 referred for screening.

There were no exclusion criteria.

In this study, all preschool children referred to East Tehran Health Center were examined during a certain period of time in the fall of 2020. This descriptive cross-sectional study was performed for the purpose of screening.

A researcher-made questionnaire was used for data collection. This questionnaire contains 3 separate sections as follows: (1) demographic information, including age, gender, height, and weight, which was measured by a physician; (2) data on musculoskeletal disorders, including a question with scores of 0 for "no history of the disorder" and 1 for "a history of the disorder."; (3) systemic disease information, including 12 questions with scores of 0 for "no history of the disease" and 1 for "a history of the disease."

Evaluation of head forward, drooping shoulders, flat back, scoliosis, thoracic kyphosis, lordosis, genu valgus, genu varus, flatfoot, and hallux valgus was performed using instruments such as plumb line, checkerboard, mirror box, and questionnaire of personal characteristics. Examinations of the sole for flatfeet and the spine for scoliosis and kyphosis were performed in the presence of parents; a physician measured and recorded the weight and height of children, as well as answered the questions about the musculoskeletal disorders. The rest of the health questionnaire, including demographic information and systemic disease records, was filled out by the physician based on parental statements.

To describe the qualitative variables, the data obtained to answer the research questions and hypotheses were analyzed using SPSS (SPSS Inc, Chicago, Ill, USA) to calculate the frequency and percentage. To show the relationship between the prevalence of diseases and demographic variables, the chi-square or t test was used.

3.1. Ethical Consideration

The Ethics Committee of Tehran University of Medical Sciences approved this study (IR.TUMS.IKHC.REC.1398.174).

4. Results

The frequency of variables and their relationship with demographic information are as follows:

Gender: Of the 344 patients, 173 (50.3%) were female, and 171 (49.7%) were male.

The frequency and percentage distribution of the age of the patients are shown in [Table 1](#).

Table 1. Frequency and Percentage Distribution of Age

Age	No. (%)
5	68 (19.8)
6	263 (76.5)
7	12 (3.5)
8	1 (0.2)
Total	344 (100)

Body mass index (BMI): Of the 344 patients, 35 (10.2%) were underweight (BMI < 5%), 242 (70.3%) were normal (BMI between 5% and 85%), 37 (10.8%) were overweight (BMI between 85% and 95%), and 30 (8.7%) were obese (BMI above 95%; see [Table 2](#)).

Table 2. Frequency of Body Mass Index

BMI	No. (%)
Underweight	35 (10.2)
Normal	242 (70.3)
Overweight	37 (37)
Obese	30 (30)
Total	344 (344)

There was no significant relationship between age and BMI groupings ($P = 0.677$); thus, 16.2% of 5-year-old children were underweight, 10.3% were overweight, and 8.8% were obese; 9.1% of 6-year-old children were underweight, 11% were overweight, and 9.1% were obese; further, 8.3% of 7-year-old children were overweight.

There was no significant relationship between gender and BMI groupings ($P = 0.584$); thus, 11.6%, 11%, and 6.9% of girls and also 8.8%, 10.5%, and 10.5% of boys were underweight, overweight, and obese, respectively.

Of the 344 patients, 28 (8.1%) had musculoskeletal disorders, all of whom had flatfeet.

Regarding the relationship between musculoskeletal disorders and BMI, the mean BMI in those with and without musculoskeletal disorders was reported to be 16.11 ± 2.6 and 15.94 ± 2.44 , respectively; thus, no significant relationship was observed between BMI and the rate of musculoskeletal disorders ($P = 0.716$).

5. Discussion

A group of known social, environmental, behavioral, and underlying (hereditary) risk factors are observed in

these diseases, increasing the likelihood of disease or damage. If modifiable risk factors are eliminated or controlled, a percentage of such disorders and disabilities can be prevented.

In the study by Zakeri et al, the most prevalent skeletal disorder observed in both females and males was scoliosis (85.4%), followed by drooping shoulders (81.7%). Flat back (1.6%) was the less common skeletal disorder. Also, head forward, genu valgus, and hallux valgus were significantly higher in girls than boys. The prevalence of flatfoot in the total population was 22.5% (9).

In the study by Askary et al, the prevalence of flatfoot was evaluated and compared between the genders among primary school children. The overall prevalence of flatfoot was 74%, out of which 23% were mild, 34% were moderate, and 17% were severe. The prevalence of flatfoot in girl and boy students was 75.2% and 72.6%, respectively (11). In our study, the prevalence of musculoskeletal disorders, which only included flatfoot, was 8.1% and had no significant relationship with age, gender, and BMI.

In a cross-sectional study, Larrañaga et al estimated the prevalence of overweight and obesity. In this study, 55.4% of the population was obese, which included 6% of boys and 4.7% of girls, and the highest obesity rates were seen in the 11 to 14-year-old group of boys (7.2%) and the 4 to 6-year-old group of girls (12.5%). The prevalence of obesity was higher in the less privileged socioeconomic strata (6.9% vs. 5.2%) for both boys and girls (12).

In the cross-sectional study by Gaeini et al., the prevalence of underweight, overweight, and obesity in preschool children of Tehran in 2009-2010 was 4.77%, 9.81%, and 4.77% in boys and 4.77%, 10.31%, and 4.49% in girls, respectively (10).

In our study, 10.2% were underweight (BMI < 5%), 70.3% were normal (BMI between 5% and 85%), 10.8% were overweight (BMI between 85% and 95%), and 8.7% were obese (BMI above 95%).

A total of 19.5% suffered from overweight and obesity, and the importance of addressing this health issue is obvious (9).

Due to the high prevalence of musculoskeletal disorders (especially flatfoot) and obesity in preschool children, screening programs with more detailed examinations are recommended for the prevention and early diagnosis of this disorder (10). In order to reduce the rate of musculoskeletal disorders and obesity in children, identifying and following them in the early stages of the disorders seem to be a necessary solution for early diagnosis, appropriate treatment, and recovery of children. Also, more detailed investigations, such as determining the degree of flatfoot (mild, moderate, and severe) and the resulting physical problems, as well as factors affecting childhood

obesity, are recommended for future studies.

5.1. Conclusions

In this study, 344 preschool children referred to the East Tehran Health Center in 2020 were examined for the prevalence of musculoskeletal disorders.

In terms of BMI, 10.2% were underweight, 70.3% were normal, 10.8% were overweight, and 8.7% were obese. In this study, the most prevalent disorder was obesity (8.7%), followed by musculoskeletal disorders, which only involved flatfoot (8.1%).

Acknowledgments

We thank the personnel of the Physical Medicine and Rehabilitation Clinic in Imam Khomeini Hospital, Tehran, Iran, who helped us in performing this study.

Footnotes

Authors' Contribution: Study concept and design, M. A. and S. Z. E. R.; Acquisition of data, M. A. and F. B.; Analysis and interpretation of data, M. A. and S. Z. E. R.; Drafting of the manuscript, S. Z. E. R.; Critical revision of the manuscript for important intellectual content, M. A. and S. Z. E. R.; Statistical analysis, M. A. and S. Z. E. R.; Administrative, technical, and material support, S. Z. E. R. and F. B.; Study supervision, M. A. and S. Z. E. R.

Conflict of Interests: The authors declare no conflict of interests.

Data Reproducibility: The dataset presented in the study is available on request from the corresponding author during submission or after its publication.

Ethical Approval: The Ethics Committee of Tehran University of Medical Sciences approved this study (IR.TUMS.IKHC.REC.1398.174) (ethics.research.ac.ir/EthicsProposalViewEn.php?id = 87999).

Funding/Support: This study was supported by Tehran University of Medical Sciences without any financial support.

Informed Consent: All parents of the participant signed informed consent.

References

1. Karimi S, Javadi M, Jafarzadeh F. Economic burden and costs of chronic diseases in Iran and the world. *Health Inf Manag.* 2012;**8**(7):984-96.
2. Nolte E, McKee M. *Caring for people with chronic conditions – a health systems perspective.* 71. UK: McGraw-Hill Education; 2008. doi: [10.1055/s-0029-1239177](https://doi.org/10.1055/s-0029-1239177).
3. Ghergherechi R, Hazhir N, Samadi M, Rafeey M. Cardiovascular risk factors in obese children and adolescents. *Med J Tabriz Univ Med Sci Health Serv.* 2018;**40**(1):52-8.
4. Rezakhaniha S, Rezakhaniha B, Siroosbakht S. Which of the Anthropometric Parameters and Hormonal Variables in Prepubertal Children Are Correlated to True Micropenis? *Iran J Pediatr.* 2021;**31**(1). e108534. doi:[10.5812/ijp.108534](https://doi.org/10.5812/ijp.108534).
5. Rezakhaniha S, Rezakhaniha B, Aarabi N, Siroosbakht S. Is It Necessary to Weight Loss in Obese Boys with Small Penile Length? A Case-Control Study. *J Compr Pediatr.* 2020;**11**(4). doi: [10.5812/comprped.107272](https://doi.org/10.5812/comprped.107272).
6. Cherney K. *Musculoskeletal Disorders.* New York, USA: Healthline Media; 2018, [cited 2022]. Available from: <https://www.healthline.com/health/fibromyalgia#diet>.
7. Karimian R, Karimian M, Hadipour M, Heyat F, Janbozorgi A. [The prevalence of children's postural abnormalities and its association with sport activity]. *Journal of Fasa University of Medical Sciences.* 2016;**6**(1):106-12. Persian.
8. Ogden CL, Carroll MD, Kit BK, Flegal KM. Prevalence of childhood and adult obesity in the United States, 2011-2012. *JAMA.* 2014;**311**(8):806-14. doi: [10.1001/jama.2014.732](https://doi.org/10.1001/jama.2014.732). [PubMed: [24570244](https://pubmed.ncbi.nlm.nih.gov/24570244/)]. [PubMed Central: [PMC4770258](https://pubmed.ncbi.nlm.nih.gov/PMC4770258/)].
9. Zakeri Y, Gheibizadeh M, Baraz S, Bijan Nejad D, Latifi SM. Relationship between features of desks and chairs and prevalence of skeletal disorders in primary school students in Abadan. *Int J Pediatr.* 2016;**4**(11):3949-56.
10. Gaeini A, Kashaf M, Samadi A, Fallahi A. Prevalence of underweight, overweight and obesity in preschool children of Tehran, Iran. *J Res Med Sci.* 2011;**16**(6):821-7. [PubMed: [22091313](https://pubmed.ncbi.nlm.nih.gov/22091313/)]. [PubMed Central: [PMC3214402](https://pubmed.ncbi.nlm.nih.gov/PMC3214402/)].
11. Askary Kachoosangy R, Aliabadi F, Ghorbani M. Prevalence of flat foot: comparison between male and female primary school students. *Iran Rehabil J.* 2013;**11**(3):22-4.
12. Larranaga N, Amiano P, Arrizabalaga JJ, Bidaurrezaga J, Gorostiza E. Prevalence of obesity in 4-18-year-old population in the Basque Country, Spain. *Obes Rev.* 2007;**8**(4):281-7. doi: [10.1111/j.1467-789X.2006.00306.x](https://doi.org/10.1111/j.1467-789X.2006.00306.x). [PubMed: [17578378](https://pubmed.ncbi.nlm.nih.gov/17578378/)].