**Research Article** 



# Identification Based on Height, Age, and Gender Estimation Using the Measured Parameters of the T10 and T11 Vertebrae in Three-Dimensional Computed Tomography Scan Among the Iranian Population

Mohammad Reza Khaleghi<sup>1</sup>, Maryam Vasheghani Farahani <sup>(D) 2</sup>, Azadeh Memarian <sup>3</sup>, Nazanin Maleki<sup>4</sup>, Dariush Taherkhani <sup>5</sup>, Babak Shekarchi<sup>1,\*</sup>

<sup>1</sup> Radiation Sciences Research Center, AJA University of Medical Sciences, Tehran, Iran

<sup>2</sup> Department of Forensic Medicine, School of Medicine, AJA University of Medical Sciences, Tehran, Iran

<sup>3</sup> Department of Forensic Medicine and Toxicology, Mazandaran University of Medical Sciences, Sari, Iran

<sup>4</sup> Department of Radiology, School of Medicine, Iran University of Medical Sciences, Tehran, Iran

<sup>5</sup> Shahed University of Medical Sciences, Tehran, Iran

Corresponding author: Radiation Sciences Research Center, AJA University of Medical Sciences, Tehran, Iran. Email: shekarchi.babak@yahoo.com

Received 2023 May 24; Revised 2023 July 25; Accepted 2023 July 25.

## Abstract

The present study aimed to estimate the age, gender, and height of the Iranian population based on the characteristics of the 10th and 11th thoracic vertebrae for identification. This cross-sectional study measured the T10 and T11 indices in axial, sagittal, and coronal computed tomography (CT) scan images. The measured indices included vertebral length (VL), endplate depth (EPD), endplate width (EPW), foramen diameter (FD), foramen width (FW), transverse process distance (TDm), spinal process height (SPH), spinal process length (SPL), vertebral body height (VBH), pedicle height (PH), pedicle width (PW), articular process height superior (APHs), and maximum distance between articular processes (ADm). The mean differences of the T10 and T11 indices between the two groups of men and women, except FW, FD, and APHi, were higher in men, which were statistically significant (P < 0.006). VL, EPD, EPW, FW, TDm, VBH, PH, and ADm in men, and VL, EPD, EPW, TDm, SPH, SPL, PH, APHs, and ADm in women were significantly correlated with height in the T10 vertebra (P < 0.05). Moreover, FD in T10 was found to be negatively correlated with women's height (P = 0.04). In addition, in the T11, the VL, EPD, EPW, FW, FD, TDm, and VBH indices in men and the VL, EPW, TDm, SPL, PH, APHs, and PW in women were significantly correlated with height (P < 0.05). Moreover, VL, EPD, TDm, SPH, SPL, and APHi in women and VL in men were significantly and positively correlated with age in T10 (P < 0.05), similar to the T11 APHs correlation with age in men and VL, EPW, EPD, SPL, and APHi in women. The current study showed that the T10 and L11 vertebral indices were significantly correlated with gender, height, and age.

Keywords: Forensic Anthropology, Computed Tomography Scan, Spine, Identification, Population

#### 1. Background

Identification is known as a chief mission of the forensic practice. Determination of biological factors such as gender, age, and height reduces the probable victim matches in forensic search procedures and offers several helpful clues in an individual's identification (1).

Various bones, including the pelvis, skull, humerus, and talus, have been metrically studied to evaluate the

subjects' biological factors (2-4). Occasionally, due to mass disasters or environmental conditions, bones are broken or commingled; in these situations, identification will be carried out based on small components of the skeleton (5). Therefore, it is essential to develop metric approaches related to different types of bones.

The vertebral column is the greatest conserved bone among all skeletal components (6). It is also used in

Copyright © 2023, Annals of Military and Health Sciences Research. This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (http://creativecommons.org/licenses/by-nc/4.0/) which permits copy and redistribute the material just in noncommercial usages, provided the original work is properly cited.

forensic assessments due to its comparative thickness and capability to resist heaviness. Correspondingly, the vertebral column contains an outer dense cortical bone and an inner cancellous bone, which help bear weight by vertebrae (7-9).

A limited number of previous studies have evaluated the 12th thoracic vertebra in gender, age, and height estimations (7, 9-11), and only one study has examined the 10th thoracic vertebra in gender determination but not in age and height (12). Moreover, no study was found to measure the 11th thoracic vertebra and its correlations with gender, age, and height.

Diagnosing and identifying differences in spinal dimension among adults could be very helpful in determining height, age, and gender in forensic medicine. In this regard, each one of the components of the spinal cord at any stage can greatly help identify the person. Accordingly, estimating age, gender, and height based on the characteristics of the thoracic vertebrae is practical.

# 2. Objectives

In the present study, we aimed to investigate the height, age, and gender and finally to determine the identity of the military personnel by examining various parameters in a three-dimensional (3D) computed tomography (CT) scan in the T10 and T11 vertebrae.

#### 3. Methods

This cross-sectional study was conducted after obtaining the approval of the AJA University of Medical Sciences Ethics Committee in 2020.

#### 3.1. Study Population

In this cross-sectional study, 100 patients over 16 years of age were investigated, and for any reason, CT scans of the abdomen and pelvis were performed for them in Imam Reza Hospital.

The exclusion criteria included being under 16 years old, a history of congenital anomalies of the thoracic spine, a history of thoracic trauma with any severity, structural, metabolic, or rheumatic disorder in the thoracic spine, and a history of any surgical intervention and reconstructive therapy in the thoracic region. Badr-al-Dine et al. defined the sample size in their study as 100 (7).

#### 3.2. Computed Tomography Scan Protocol

The Philips Ingenuity Flex multidetector scanner was applied. Abdominopelvic CT without any contrast was utilized. Image acquisition parameters were a tube voltage of 120 kVp, a tube current of 150 mAs, slice thickness of 2 mm at a -1 mm increment, collimation of 16\*1.5, pitch of 1.188, and rotation time of 0.5 sec. All images were imported to the local picture archiving and communicative system (PACS). Then, the images were reconstructed in the sagittal and coronal axes in the local PACS. After that, an expert radiologist measured the indices in the CT images (including coronal, sagittal, and axial images) in the bone window (width: 2000 HU, level: 250 HU) reconstructed images.

## 3.3. Measurements and Indices

All CT scan images were reviewed by a radiologist, and the following indices were then measured. If there was a suspicion regarding any measurement error, the images were reviewed by a musculoskeletal expert attending radiologist, and his opinion was applied as a criterion.

We measured the indices of the 10th and 11th thoracic vertebrae. The measured indices were as follows:

Vertebral length (VL): The maximum distance between two parallel lines, one passing through the anterior of the vertebral body and the other passing through the apex of the spinous process in sagittal images.

Upper endplate depth (EPDu): The maximum anterior-posterior distance of the upper surface of the vertebral body in axial images.

Upper endplate width (EPWu): The maximum lateral distance of the upper surface of the vertebral body in axial images.

Foramen diameter (FD): The maximum anteriorposterior distance of vertebral foramen in axial images.

Foramen width (FW): The maximum lateral distance of the vertebral body in axial images.

Transverse process distance (TDm): The maximum distance between transverse processes in axial images.

Spinal process height (SPH): The maximum distance between the superior-inferior aspects of the spinal process in sagittal images.

Spinal process length (SPL): The maximum distance between the anterior-posterior aspects of the spinal process in axial images.

Verbal body height anterior (VBHa): Anterior height of the vertebral body in sagittal images.

Pedicle height (PH): Pedicle height in sagittal images. Pedicle width (PW): Pedicle width in axial images.



Figure 1. Measured dimensions of the thoracic vertebrae in coronal, sagittal, and axial views: Maximum distance between the articular process (ADm), inferior articular process height (APHi), superior articular process height (APHs), upper endplate depth (EPDu), upper endplate width (EPWu), foramen diameter depth (FDd), foramen diameter width (FDw), pedicle height (PH), pedicle widths (PW), transverse process distance (TDm), spinous process length (SPL), spinous process height (SPH), vertebral body height anterior (VBHa), vertebral length (VL).

Articular process height inferior (APHi): Inferior articular process height in sagittal images.

Articular process height superior (APHs): Superior articular process height in sagittal images.

Maximum distance between articular processes (ADm): The maximum distance between articular processes in the coronal image.

The measured indices are shown in Figure 1.

Finally, the evaluated indices were statistically compared, and the diagnostic accuracy of each one of them was evaluated to estimate height, age, and gender.

# 3.4. Statistical Analysis

The IBM<sup>®</sup> SPSS<sup>®</sup> software version 23 was used for data analysis. Quantitative variables were described as mean and standard deviation, and stratified qualitative variables as percentages. The *t*-test or analysis of variance (ANOVA) was used for quantitative comparison, and the chi-square test was used for comparing qualitative variables. The significance level was considered a p-value less than 0.05.

# 4. Results

The frequency distributions of age and height of the subjects by gender are shown in Table 1.

The results of the two independent samples *t*-test showed that the mean age difference between the two groups, including men and women, was not statistically significant at the 95% confidence level (P > 0.001). However, the mean height difference between male and female patients was statistically significant at a 95% confidence level (P < 0.0001).

The results of the two independent samples *t*-test regarding normal data and the Mann-Whitney test regarding non-normal data showed that the mean difference of all T10 indices, except FW, FD, and APHi, was statistically significant at a 95% confidence level between male and female patients (Table 2).

Table 1. Frequency Distribution of Age and Height of the Subjects by Gender							
	Ν	Mean $\pm$ SD	Min	Max	P-Value		
Age					0.1008		
Men	54	$35.04 \pm 10.11$	17	56			
Women	46	$38.28\pm9.333$	16	55			
Height					< 0.0001		
Men	54	$176.1 \pm 7.480$	160	195			
Women	46	$161.6\pm6.053$	152	179			

f <b>able 2.</b> Compari	son of the T10 and T11 Indices i	n Men and Women						
Ter ditana		T10		Tu				
mulces	Men (n = 54)	Women (n = 46)	P-Value	Men (n = 54)	Women (n = 46)	P-Value		
VL	70.89	63.55	< 0.0001	71.89	65.03	< 0.0001		
EPD	28.99	25.90	< 0.0001	29.93	26.55	< 0.0001		
EPW	35.21	30.75	< 0.0001	38.51	33.47	< 0.0001		
FD	14.60	14.34	0.2794	15.39	15.28	0.7155		
FW	17.60	17.33	0.4479	18.89	18.81	0.8694		
TDm	59.49	53.11	< 0.0001	55.61	49.05	< 0.0001		
SPH	10.36	9.654	0.0116	11.18	9.978	0.0009		
SPL	20.52	17.30	< 0.0001	20.94	17.78	< 0.0001		
VBH	22.65	21.01	< 0.0001	23.09	21.75	0.0003		
РН	12.28	11.56	0.0056	13.36	12.14	< 0.0001		
PW	6.496	5.389	< 0.0001	7.954	6.702	< 0.0001		
APHi	18.42	17.72	0.2128	20.03	19.08	0.1007		
APHs	13.62	11.41	0.0008	14.24	12.15	0.0035		
ADm	29.19	26.37	0.0005	30.10	27.29	0.0009		

Abbreviations: VL, vertebral length; EPD, endplate depth; EPW, endplate width; FD, foramen diameter; FW, foramen width; TDm, transverse process distance; SPH, spinous process height; SPL, spinous process length; VBH, vertebral body height; PH, pedicle height; PW, pedicle widths; APHi, inferior articular process height; APHs, superior articular process height; Adm, maximum distance between the articular process.

The results of the two independent samples *t*-test regarding normal data and the Mann-Whitney test regarding non-normal data showed that the mean difference of all T11 indices, except FW, FD, and APHi, was statistically significant at a 95% confidence level in the two groups (Table 2).

According to Table 3, out of the 14 indices, 8 indices measured in the T10 vertebra in men were shown to have a statistically significant relationship with height. Additionally, the VL, EPD, EPW, FW, TDm, VBH, PH, and ADm indices had significant positive correlations with height in men. Of the 14 indices, 9 indices measured in the T10 vertebra in women were found to have a statistically significant relationship with height. Moreover, VL, EPD, EPW, TDm, SPH, SPL, PH, APHs, and ADm were significantly and positively correlated, and FD was negatively correlated with height in women.

According to Table 3, of the 14 indices, 7 indices measured in the T11 vertebra among men had a

statistically significant relationship with height. Furthermore, VL, EPD, EPW, FW, FD, TDm, and VBH in men were significantly and positively correlated with height. Out of the 14 indices, 6 indices measured in the T11 vertebra in women had a statistically significant relationship with height. Notably, VL, EPW, TDm, SPL, PH, APHs, and PW were significantly and positively correlated with height in women.

According to Table 4, one of the indices measured in the T10 vertebra among men had a statistically significant relationship with age. The VL index in men was significantly and positively correlated with age. Of the 14 indices, 6 indices measured in the T10 vertebra in women had a statistically significant relationship with age. Moreover, VL, EPD, TDm, SPH, SPL, and APHi were observed to be significantly and positively correlated with age in women.

According to Table 4, one of the indices measured in the T11 vertebra in men had a statistically significant

able 3. Correlations of the T10 and T11 Vertebral Indices with Height in Men and Women										
	T10					T11				
	Men		Women		Men		Women			
	Correlation Coefficient	P-Value	Correlation Coefficient	P-Value	Correlation Coefficient	P-Value	Correlation Coefficient	P-Value		
VL	0.4458	0.0007	0.3766	0.0099	0.3312	0.0144	0.3860	0.0081		
EPD	0.4020	0.0026	0.1132	0.4538	0.3029	0.0260	0.2370	0.1128		
EPW	0.2984	0.0284	0.3393	0.0211	0.3514	0.0092	0.3954	0.0065		
FD	0.2438	0.0756	-0.2959	0.0458	0.4051	0.0024	-0.03126	0.8366		
FW	0.3864	0.0039	-0.2206	0.1406	0.3110	0.0221	0.07781	0.6073		
TDm	0.3948	0.0031	0.3779	0.0096	0.3347	0.0134	0.1360	0.3675		
SPH	0.2475	0.071	0.2882	0.0521	-0.08733	0.5300	0.1310	0.3854		
SPL	-0.006489	0.9629	0.3452	0.0188	-0.005362	0.9693	0.2895	0.0510		
VBH	0.2713	0.047	0.2708	0.0687	0.3569	0.0081	0.1557	0.3013		
PH	0.3633	0.0069	0.6308	< 0.0001	0.2180	0.1133	0.4778	0.008		
PW	0.06987	0.6156	0.2282	0.1272	0.2405	0.0799	0.3334	0.0236		
APHi	0.1246	0.3693	0.1850	0.2185	0.2275	0.0974	0.06489	0.6683		
APHs	-0.002045	0.9883	0.2934	0.0478	-0.00256	0.9871	0.3348	0.0230		
ADm	0.3866	0.0039	0.4312	0.0028	0.1941	0.1595	0.2824	0.0611		

Abbreviations: VL, vertebral length; EPD, endplate depth; EPW, endplate width; FD, foramen diameter; FW, foramen width; TDm, transverse process distance; SPH, spinous process height; SPL, spinous process length; VBH, vertebral body height; PH, pedicle height; PW, pedicle widths; APHi, inferior articular process height; APHs, superior articular process height; Adm, maximum distance between the articular process.

relationship with age. The APHs index among men was found to be significantly and positively correlated with the subjects' age. Among the 14 indices, 5 indices measured in the T11 vertebra in women had a statistically significant relationship with age. Furthermore, VL, EPW, EPD, SPL, and APHi were significantly and positively correlated with age in women.

Cut-off points and the areas under the curve (AUC) for VL, EPD, EPW, FD, FW, TDm, SPH, SPL, VBH, PH, PW, APHi, APHs, ADm were determined for the T10 vertebra using receiver operating characteristic (ROC) analysis, as shown in Table 5.

Cut-off points and AUCs for VL, EPD, EPW, FD, FW, TDm, SPH, SPL, VBH, PH, PW, APHi, APHs, and ADm for the T11 vertebra were determined using ROC analysis, as shown in Table 6.

#### 5. Discussion

The present study aimed to evaluate height, age, and gender by measuring 3D CT scan indices of the T10 and T11 vertebrae in adults. The relationship between the individuals' height, age, and gender, and the spine's morphology is known as a helpful factor in identifying a person in forensic medicine. For this purpose, in cases where only some parts of the spine are available, the morphology of the spine's vertebrae can be used to understand various skeletal features. According to the present study, there was a significant statistical difference (P < 0.0001) in the mean heights of male and female patients, with the height being higher in men. This finding is consistent with Tsubaki et al.'s research, reporting that the area of the 10th thoracic vertebra was 970.5 mm<sup>2</sup> in male subjects and 774.0 mm<sup>2</sup> in female subjects. The difference in the area of the 10th thoracic vertebra between male and female subjects was approximately 25% (12). Similarly, Ramadan et al. found that male subjects had higher mean values than female subjects across all measured dimensions of the T12th vertebra (10).

The mean difference of all T10 indices, except FW, FD, and APHi, was statistically significant in the two male and female groups. There was also a significant mean difference between men and women regarding all T11 indices except FW, FD, and APHi. Tsubaki et al.'s study showed that five indices of the 10th thoracic vertebra were significantly different in male and female patients. The accuracy of gender determination in male, female, and total subjects was 85.0%, 88.0%, and 86.5%, respectively (12). In Badr El Dine and El Shafei's study on the T12 vertebra, among 24 linear measurements, 14 measurements displayed a significant gender difference; the predictive accuracy was between 49.0% and 85.5%; and lower EPD, upper EPW, and superior VL had the higher predictive accuracy (7). In Yu et al.'s study in Korea, from 35 traits of the T12 vertebra, 23 showed

Fable 4. Correlations of the T10 and T11 Vertebral Indices with Age in Men and Women								
	T10						T11	
	Me	n	Won	Women		Men		ien
	Correlation Coefficient	P-Value	Correlation Coefficient	P-Value	Correlation Coefficient	P-Value	Correlation Coefficient	P-Value
VL	0.7978	0.03568	0.3944	0.0067	0.007338	0.9580	0.4013	0.0057
EPD	-0.1353	0.3293	0.4474	0.0018	0.01007	0.9424	0.3809	0.0090
EPW	0.1207	0.3846	0.2755	0.0638	0.06066	0.6630	0.3048	0.0394
FD	-0.1117	0.4213	-0.1138	0.4514	-0.2384	0.0826	-0.2285	0.1267
FW	-0.1595	0.2494	-0.03796	0.8022	-0.1289	0.3528	-0.1727	0.2512
TDm	-0.1062	0.4447	0.1212	0.4224	-0.2633	0.0610	0.08977	0.5530
SPH	0.09183	0.5090	0.4149	0.0041	0.1873	0.1751	0.2066	0.1683
SPL	0.1703	0.2183	0.3884	0.0076	0.1336	0.3356	0.4428	0.0021
VBH	-0.01568	0.9104	0.1155	0.4448	-0.1171	0.3989	0.07705	0.6108
РН	-0.08941	0.5202	0.2413	0.1062	-0.04809	0.7299	-0.02124	0.8886
PW	-0.09618	0.4890	0.07942	0.5998	-0.03391	0.8077	0.2095	0.1623
APHi	0.1106	0.4258	0.3810	0.0090	0.01879	0.8927	0.3760	0.0100
APHs	0.2092	0.1290	0.2834	0.0611	0.2997	0.0277	0.2771	0.0622
ADm	-0.1717	0.2145	0.0716	0.5207	-0.03913	0.7788	0.05989	0.6926

Abbreviations: VL, vertebral length; EPD, endplate depth; EPW, endplate width; FD, foramen diameter; FW, foramen width; TDm, transverse process distance; SPH, spinous process height; SPL, spinous process length; VBH, vertebral body height; PH, pedicle height; PW, pedicle widths; APHi, inferior articular process height; APHs, superior articular process height; Adm, maximum distance between the articular process.

 Table 5. Cut-off Points and the Areas Under the Curve Using Receiver Operating Characteristic Analysis for Vertebral Length, Endplate Depth, Endplate Width, Foramen Diameter, Foramen Width, Transverse Process Distance, Spinous Process Height, Spinous Process Length, Vertebral Body Height, Pedicle Height, Pedicle Widths, Inferior Articular Process Height, Superior Articular Process Height, and Maximum Distance Between the Articular Process for the TiO Vertebra<sup>a</sup>

Indices	Cut off Point	Sensitivity	Specificity	AUC	P-Value
VL	Male < 68.10 < female	91.30	83.33	0.9165	< 0.0001
EPD	Male < 26.15 < female	71.74	90.74	0.8364	< 0.0001
EPW	Male < 33.70 < female	91.30	72.22	0.9080	< 0.0001
FD	Male < * < female	*	*	*	*
FW	Male < * < female	*	*	*	*
TDm	Male < 55.40 < female	82.61	85.19	0.8722	< 0.0001
SPH	Male < 10.05 < female	67.39	57.41	0.6443	0.01319
SPL	Male < 19.90 < female	89.13	70.37	0.8428	< 0.0001
VBH	Male < 21.75 < female	78.26	66.67	0.7820	< 0.0001
РН	Male < 11.95 < female	73.91	62.96	0.7180	0.0001824
PW	Male < 5.850 < female	69.57	64.81	0.7371	< 0.0001
APHi	Male < 18.55 < female	71.74	57.41	0.6163	0.04569
APHs	Male < 12.25 < female	76.09	66.67	0.7534	< 0.0001
ADm	Male < 29.20 < female	93.48	53.70	0.7933	< 0.0001

Abbreviations: VL, vertebral length; EPD, endplate depth; EPW, endplate width; FD, foramen diameter; FW, foramen width; TDm, transverse process distance; SPH, spinous process height; SPL, spinous process length; VBH, vertebral body height; PH, pedicle height; PW, pedicle widths; APHi, inferior articular process height; APHs, superior articular process height; Adm, maximum distance between the articular process; AUC, areas under the curve.

<sup>a</sup> \*: No cut off point was defined with statistical analysis.

significant gender differences. The linear dimensions of the vertebral body, pedicle, mammillary, and transverse processes were found to have significant differences between men and women. In addition, it was demonstrated that the coronal diameters of the endplates were the most effective with gender determination, with an accuracy of over 80.0% (9). In Ramadan et al.'s study, all dimensions of the T12 vertebra significantly differed between the two genders (P < 0.001), and the gender prediction accuracy rate was

 Table 6. Cut-off Points and the Areas Under the Curve Using Receiver Operating Characteristic Analysis for Vertebral Length, Endplate Depth, Endplate Width, Foramen Diameter,

 Foramen Width, Transverse Process Distance, Spinous Process Height, Spinous Process Length, Vertebral Body Height, Pedicle Height, Pedicle Widths, Inferior Articular Process

 Height, Superior Articular Process Height, and Maximum Distance Between the Articular Process for the Til Vertebra<sup>a</sup>

Indices	Cut off Point	Sensitivity	Specificity	AUC	P-Value
VL	Male < 68.25 < female	80.43	88.89	0.8861	< 0.0001
EPD	Male < 28.85 < female	84.78	66.67	0.8404	< 0.0001
EPW	Male < 36.00 < female	84.78	88.89	0.9034	< 0.0001
FD	Male <* < female	*	*	*	*
FW	Male < * < female	*	*	*	*
TDm	Male < 54.35 < female	93.48	74.07	0.8786	< 0.0001
SPH	Male < 10.35 < female	67.39	70.37	0.7105	0.0003009
SPL	Male < 20.40 < female	89.13	66.67	0.8058	< 0.0001
VBH	Male < 22.75 < female	76.09	62.96	0.7182	0.0001799
РН	Male < 13.10 < female	86.96	61.11	0.7701	< 0.0001
PW	Male < 7.450 < female	69.57	66.67	0.7361	< 0.0001
APHi	Male < 19.95 < female	69.57	62.96	0.6634	0.005006
APHs	Male < 12.35 < female	63.04	74.07	0.7148	0.0002269
ADm	Male < 28.45 < female	78.26	72.22	0.8037	< 0.0001

Abbreviations: VL, vertebral length; EPD, endplate depth; EPW, endplate width; FD, foramen diameter; FW, foramen width; TDm, transverse process distance; SPH, spinous process height; SPL, spinous process length; VBH, vertebral body height; PH, pedicle height; PW, pedicle widths; APHi, inferior articular process height; APHs, superior articular process; AUCs, area

<sup>a</sup> \*: No cut off point was defined with statistical analysis.

defined at 88.6% (10). Hou et al. have reported that all dimensions of T12 significantly differed between genders, except in terms of sagittal FD and the right inferior APH. All dimensions had higher mean values in male subjects than in females. The prediction value for gender was determined to be between 56.4 and 90.1% (11).

According to the results of this study, in the T10 vertebra, VL, EPD, EPW, FW, TDm, VBH, PH, and ADm were significantly and positively correlated with height in men, and the VL, EPD, EPW, TDm, SPH, SPL, PH, APHs, and ADm indices were found to be significantly and positively correlated with height in women. The results of the current study show that the VL, EPD, EPW, FW, FD, TDm, and VBH indices in the T11 vertebra were significantly and positively correlated with height in men. Moreover, VL, EPW, TDm, SPL, PH, APHs, and PW in the T11 vertebrae were positively correlated with height in women. Therefore, in the T11 vertebra, the VL, EPW, and TDm indices in the whole study population (including men and women) had a positive relationship with height. Zyoud et al.'s reported that the linear regression formula derived from post-mortem CT (PMCT) thoracic length parameters gave a good correlation with autopsy body length, so it could be used for a precise estimation of the body height (13). Moreover, Yudianto and Wibowo's study showed that the correlation between the T1-T12 vertebral indices and the

body height was reasonably strong, and the regression formula could be used in the height estimation (14).

Concerning the correlation between age and thoracic vertebral indices, the present study showed that the VL index in the T10 vertebra in men and the VL, EPD, TDm, SPH, SPL, and APHi indices in women had a positive correlation with the subjects' age. Additionally, in the T11 vertebra, APHs in men and VL, EPW, EPD, SPL, and APHi in women were positively correlated with age. According to Ramadan et al.'s study in Egypt, a significant positive correlation was found between age and all dimensions of the T12 vertebra, except length of the vertebral foramen (LVF), transverse distance (TD), PW, and TDm (10). Furthermore, in their study using 3D CT imaging, Kacar et al. reported that the middle thoracic region had the highest correlation with 20 to 70 years of age, and the lower thoracic region revealed a higher correlation with 70 to 90 years of age (15).

This study showed that most of the T10 and T11 vertebral indices were significantly correlated with gender, height, and age, so they can be used to determine height, age, and gender and, consequently, for the patient's identification. Given that the participants' mean age was about 33 years, and they were young and had no degenerative vertebra changes, the results of this study can be useful and even be used as criteria for determining bone age.

#### Footnotes

**Authors' Contribution:** Study concept and design: D. T., B. SH., and A. M.; acquisition of data: N. M. and A. M.; analysis and interpretation of data: MR. KH., M. V. F., and N. M.; drafting of the manuscript: MR. KH.; critical revision of the manuscript for important intellectual content: M. V. F. and N. M.; statistical analysis: A. M.; administrative, technical, and material support: MR. KH.; study supervision: D. T. and B. SH.

**Conflict of Interests:** The authors confirm that there was no conflict of interest.

**Data Availability:** The dataset presented in the study is available on request from the corresponding author during submission or after publication.

**Ethical Approval:** This study was approved by the Ethics Committee of AJA University of Medical Sciences, Tehran, Iran (Code: IR.AJAUMS.REC.1400.199).

**Funding/Support:** The present study received no funding.

**Informed Consent:** Informed consent was obtained from the patients for the anonymous use of their information and group publication of the results.

#### References

- 1. Pal A, De S, Sengupta P, Maity P, Dhara PC. Estimation of stature from hand dimensions in Bengalee population, West Bengal, India. *Egypt J Forensic Sci.* 2016;6(2):90-8. https://doi.org/10.1016/j.ejfs.2016.03.001.
- Iscan MY, Loth SR, King CA, Shihai D, Yoshino M. Sexual dimorphism in the humerus: a comparative analysis of Chinese, Japanese and Thais. *Forensic Sci Int.* 1998;**98**(1-2):17-29. [PubMed ID: 10036756]. https://doi.org/10.1016/s0379-0738(98)00119-4.
- Akhlaghi M, Sheikhazadi A, Naghsh A, Dorvashi G. Identification of sex in Iranian population using patella dimensions. *J Forensic Leg Med.* 2010;17(3):150-5. [PubMed ID: 20211456]. https://doi.org/10.1016/j.jflm.2009.11.005.
- 4. Ozer I, Katayama K. Sex determination using the femur in an ancient Japanese population. *Coll Antropol.* 2008;**32**(1):67-72. [PubMed ID: 18494190].

- Kranioti EF, Michalodimitrakis M. Sexual dimorphism of the humerus in contemporary Cretans-a population-specific study and a review of the literature\*. J Forensic Sci. 2009;54(5):996-1000. [PubMed ID: 19627412]. https://doi.org/10.1111/j.1556-4029.2009.01103.x.
- Padovan L, Ulbricht V, Groppo FC, Neto JSP, Andrade VM, Junior LF. Sexual dimorphism through the study of atlas vertebra in the Brazilian population. *J Forensic Dent Sci.* 2019;**11**(3):158-62. [PubMed ID: 32801589]. [PubMed Central ID: PMC7398361]. https://doi.org/10.4103/jfo.jfds\_85\_19.
- Badr El Dine FMM, El Shafei MM. Sex determination using anthropometric measurements from multi-slice computed tomography of the 12th thoracic and the first lumbar vertebrae among adult Egyptians. Egypt J Forensic Sci. 2015;5(3):82-9. https://doi.org/10.1016/j.ejfs.2014.07.005.
- 8. Gülek B, Durgun B, Özer HTE, Nazan Alparslan Z, Sarpel T, Erken E. CTbased Morphometric Data of L3-L5 Vertebrae: Anatomic and Surgical Approach. *Neurosurg* Q. 2007;**17**(2):92-7. https://doi.org/10.1097/WNQ.0b013e318032e0ca.
- Yu SB, Lee UY, Kwak DS, Ahn YW, Jin CZ, Zhao J, et al. Determination of sex for the 12th thoracic vertebra by morphometry of threedimensional reconstructed vertebral models. *J Forensic Sci.* 2008;**53**(3):620-5. [PubMed ID: 18471207]. https://doi.org/10.1111/j.1556-4029.2008.00701.x.
- Ramadan N, Abd El-Salam MH, Hanon AF, El-Sayed NF, Al-Amir AY. Age and Sex Identification Using Multi-slice Computed Tomography of the Last Thoracic Vertebrae of an Egyptian Sample. J Forensic Res. 2017;8(4):1000386. https://doi.org/10.4172/2157-7145.1000386.
- Hou WB, Cheng KL, Tian SY, Lu YQ, Han YY, Lai Y, et al. Metric method for sex determination based on the 12th thoracic vertebra in contemporary north-easterners in China. *J Forensic Leg Med.* 2012;**19**(3):137-43. [PubMed ID: 22390999]. https://doi.org/10.1016/j.jflm.2011.12.012.
- Tsubaki S, Morishita J, Usumoto Y, Sakaguchi K, Matsunobu Y, Kawazoe Y, et al. Sex determination based on a thoracic vertebra and ribs evaluation using clinical chest radiography. *Leg Med (Tokyo)*. 2017;**27**:19-24. [PubMed ID: 28668479]. https://doi.org/10.1016/j.legalmed.2017.06.003.
- 13. Zyoud TYT, Abdul Rashid SN, Suppiah S, Mahmud R, Kabeer A, Abd Manaf R, et al. Estimation of body height from spinal length measurements using post-mortem computed tomographic images. *Malays J Pathol*. 2020;**42**(3):423-31. [PubMed ID: 33361724].
- Yudianto A, Wibowo A. Correlation between Vertebral Length and Body Height (Os. Vertebralis) in Indonesian People. *Majalah Kedokteran Bandung.* 2019;51(1):13-6. https://doi.org/10.15395/mkb.v51n1.1472.
- Kacar E, Unlu E, Beker-Acay M, Balcik C, Gultekin MA, Kocak U, et al. Age estimation by assessing the vertebral osteophytes with the aid of 3D CT imaging. *Aust J Forensic Sci.* 2016;49(4):449-58. https://doi.org/10.1080/00450618.2016.1167241.