Published Online: 2025 May 12

Research Article



Validation of GLIM Criteria for Diagnosing Malnutrition in Patients at the Central Militar Hospital in Mexico City

Anahi Merceda-Hernández¹, Ximena Márquez-Ibarra¹, Sandra O'Neill-Suárez², Luis Enrique Orozco-Rivera², Paola Sánchez-Corrales², José Antonio García-Castellanos³, Exsal M. Albores-Méndez¹, Rodolfo David Mayen-Quinto¹, Marco A. Vargas-Hernández¹, Gabriela Gutierrez-Salmean^{2,4,*}

¹ Military Medical School, Ministry of National Defense, Mexico City, Mexico

² Center for Health Sciences Research (CICSA), Faculty of Health Sciences, Anáhuac University, Mexico City, Mexico

³ Central Military Hospital, Secretariat of National Defense, Mexico City, Mexico

⁴ Military School of Health Graduates, Ministry of National Defense, Mexico City, Mexico

* Corresponding Author: Center for Health Sciences Research (CICSA), Faculty of Health Sciences, Anáhuac University, Mexico City, Mexico. Email: gabrielasalmean@yahoo.com

Received: 10 January, 2025; Revised: 8 April, 2025; Accepted: 8 May, 2025

Abstract

Background: Hospital malnutrition remains a significant global issue. Accurate screening and diagnosis within 24 to 48 hours of admission are crucial for reducing morbidity and mortality. The global leadership initiative on malnutrition (GLIM) criteria provide a standardized framework for diagnosing malnutrition, but their validation in military hospitals has been limited.

Objectives: To validate the GLIM criteria for diagnosing malnutrition in hospitalized patients at a third-level military hospital.

Methods: A cross-sectional study was conducted involving hospitalized patients at Mexico City's Central Military Hospital. The GLIM criteria were compared to the subjective global assessment (SGA). Sensitivity, specificity, predictive values, and concordance were assessed.

Results: Nutritional screening indicated that 19.5% of patients were malnourished according to the GLIM criteria, while 19.8% were identified as malnourished by the SGA. Regarding severity, 18.3% had moderate malnutrition and 1.2% had severe malnutrition according to the GLIM criteria; for the SGA, the corresponding prevalences were 18.2% and 1.5%. In terms of validation, compared to the SGA, the GLIM criteria demonstrated a sensitivity of 81.25% and a specificity of 95.37%, with moderate agreement (Kappa = 0.755).

Conclusions: The GLIM criteria are a reliable diagnostic tool for hospital-acquired malnutrition and should be implemented in accordance with global standardization.

Keywords: Malnutrition, Military Hospital, Validation Studies

1. Background

Malnutrition (undernutrition) is defined as the state resulting from an insufficient intake, uptake, or use of nutrients that leads to altered body composition (decreased fat-free mass and body cell mass), which in turn limits physical and mental function and is associated with clinical outcomes (1). Three different types of malnutrition have been proposed and accepted as a consensus: disease-related malnutrition (DRM) with or without inflammation, and malnutrition without disease, such as hunger-related malnutrition (1). Within the hospital setting, all of these types may develop.

Hospital malnutrition is a common worldwide problem, with prevalence rates ranging from 19% to 80%, according to extensive documentation. To identify malnutrition risk in a timely manner, nutritional screening must be implemented for all patients within the first 24 to 48 hours of hospital admission. Consequently, appropriate nutritional therapy can be

Copyright © 2025, Annals of Military and Health Sciences Research. This open-access article is available under the Creative Commons Attribution-NonCommercial 4.0 (CC BY-NC 4.0) International License (https://creativecommons.org/licenses/by-nc/4.0/), which allows for the copying and redistribution of the material only for noncommercial purposes, provided that the original work is properly cited.

How to Cite: Merceda-Hernández A, Márquez-Ibarra X, O'Neill-Suárez S, Orozco-Rivera L E, Sánchez-Corrales P, et al. Validation of GLIM Criteria for Diagnosing Malnutrition in Patients at the Central Militar Hospital in Mexico City. Ann Mil Health Sci Res. 2025; 23 (2): e159541. https://doi.org/10.5812/amh-159541.

implemented promptly, thereby lowering mortality, morbidity, and length of hospital stay (2).

Several nutritional screening tools have been developed for the prompt identification of at-risk patients and to facilitate referrals for more comprehensive nutritional care (3), such as the Mini Nutritional Assessment, Nutritional Risk Screening 2002, Malnutrition Universal Screening Tool, and subjective global assessment (SGA), among others. Although these tools include overall similar indicators, not all have the same concordance, which explains the wide range of hospital malnutrition prevalence. In this context, the SGA has been extensively validated in hospitalized patients with different conditions (e.g., cancer, cirrhosis, kidney disease, major surgery), yielding excellent reproducibility and reliability (i.e., kappas and rhos > 0.7). It correlates not only with anthropometric data (percent weight loss, BMI, tricipital fold, and arm circumference) but also with biochemical indicators (albumin, transferrin, and total cholesterol) as it includes data from clinical history (food and fluid intake, weight change, and gastrointestinal symptoms), а physical exam (evaluating muscle and fat stores), and metabolic demand.

Nevertheless, in 2018, an international group of experts in hospital nutrition proposed the global leadership initiative on malnutrition (GLIM) criteria. This initiative aimed to establish a unified and more global practical approach for diagnosing and classifying hospital malnutrition. The framework drew from the consensus of the Academy of Nutrition and Dietetics (AND), the American society of parenteral and enteral nutrition (ASPEN), and the European society for clinical nutrition and metabolism (ESPEN). According to this global consensus, diagnosing malnutrition requires at least one phenotypic criterion and one etiological criterion. The severity of malnutrition (moderate or severe) is further determined based on phenotypic criteria (e.g., moderate malnutrition when weight loss is > 5%) (4). Although this consensus report was published in 2019, not all institutions have implemented this tool; however, different studies have validated the GLIM criteria – indeed, versus the SGA – yielding even higher feasibility and patient acceptability (5). Moreover, the GLIM criteria have also proven to be a good predictor of short-term (i.e., 30 days) mortality and hospital readmission (6, 7).

2. Objectives

Given the aforementioned, the objective of our study was to validate the GLIM criteria in a third-level military hospital, considering that, worldwide, no information has been reported within this population other than in studies by a group assessing retrospective multicentered cohorts, which include army facilities (4, 8).

3. Methods

We conducted a cross-sectional study in hospitalized patients at a third-level institution (Mexico's Central Military Hospital). Consecutive case sampling was used. Once participants agreed to be part of the study, they were asked to sign an informed consent form. The same interviewer administered both tools (GLIM and SGA). In all cases, data were obtained from primary sources (i.e., the patient and their companions), with the exception of laboratory values, which were obtained from patients' chart registers. Body weight was measured using an InBody Dual scale, and height was assessed with a SECA model 213 stadiometer, using the techniques described by Roberts et al. (5). A physical examination was performed to complete the SGA questionnaire, and patients were then classified into one of three categories (<mark>6</mark>):

- A: No malnutrition
- B: Moderate malnutrition
- C: Severe malnutrition

The SGA assessment includes weight loss during the past 6 months and the last 2 weeks, sufficient dietary intake and changes with respect to the normal diet (liquid, hypocaloric, appetite perception), gastrointestinal symptoms such as nausea, vomiting, diarrhea, and functional capacity, together with the physical examination of fat (suborbital, triceps, and bicipital) and muscle (temporal, clavicular, scapular, calf circumference, interosseous) reserves, and the presence of fluid overload (9).

GLIM was further completed, and patients were classified as malnourished if they fulfilled at least one phenotypic criterion and one etiological criterion. The severity of malnutrition was also staged according to GLIM guidelines. In brief, GLIM phenotypic criteria include involuntary weight loss and/or low BMI and/or reduced muscle mass, whereas etiological criteria include either reduced food intake or an inflammatory condition (4).

For the criterion of reduced muscle mass, bioimpedance (with InBody Dual) was used, employing the cut-off points of the described parameters. The values indicated by the European Society of Clinical Nutrition and Metabolism (ESPEN) for fat-free mass (fatfree mass/height in meters squared) were used, indicating that for men it is 17 kg/m^2 and for women, 15 kg/m^2 (4). Hand-grip strength was measured using a CAMRY EH101 digital hand dynamometer. The patient stood upright with arms slightly away from the body. Two measurements were taken on each hand, alternating between hands, and recording the highest value. Results were classified as "weak" or "normal" according to gender. Finally, the calf circumference of each leg was measured – as described elsewhere – using a non-elastic tape placed horizontally, touching the entire circumference without pinching the skin. The tape was moved up and down to determine the maximum circumference perpendicular to the axis of the calf (10). The present protocol was approved and registered by IRB number 2023FNV082.

3.1. Statistical Analysis

Quantitative variables are presented as measures of central tendency and dispersion, in accordance with their distribution. Qualitative variables are presented as frequency and percentages. For inferential statistics (11), concordance reliability was assessed with Cohen's Kappa, considering > 0.80 as a substantial strong correlation. Finally, for construct validity, we used hospital length of stay (LOS) and considered it to be significant with a P-value < 0.05.

4. Results

During June to July 2024, all hospitalized patients who consented to participate were considered. A total of 323 patients were screened; 48.6% were male, and 78% were under 65 years old. Patients were hospitalized in different departments: 6.8% from Cardiology, 8.4% from Men's Surgery, 9.9% from Women's Surgery, 1.5% from Plastic Surgery, 3.1% from Vascular Surgery, 4% from Endocrinology, 6.2% from Gastroenterology, 6.5% from Hematology, 6.5% from Infectology, 6.2% from Men's Medicine, 7.4% from Women's Medicine, 2.8% from Nephrology, 4.3% from Pneumology, 1.2% from Neurosurgery, 3.1% from Neurology, 11.1% from Oncology, 2.8% from Orthopedics, 2.2% from Urology. Oncology had the highest number of patients screened, while Neurosurgery had the lowest availability of patients for screening. Demographic data are described in Table 1.

Table 1. Demographic Data of Studied Sample ^a		
Variables	Values	
Gender (male); No. (%)	155 (48.6)	
Age (y)	53 ± 8	
Body Mass Index (kg/m ²)	26.3 ± 4.3	
Fat mass (%)	34.5 ± 9.1	
Muscle mass (kg)	24.1 ± 5.4	
Calf circumference (cm)	33.7 ± 3.9	
Hand grip strength (kg)	19.4 ± 10.3	
Length of stay(d)	5.5 ± 6.3	

^a Values are expressed as mean ± SD unless otherwise indicated.

Nutritional screening identified 19.5% of patients as malnourished according to the GLIM criteria, while 19.8% were identified as such by the SGA. The most frequent indicator in both tools was unintentional weight loss, followed by the presence of an inflammatory condition (e.g., major surgery, neoplasia). As shown in Figure 1, regarding severity, similar results were found: 18.3% had moderate malnutrition and only 1.2% presented severe malnutrition according to the GLIM criteria; for the SGA, the corresponding prevalences were 18.2% and 1.5%.



Figure 1. Prevalence of malnutrition, according to global leadership initiative on malnutrition (GLIM) and subjective global assessment (SGA) criteria.

According to the validation guidelines designed by the GLIM group (9), the tool is accepted as valid when, after a significant association (P < 0.0001 for chi-square, and OR 89.19, CI 37.95 - 209.6), concordance (i.e., Kappa) versus the gold standard (in this case, SGA) is at least moderate (0.6 - 0.8). Moreover, as with any other diagnostic test, the sum of its sensitivity and specificity should be at least 1.5, and its predictive values should be more than 80%. All these criteria are met, as shown in Table 2.

Table 2. Diagnostic Performance Between Global Leadership Malnutrition and Subjective Global Assessment Seesement Seesement	Initiative or
Statistical Indicator	Values
χ^2	184.128
OR	89.19
Kappa Coefficient	0.755
Sensitivity	0.8125
Specificity	0.9537
Positive Predictive Value	0.8095
Negative Predictive Value	0.9500

Abbreviations: GLIM, Global leadership initiative on malnutrition; SGA, subjective global assessment.

Thus, with this validation, we compared the length of stay, confirming that malnutrition is positively associated with it: as seen in Figure 2, patients with a GLIM-diagnosed nutritional status depletion had a significantly higher length of stay (8 vs. 5 days, P < 0.001).



Figure 2. Comparison in LOS (days) between patients with/without malnutrition, according to Global Leadership Initiative on Malnutrition (GLIM) criteria. (P < 0.001 after independent *t*-test)

5. Discussion

Our study screened a total of 323 patients from all services at a military hospital and validated the GLIM criteria for the diagnosis of malnutrition. When compared to the SGA, the percentages of altered nutritional status were fairly similar (19.8% with SGA and 19.5% with GLIM). Likewise, the severity of malnutrition was also concordant (18.3% presented moderate malnutrition according to SGA and 18.3% according to GLIM; 1.5% presented severe malnutrition according to SGA and 1.2% according to GLIM).

The above is extremely important since, at an international level, there is an attempt to obtain a unified and valid scale to improve and optimize the timely diagnosis of malnutrition in patients from different populations. This is in addition to avoiding the consequences that may otherwise be caused, such as increased days of hospital stay, complications of the underlying disease, risk of contracting nosocomial diseases, risk of falls, and increased hospital expenses, both for the institution that provides medical services and for patients and their families (12).

These findings are in line with other validation studies of the GLIM criteria. For example, a study by Maeda et al. reported a similar prevalence of malnutrition (22.4%) using GLIM in hospitalized elderly patients, with high agreement with SGA ($\kappa = 0.84$) (13). Likewise, another group found prevalences of 23.7% with GLIM criteria and 24.5% with SGA in internal medicine patients in Spain, reinforcing the reliability of GLIM in clinical settings (14). Our results are slightly lower, which may be due to differences in population characteristics or care protocols in a military hospital setting.

Regarding severity classification, our data also showed strong concordance. This supports observations by other studies, which have found that GLIM provides comparable gradation to SGA, though it may slightly underestimate severe cases when inflammatory markers are not routinely used (15). Similarly, Silva et al. reported 20.6% of cancer patients diagnosed with malnutrition by GLIM versus 22% by SGA, with good diagnostic agreement ($\kappa = 0.76$) (16).

Nevertheless, some studies show greater discrepancies. For example, when studying critically ill patients, malnutrition rates are around 34% with GLIM and 41% with SGA, suggesting that SGA may be more sensitive in critically ill patients (17). These differences highlight the importance of evaluating diagnostic tools within specific patient contexts to ensure accurate and timely identification.

Finally, malnutrition is associated with a longer hospital stay. This implies a greater need for resources and care for malnourished patients. Furthermore, as shown in Figure 2, significant weight loss in malnourished patients indicates a deterioration in nutritional status and the need for more detailed nutritional and medical intervention (18).

5.1. Conclusions

The GLIM criteria proved to be a valid instrument for the diagnosis of malnutrition, showing a strong correlation and high level of agreement with the SGA when applied under the same conditions. We validated this tool at the Central Military Hospital, and it should be used to assess malnutrition to obtain comparable results to those reported worldwide.

5.2. Limitations

The study was conducted in a center that serves only military beneficiaries or insured individuals, so it does not include a sample from all areas of the country.

Footnotes

Authors' Contribution: Concept and design of study: G. G. S., P. S. C., and J. A. G. C.; Data acquisition and analysis: A. M. H., X. M. I., and S. O. S.; Statistical analysis: G. G. S., E. M. A. M., and M. A. V. H.; Wrote the manuscript: S. O. S., L. E. O. R., and R. D. M. Q.; Coordinated the analysis and reviewed the manuscript: G. G. S. All authors approve final manuscript.

Conflict of Interests Statement: The authors declared no conflict of interests.

Data Availability: The dataset presented in the study is available on request from the corresponding author during submission or after publication.

Ethical Approval: IRB 20234352.

Funding/Support: The authors declare that no funding was received for this study.

Informed Consent: Informed consent was obtained from all participant.

References

- Cederholm T, Barazzoni R, Austin P, Ballmer P, Biolo G, Bischoff SC, et al. ESPEN guidelines on definitions and terminology of clinical nutrition. *Clin Nutr.* 2017;**36**(1):49-64. [PubMed ID: 27642056]. https://doi.org/10.1016/j.clnu.2016.09.004.
- Navarro P, Capelli O, Adaglio J, Barritta R. [GLIM criteria: analysis of their internal consistency and a comparison with the subjective global assessment]. *Rev Nutr Clin Metab.* 2021;4(4):14-23. ES. https://doi.org/10.35454/rncm.v4n4.304.
- Lobatón E. [Hospital malnutrition: etiology and criteria for diagnosis and classification]. *Rev Nutr Clin Metab.* 2020;3(1):121-7. ES. https://doi.org/10.35454/rncm.v3n1.019.
- Cederholm T, Jensen GL, Correia M, Gonzalez MC, Fukushima R, Higashiguchi T, et al. GLIM criteria for the diagnosis of malnutrition -A consensus report from the global clinical nutrition community. *Clin Nutr.* 2019;**38**(1):1-9. [PubMed ID: 30181091]. https://doi.org/10.1016/j.clnu.2018.08.002.
- Roberts S, Nucera R, Dowd T, Turner K, Langston K, Keller H, et al. Prospective validation of the Global Leadership Initiative on Malnutrition criteria for identifying malnutrition in hospitals: A protocol and feasibility pilot study. *Nutr Clin Pract.* 2024;**39**(6):1406-18. [PubMed ID: 38711239]. https://doi.org/10.1002/ncp.11156.
- Gersely GD, Klein RCM, da Rocha G, Bruzaca WFS, Ribeiro LMK, Santos BC, et al. GLIM criteria validation and reliability in critically ill patients with cancer: A prospective study. *JPEN J Parenter Enteral Nutr.* 2024;**48**(6):726-34. [PubMed ID: 38850511]. https://doi.org/10.1002/jpen.2657.
- Jazinaki MS, Safarian M, Arabi SM, Jamali J, Norouzy A. Validation of GLIM criteria for hospital malnutrition diagnosis by comparison of three different anthropometric approaches to evaluate reduced muscle mass: a prospective cohort study. *Front Nutr.* 2024;**11**:1438158.
 [PubMed ID: 39703335]. [PubMed Central ID: PMC11655236]. https://doi.org/10.3389/fnut.2024.1438158.
- Zhang Q, Zhang KP, Zhang X, Tang M, Song CH, Cong MH, et al. Scored-GLIM as an effective tool to assess nutrition status and predict survival in patients with cancer. *Clin Nutr.* 2021;40(6):4225-33. [PubMed ID: 33579553]. https://doi.org/10.1016/j.clnu.2021.01.033.
- 9. Duerksen DR, Laporte M, Jeejeebhoy K. Evaluation of Nutrition Status Using the Subjective Global Assessment: Malnutrition, Cachexia, and Sarcopenia. *Nutr Clin Pract*. 2021;**36**(5):942-56. [PubMed ID: 33373482]. https://doi.org/10.1002/ncp.10613.
- Carrasco Llatas M, Palomino Quispe LP, Niño Montero J. [Calf circumference as a predictor of hospital malnutrition in patients with type 2 diabetes mellitus.]. Nutr Clín Diet Hosp. 2023;43(2). ES. https://doi.org/10.12873/432carrasco.
- de van der Schueren MAE, Keller H, Glim Consortium, Cederholm T, Barazzoni R, Compher C, et al. Global Leadership Initiative on Malnutrition (GLIM): Guidance on validation of the operational criteria for the diagnosis of protein-energy malnutrition in adults. *Clin Nutr.* 2020;**39**(9):2872-80. [PubMed ID: 32563597]. https://doi.org/10.1016/j.clnu.2019.12.022.
- Alvarez-Altamirano K, Bejarano-Rosales MP, Gonzalez-Rodriguez BK, Mondragon-Nieto G, Alatriste-Ortiz G, Noguez LJJ, et al. Prevalence of nutritional risk and malnutrition in hospitalized patients: a retrospective, cross-sectional study of single-day screening. *Appl Physiol Nutr Metab.* 2024;49(6):838-43. [PubMed ID: 38700079]. https://doi.org/10.1139/apnm-2023-0190.

- Maeda K, Ishida Y, Nonogaki T, Mori N. Reference body mass index values and the prevalence of malnutrition according to the Global Leadership Initiative on Malnutrition criteria. *Clin Nutr.* 2020;**39**(1):180-184. [PubMed ID: 30712782]. https://doi.org/10.1016/j.clnu.2019.01.011.
- Contreras-Bolívar V, Sánchez-Torralvo FJ, Ruiz-Vico M, González-Almendros I, Barrios M, Padín S, et al. GLIM Criteria Using Hand Grip Strength Adequately Predict Six-Month Mortality in Cancer Inpatients. *Nutrients.* 2019;11(1):2043. https://doi.org/10.3390/nu11092043.
- van der Schueren MAE, Mareschal J, Haenni A. Application of the GLIM criteria for the diagnosis of malnutrition in clinical practice: A prospective study. *Clin Nutr ESPEN*. 2020;35:91-6. https://doi.org/10.1016/j.clnesp.2019.11.010.
- Henriksen C, Paur I, Pedersen A, Sørlie Kværner A, Ræder H, Berg Henriksen H, et al. Agreement between GLIM and PG-SGA for diagnosis of malnutrition depends on the screening tool used in GLIM. Clin Nutr ESPEN. 2022;41(2):329-336. https://doi.org/10.1016/j.clnu.2021.12.024.
- 17. Ribeiro PC, de Brito BL, Gomes ML. Nutritional assessment in intensive care unit: Agreement between Subjective Global Assessment and GLIM criteria. *Nutr Clin Pract.* 2022;**37**(1):180-8. https://doi.org/10.1002/ncp.10696.
- Lorden H, Engelken J, Sprang K, Rolfson M, Mandelbrot D, Parajuli S. Malnutrition in solid organ transplant patients: A review of the literature. *Clin Transplant*. 2023;**37**(11). e15138. [PubMed ID: 37715587]. https://doi.org/10.1111/ctr.15138.