



Effects of Whole Body Vibration Training on Inflammatory Markers in Young Healthy Males

Farshad Ghazalian^{1,*}

¹Department of Physical Education, Science and Research Branch, Islamic Azad University, Tehran, Iran

*Corresponding author: Department of Physical Education, Science and Research Branch, Islamic Azad University, Tehran, Iran. Email: phdghazalian@gmail.com

Received 2019 January 15; Revised 2019 April 21; Accepted 2019 April 29.

Abstract

Background: The intensity of the vibration load refers to frequency and amplitude.

Objectives: The present study aimed at reviewing the effects of high and low intensity whole body vibration training on inflammatory markers.

Methods: Overall, 30 healthy male patients were divided to three groups, including control (N = 8), high intensity (N = 11), and low intensity (N = 11) groups. Training included consecutive weeks, three times each week with low intensity (amplitudes of 2 mm), high intensity (amplitudes of 4 mm), and progressive frequencies. Blood samples were analyzed for concentrations of high sensitivity C-reactive protein (hsCRP) and fibrinogen before and after completion of training. Paired sample *t*-test and One-way Analysis of Variance (ANOVA) were used to analyze the research findings ($P \leq 0.05$).

Results: Whole body vibration training with low and high intensities had no significant effects on hsCRP and fibrinogen levels ($P \geq 0.05$).

Conclusions: It appears that six weeks of low and high intensity vibration training has the same effects on CRP and fibrinogen.

Keywords: Vibration Training, Amplitude, Inflammatory Markers

1. Background

Whole Body Vibrational Training (WBVT) could provide health benefits. In elite athletes, WBVT has been used as preventive medicine and resistance training (1). Furthermore, WBVT regardless of age or fitness status can be beneficial for everyone and it represents a high revolution in health and fitness science. One of the major risk factors for cardiovascular disease is a sedentary lifestyle (2, 3). Besides numerous benefits of exercise, it is well-known that regular exercise minimizes cardiovascular morbidity and mortality (4, 5). Watanabe et al. showed WBVT training along with aerobic training in comparison with aerobic training alone significantly reduced arterial stiffness and increased cardio-respiratory fitness (6). For some groups, such as elderly people, WBVT rather than aerobic training is more safe and effective and has more anti-inflammatory effects (7). So et al. showed that WBVT significantly decreased TC, LDL, and TG (1). Hypertension, smoking, and hyperlipidemia can induce coronary heart disease (8, 9). There are some studies indicating that there is a significant correlation between inflammatory markers and coronary heart disease (9). Furthermore, CRP is a marker of sub-clinical

inflammation. An increase in CRP concentration results in a four- to five-fold increase in coronary risks (10). Fibrinogen is a high molecular weight protein and is formed in the liver (11). Evidence shows that reduction of 1 g/L in fibrinogen concentration results in 15% reduction of coronary disease (12, 13). A wealth of data during the last decade suggests that fibrinogen level represents an independent cardiovascular risk factor. Its impact seems to be as strong as, or even stronger than, that of total cholesterol concentration. If this view is accepted, the question arises as to how fibrinogen and/or CRP levels can be altered in order to reduce the risk. Clearly, this aspect has not yet been fully investigated (14). Studies showed that WBVT have utilized training time periods ranging from ten days to six months, frequencies between 12 to 60 Hz, and amplitudes ranging from 1.7 to 10 mm (15, 16). Various frequencies, amplitudes, and durations have been utilized yet an optimal pattern is not recognized. For proper doses of WBVT, attention should be paid to intensity, strength, and volume of the vibration, which can effect positive biological and physical responses. The volume refers to time under vibration stimulus; intensity refers to the mechanical char-

acteristics of frequency and amplitude and strength refers to observational characteristics of the muscle impacted by the vibration load (17). Little modification frequency and amplitude can lead to relatively large changes in acceleration and magnitude of vibration being transmitted to the body. Furthermore, WBVT in health centers has been recognized as a well-known training method and effect of vibration exercises on risk factors are widely studied in humans and it has also been suggested that mechanical frequency and low range stimulation can be an effective way to prepare muscle structures; nevertheless, the results of the reported studies are contradictory. The reason for the inconsistency in researches is due to the difference in the subject's gender, the time of the test, the age of the subjects, duration and frequency of the vibration, the type of exercise, and the amount of motivation of the subjects. Therefore, a study that can control disruptive factors as much as possible can provide valuable findings regarding the effects of vibration exercises on inflammatory factors. Therefore the main hypothesis of the present study is that high intensity and low intensity vibration have the same influence on inflammatory markers (CRP and fibrinogen).

2. Objectives

Since different intensities of WBVT may have different effects on inflammatory responses; the present study aimed at reviewing the effects of high and low intensity WBVT on inflammatory markers in young healthy males.

3. Methods

3.1. Subject Selection and Measurement

Overall, 36 healthy and sedentary students of Islamic Azad University, Science and Research Branch (in 2018) were selected voluntarily. At first, all subjects filled the physical activity behavior questionnaire. Inclusion criteria were age between 20 and 25 years, no history of physical exercise in the last three months, no smoking, no consumption of systemic corticosteroids. The exclusion criteria included severe co-morbidities or self-reported contraindications for WBV (e.g. deep vein thrombosis, metal implants, pacemaker, epilepsy, tumors, arterial aneurysm, and arrhythmia). All participants were paired for gender, age, and body mass index (BMI). They were informed about the procedures, risks, and benefits of the research in advance.

As six subjects were not able to continue exercise sessions, they were excluded from the study and the rest of the subjects were randomly divided to three groups including high intensity vibration ($N = 11$), low intensity vibration ($N = 11$), and control ($N = 8$). It should be noted that

the present study was blind study, which means none of the subjects knew about the amplitudes of training; however, only trained staff knew about differentiation between groups.

In pre-test and post-test (48 hours after the last training session) all subjects referred to the laboratory (eight hours of overnight fast). Blood samples (5 mL), taken by venipuncture, were then analyzed for concentrations of hsCRP and fibrinogen. Both tests measured the same protein in blood. The hsCRP concentrations were between 0.5 and 10 mg/L. Plasma levels of hsCRP was measured by immunoturbidimetry 5+1 Greiner, Germany, and fibrinogen was measured by Clauss, MahsaYaran, Iran.

3.2. Whole Body Vibration Training

During the six-week experimental period, all subjects continued their conventional living schedule, yet the intervention groups (low intensity and high intensity groups) additionally performed a WBVT program, three times per week with amplitudes of 4 or 2 mm and progressive frequencies from 25 Hz with increments of 5 Hz, weekly. Subjects completed a standardized warm-up. This consisted of five minutes of cycling on a cycle ergometer (Technogym Italy) at 50 W. Furthermore, WBVT was performed on a Vibofit vibration fitness machine (Figure 1). The WBVT program is summarized in Table 1. It should be noted that in this period of time, the control group did not contribute to any training and only had their daily activity.



Figure 1. Vibofit vibration fitness machine

Table 1. The WBVT Training Schedule

Position of Training	Set	Repetition	Frequency in the First Week ^a , Hz	Amplitudes ^b , mm	Duration of Each Repetition, s	Duration of Rest Between Sets, s	Total Duration of Training in the Session, min
1st session of the week			25	4 or 2	30	30	9
Squat	2	3					
Lunges	2	3					
Deep squat	2	3					
2nd session of the week			25	4 or 2	45	30	13.5
Squat	2	3					
Lunges	2	3					
Deep squat	2	3					
3rd session of the week			25	4 or 2	60	30	18
Squat	2	3					
Lunges	2	3					
Deep squat	2	3					

^aThe frequency at beginning of the program was 25 Hz which was progressively increased (5 Hz each week) to the final frequency of 50 Hz.

^b4 mm for the high amplitude and 2 mm for the low amplitude groups.

3.3. Ethical Issues

The study protocol was evaluated and accepted by the Ethical Committee of the Sports Medicine Federation of Iran and complied with the Helsinki Declaration of 1975, revised in 2002. All the participants received adequate information about the study protocol and possible advantages and disadvantages of training. They entered the study deliberately and were free to quit the protocol upon their request.

3.4. Statistical Analysis

All data were found to be normally distributed; therefore, analysis was carried out using parametric statistical tests. Paired sample *t*-test and One-way ANOVA was used to identify any statistically significant differences within and between groups. Data are expressed as mean \pm standard deviation (SD). All statistical analysis was done by SPSS version 18 (SPSS Inc., USA) The level of significance was set at $P \leq 0.05$.

4. Results

Demographic characteristics of all subjects are reported in Table 2. The results of One-way ANOVA showed that there were no significant differences between age ($P = 0.67$), height ($P = 0.90$), weight ($P = 0.55$), and BMI ($P = 0.45$) between high intensity, low intensity, and control groups (Table 2).

The results of One-way ANOVA test showed that WBVT with progressive frequencies between low and high intensity groups caused no significant differences in hsCRP ($P \geq 0.05$) (Figure 2) and fibrinogen (Figure 3) concentrations ($P \geq 0.05$). The results of paired sample *t*-test showed that

there were no significant differences between pre-test and post-test levels of hsCRP and fibrinogen in low intensity ($P \geq 0.05$) and control ($P \geq 0.05$) groups; also, there was no significant difference between pre-test and post-test levels of fibrinogen in the high intensity group ($P \geq 0.05$). Nevertheless, hsCRP in the post-test of high intensity group was significantly increased compared to the pre-test.




Figure 2. hsCRP levels in three groups of research. *Significant increase in post-test compared to pre-test

5. Discussion


Increased CRP and fibrinogen concentrations can reflect low grade chronic inflammation (18). It has been reported that measurement of CRP and fibrinogen can be a method for detection of cardiovascular risk. The present study showed that high and low intensity WBVT caused no

Table 2. The demographic characteristics of groups^a

Variable	Group			P Value ^b
	High Intensity (N = 11)	Low Intensity (N = 11)	Control (N = 8)	
Age, y	21.14 ± 1.51	20.45 ± 1.39	20.88 ± 0.83	0.67
Height, cm	177.20 ± 8.01	174.56 ± 4.30	176.42 ± 2.97	0.90
Weight, kg	70.68 ± 9.21	69.80 ± 15.38	65.20 ± 9.50	0.55
BMI, kg/m ²	22.69 ± 3.23	22.83 ± 4.69	20.89 ± 2.47	0.45

^aValues are expressed as mean ± SD.

^bResults of One-way ANOVA test.

**Figure 3.** Fibrinogen levels in three groups of research

significant differences in hsCRP and fibrinogen concentrations. However, the increase of hsCRP in the high intensity group was significant. In relation to WBVT, various studies have been reported, for example Rittweger et al. reported that WBVT by increase in oxygen uptake significantly increased basal energy metabolism rate (17); also, it has been suggested that in WBVT, the energy expenditure is similar to light work of 0.3 metabolic energy equivalents (19). Furthermore, it has been reported that WBVT similar to those experienced during moderate walking at 4 km/h provides cardiovascular stimuli (20). In one study, three seconds up and three seconds down squat training on a vibration plate similar to cycling at 70 Watt had the same metabolic rate (21) and in another study, WBV session significantly increased the 24-hour oxygen consumption (22). Also it has been reported that 10% WBVT reduced body weight, body fat, and serum leptin (23).

It appears that exercise directly and indirectly reduces CRP; indeed exercises reduce cytokine production in mononuclear cells, muscle, and fat (directly), and also increases endothelial function and insulin sensitivity while it reduces body weight (indirectly). It was found that the endothelium is essential for vasodilatation in response to increases in blood flow associated shear stress (24). In response to shear stress, several vasodilators are released by

the endothelium, which include prostaglandins and nitric oxide (25). During WBVT, blood flow of skin increases (26) and this increase in blood flow could induce reduction in shear stress on blood vessels. Increase in blood flow can increase on endothelial walls of blood vessels. It has also been shown that WBVT leads to the speculation that WBVT may also result in an increased fibrinolytic response (27). Thus, WBVT may enhance the cardio-protective benefits and influence fibrinolytic activity of a therapeutic exercise regimen. Increased vascular shear stress has been shown to increase fibrinolytic activity in cultured endothelial cells (28). The mechanism underlying the fibrinolytic responses reported in the present study may be related to vascular shear stress. As the movement of the platform is sinusoidal, the acceleration transmitted to the body is calculated as $a = A(2\pi f)^2$. Where "A" is the amplitude of the oscillations and "f" is the frequency. The intensity of the vibration load refers to the mechanical characteristics of the frequency and amplitude of the vibration stimulus. Small changes in amplitude and frequency determine relatively large changes in acceleration and magnitude of vibration being transmitted to the body. It seems that WBVT affects CRP and fibrinogen indirectly by improving endothelial function. Frequency and amplitude in WBVT are important variables that affect vascular shear stress factors. In conclusion, amplitude seems to be a less important parameter than frequency for intensity of whole body vibration training programs on inflammatory markers.

5.1. Conclusions

According to the results of the present study, it seems that six weeks of low and high intensity vibration training have the same effects on CRP and fibrinogen.

Acknowledgments

The present study was performed under the supervision of the Department of Physical Education, Science and Research Branch, Islamic Azad University, Tehran, Iran. The costs were paid by the university.

Footnotes

Authors' Contribution: Only one author contributed to the writing and revision of this paper.

Conflict of Interests: The authors declare that they had no conflicts of interest.

Ethical Approval: The present research protocol was evaluated by the Ethical Committee of the Sports Medicine Federation of Iran.

Funding/Support: The present study was funded by the Science and Research Branch, Islamic Azad University, Tehran, Iran.

Informed Consent: All subjects fulfilled the informed consent questionnaire.

References

- So R, Eto M, Tsujimoto T, Tanaka K. Acceleration training for improving physical fitness and weight loss in obese women. *Obes Res Clin Pract.* 2014;**8**(3):e201-98. doi: [10.1016/j.orcp.2013.03.002](https://doi.org/10.1016/j.orcp.2013.03.002). [PubMed: [24847665](https://pubmed.ncbi.nlm.nih.gov/24847665/)].
- Myers J. Exercise and cardiovascular health. *Circulation.* 2003;**107**(1). doi: [10.1161/01.cir.0000048890.59383.8d](https://doi.org/10.1161/01.cir.0000048890.59383.8d).
- Slavicek J, Kittnar O, Fraser GE, Medova E, Konecna J, Zizka R, et al. Lifestyle decreases risk factors for cardiovascular diseases. *Cent Eur J Public Health.* 2008;**16**(4):161-4. doi: [10.21101/cejph.a3474](https://doi.org/10.21101/cejph.a3474). [PubMed: [19256282](https://pubmed.ncbi.nlm.nih.gov/19256282/)]. [PubMed Central: [PMC2745100](https://pubmed.ncbi.nlm.nih.gov/PMC2745100/)].
- Alhassan S, Robinson TN. Objectively measured physical activity and cardiovascular disease risk factors in African American girls. *Ethn Dis.* 2008;**18**(4):421-6. [PubMed: [19157245](https://pubmed.ncbi.nlm.nih.gov/19157245/)]. [PubMed Central: [PMC3863587](https://pubmed.ncbi.nlm.nih.gov/PMC3863587/)].
- Hamer M, Stamatakis E. Physical activity and mortality in men and women with diagnosed cardiovascular disease. *Eur J Cardiovasc Prev Rehabil.* 2009;**16**(2):156-60. doi: [10.1097/HJR.0b013e32831fib77](https://doi.org/10.1097/HJR.0b013e32831fib77). [PubMed: [19276984](https://pubmed.ncbi.nlm.nih.gov/19276984/)].
- Watanabe T, Yabumoto T, Shin S, Shi B, Matsuoka T. Effect of short-term whole-body vibration training on metabolic risk factors, inflammatory markers, and arterial stiffness. *Adv Biosci Biotechnol.* 2014;**5**(5):438-45. doi: [10.4236/abb.2014.55053](https://doi.org/10.4236/abb.2014.55053).
- Mohammed E, Mawad A. Impact of whole body vibration versus aerobic exercise training in modulation of inflammatory markers in elderly. *World J Med Sci.* 2015;**12**(2):148-55. doi: [10.5829/idosi.wjms.2015.12.2.93116](https://doi.org/10.5829/idosi.wjms.2015.12.2.93116).
- Khot UN, Khot MB, Bajzer CT, Sapp SK, Ohman EM, Brener SJ, et al. Prevalence of conventional risk factors in patients with coronary heart disease. *JAMA.* 2003;**290**(7):898-904. doi: [10.1001/jama.290.7.898](https://doi.org/10.1001/jama.290.7.898). [PubMed: [12928466](https://pubmed.ncbi.nlm.nih.gov/12928466/)].
- Ridker PM. C-reactive protein: Eighty years from discovery to emergence as a major risk marker for cardiovascular disease. *Clin Chem.* 2009;**55**(2):209-15. doi: [10.1373/clinchem.2008.119214](https://doi.org/10.1373/clinchem.2008.119214). [PubMed: [19095723](https://pubmed.ncbi.nlm.nih.gov/19095723/)].
- Ridker PM, Hennekens CH, Buring JE, Rifai N. C-reactive protein and other markers of inflammation in the prediction of cardiovascular disease in women. *N Engl J Med.* 2000;**342**(12):836-43. doi: [10.1056/NEJM200003233421202](https://doi.org/10.1056/NEJM200003233421202). [PubMed: [10733371](https://pubmed.ncbi.nlm.nih.gov/10733371/)].
- Hall JE, Guyton AC. *Textbook of medical physiology.* London: Saunders; 2011.
- DeSouza CA, Jones PP, Seals DR. Physical activity status and adverse age-related differences in coagulation and fibrinolytic factors in women. *Arterioscler Thromb Vasc Biol.* 1998;**18**(3):362-8. doi: [10.1161/01.ATV.18.3.362](https://doi.org/10.1161/01.ATV.18.3.362). [PubMed: [9514404](https://pubmed.ncbi.nlm.nih.gov/9514404/)].
- Zanettini R, Bettega D, Agostoni O, Ballestra B, del Rosso G, di Michele R, et al. Exercise training in mild hypertension: Effects on blood pressure, left ventricular mass and coagulation factor VII and fibrinogen. *Cardiology.* 1997;**88**(5):468-73. doi: [10.1159/000177378](https://doi.org/10.1159/000177378). [PubMed: [9286510](https://pubmed.ncbi.nlm.nih.gov/9286510/)].
- Ernst E. *Fibrinogen: A "new" cardiovascular risk factor.* Blackwell-Mzvw; 1992.
- Wannamethee SG, Lowe GD, Whincup PH, Rumley A, Walker M, Lennon L. Physical activity and hemostatic and inflammatory variables in elderly men. *Circulation.* 2002;**105**(15):1785-90. [PubMed: [11956120](https://pubmed.ncbi.nlm.nih.gov/11956120/)].
- Albert MA, Glynn RJ, Ridker PM. Effect of physical activity on serum C-reactive protein. *Am J Cardiol.* 2004;**93**(2):221-5. doi: [10.1016/j.amjcard.2003.09.046](https://doi.org/10.1016/j.amjcard.2003.09.046). [PubMed: [14715354](https://pubmed.ncbi.nlm.nih.gov/14715354/)].
- Rittweger J, Ehrig J, Just K, Mutschelknauss M, Kirsch KA, Felsenberg D. Oxygen uptake in whole-body vibration exercise: Influence of vibration frequency, amplitude, and external load. *Int J Sports Med.* 2002;**23**(6):428-32. doi: [10.1055/s-2002-33739](https://doi.org/10.1055/s-2002-33739). [PubMed: [12215962](https://pubmed.ncbi.nlm.nih.gov/12215962/)].
- Ross R. Atherosclerosis is an inflammatory disease. *Am Heart J.* 1999;**138**(5 Pt 2):S419-20. doi: [10.1016/S0002-8703\(99\)70266-8](https://doi.org/10.1016/S0002-8703(99)70266-8). [PubMed: [10539839](https://pubmed.ncbi.nlm.nih.gov/10539839/)].
- Maikala RV, King S, Bhambhani YN. Acute physiological responses in healthy men during whole-body vibration. *Int Arch Occup Environ Health.* 2006;**79**(2):103-14. doi: [10.1007/s00420-005-0029-8](https://doi.org/10.1007/s00420-005-0029-8). [PubMed: [16175416](https://pubmed.ncbi.nlm.nih.gov/16175416/)].
- Da Silva ME, Fernandez JM, Castillo E, Nunez VM, Vaamonde DM, Poblador MS, et al. Influence of vibration training on energy expenditure in active men. *J Strength Cond Res.* 2007;**21**(2):470-5. doi: [10.1519/R-19025.1](https://doi.org/10.1519/R-19025.1). [PubMed: [17530948](https://pubmed.ncbi.nlm.nih.gov/17530948/)].
- Cochrane DJ, Sartor F, Winwood K, Stannard SR, Narici MV, Rittweger J. A comparison of the physiologic effects of acute whole-body vibration exercise in young and older people. *Arch Phys Med Rehabil.* 2008;**89**(5):815-21. doi: [10.1016/j.apmr.2007.09.055](https://doi.org/10.1016/j.apmr.2007.09.055). [PubMed: [18452726](https://pubmed.ncbi.nlm.nih.gov/18452726/)].
- Hazell TJ, Lemon PW. Synchronous whole-body vibration increases VO(2) during and following acute exercise. *Eur J Appl Physiol.* 2012;**112**(2):413-20. doi: [10.1007/s00421-011-1984-2](https://doi.org/10.1007/s00421-011-1984-2). [PubMed: [21573780](https://pubmed.ncbi.nlm.nih.gov/21573780/)].
- Maddalozzo GF, Iwaniec UT, Turner RT, Rosen CJ, Widrick JJ. Whole-body vibration slows the acquisition of fat in mature female rats. *Int J Obes (Lond).* 2008;**32**(9):1348-54. doi: [10.1038/ijo.2008.111](https://doi.org/10.1038/ijo.2008.111). [PubMed: [18663370](https://pubmed.ncbi.nlm.nih.gov/18663370/)]. [PubMed Central: [PMC2586051](https://pubmed.ncbi.nlm.nih.gov/PMC2586051/)].
- Pohl U, Holtz J, Busse R, Bassenge E. Crucial role of endothelium in the vasodilator response to increased flow in vivo. *Hypertension.* 1986;**8**(1):37-44. doi: [10.1161/01.HYP.8.1.37](https://doi.org/10.1161/01.HYP.8.1.37). [PubMed: [3080370](https://pubmed.ncbi.nlm.nih.gov/3080370/)].
- Joannides R, Haefeli WE, Linder L, Richard V, Bakkali EH, Thuillez C, et al. Nitric oxide is responsible for flow-dependent dilatation of human peripheral conduit arteries in vivo. *Circulation.* 1995;**91**(5):1314-9. doi: [10.1161/01.CIR.91.5.1314](https://doi.org/10.1161/01.CIR.91.5.1314). [PubMed: [7867167](https://pubmed.ncbi.nlm.nih.gov/7867167/)].
- Maloney-Hinds C, Petrofsky JS, Zimmerman G. The effect of 30 Hz vs. 50 Hz passive vibration and duration of vibration on skin blood flow in the arm. *Med Sci Monit.* 2008;**14**(3):CR12-6. [PubMed: [18301353](https://pubmed.ncbi.nlm.nih.gov/18301353/)].
- Kerschman-Schindl K, Grampp S, Henk C, Resch H, Preisinger E, Fialka-Moser V, et al. Whole-body vibration exercise leads to alterations in muscle blood volume. *Clin Physiol.* 2001;**21**(3):377-82. doi: [10.1046/j.1365-2281.2001.00335.x](https://doi.org/10.1046/j.1365-2281.2001.00335.x). [PubMed: [11380538](https://pubmed.ncbi.nlm.nih.gov/11380538/)].
- Hilberg T, Prasa D, Sturzebecher J, Glaser D, Schneider K, Gabriel HH. Blood coagulation and fibrinolysis after extreme short-term exercise. *Thromb Res.* 2003;**109**(5-6):271-7. doi: [10.1016/S0049-3848\(03\)00283-4](https://doi.org/10.1016/S0049-3848(03)00283-4). [PubMed: [12818250](https://pubmed.ncbi.nlm.nih.gov/12818250/)].