




Comparison of Premature Ejaculation in Men with Focal Epilepsy and Generalized Tonic-Clonic Epilepsy

Masoumeh Otaghi¹, Seyed Hossein Aghamiri^{2,*}, Amir Adibi³ 

¹ Department of Nursing, School of Nursing and Midwifery, Ilam University of Medical sciences, Ilam, Iran

² Department of Neurology, School of Medicine, Imam Hossein Hospital, Shahid Beheshti University of Medical Sciences, Tehran, Iran

³ School of Medicine, Shahid Mostafa Khomaeini Hospital, Ilam University of Medical sciences, Ilam, Iran

*Corresponding author: Department of Neurology, School of Medicine, Imam Hossein Hospital, Shahid Beheshti University of Medical Sciences, Tehran, Iran. Email: hosseinaghamiri60@gmail.com

Received 2024 June 30; Revised 2024 July 28; Accepted 2024 August 3.

Abstract

Background: Premature ejaculation (PE) is a sexual disorder in men with significant prevalence among neurology patients. Epilepsy is one such condition where the prevalence of PE has been notably high.

Objectives: This study aimed to compare the prevalence of PE in men with focal epilepsy and generalized tonic-clonic seizures (GTCS).

Methods: In this cross-sectional observational study, 200 patients with epilepsy were included, with 100 patients having focal epilepsy and 100 patients having GTCS. The study utilized the demographic profile form, Men's Sexual Health Questionnaire (MSHQ), Sexual Quality of Life-Men (SQOL-M), and premature ejaculation diagnostic tool (PEDT). Patients who visited specialized clinics and hospitals and met the inclusion criteria were recruited. Data were then analyzed using SPSS statistical software.

Results: The results showed that the mean (SD) age of patients in the focal epilepsy group was 30.18 (3.85). The mean (SD) scores for the MSHQ, SQOL-M, and PEDT tools were 51.81 (11.98), 34.75 (9.36), and 8.63 (4.79), respectively. Although the mean (SD) PE score in the focal epilepsy group was 9.17 (4.49), higher than that of the GTCS group at 8.09 (5.04), the difference was not statistically significant. Additionally, there was a significant relationship between the PEDT score, SQOL-M score, and MSHQ score ($P < 0.05$). The MSHQ and SQOL-M scores were lower in patients with PE disorder.

Conclusions: Sexual disorders, including PE, are significantly prevalent in both groups of patients with epilepsy. Therefore, it is essential to implement both pharmacological and non-pharmacological interventions to reduce the incidence of PE in these patients.

Keywords: Premature Ejaculation, Focal Epilepsy, Epilepsy

1. Background

Epilepsy is a neurological disorder characterized by spontaneous and recurrent seizures. It affects approximately 3 to 5 percent of populations worldwide, leading to significant psychological and social pressures on patients (1, 2). Repeated and unpredictable seizures result in physical and mental challenges that ultimately diminish the quality of life for those affected (3). The global incidence of epilepsy is estimated at 61.4 per 100,000 person-years, with notably higher rates in low- and middle-income countries (4).

Neurological disorders, including epilepsy, are prevalent and can cause numerous side effects for

patients (5-7). Epilepsy, in particular, can impact sexual function (8, 9). Research on sexual issues in epilepsy patients is limited, but available studies indicate that sexual problems are more common among these patients compared to the general population (9-11). For instance, patients with epilepsy often experience reduced sperm concentration and seminal fluid volume, which can contribute to decreased sexual satisfaction (12).

Among men, various sexual disorders include libido disorders, erectile dysfunction (ED), and ejaculation disorders. Premature ejaculation (PE) is one of the most common sexual disorders in men (13). Due to varying definitions of PE, its prevalence differs across studies. PE

can severely impact a person's sexual quality of life and mental health, particularly when it coexists with ED (14, 15).

Premature ejaculation is generally defined by a short time interval from penetration to ejaculation, occurring either before or shortly after penetration. Some sources define PE as ejaculation occurring within one minute of intercourse, while others consider ejaculation within two minutes as premature. Additionally, PE is characterized by intermittent or consistent orgasm before the desired time (16-18).

Premature ejaculation is categorized into two types. The first type involves men who are physiologically predisposed to rapid peak sexual pleasure due to shorter neural latency. The second type includes cases conditioned by psychological or behavioral factors that result in PE (19).

In diagnosing PE, doctors should consider factors such as age, the novelty of the sexual partner, and the frequency and duration of intercourse, all of which can affect the duration of stimulation (20, 21). The diagnosis of PE is primarily based on the patient's or their partner's history. It has been observed that if a doctor attempts to measure ejaculation time in a clinical setting, it often appears normal or longer than reported by the patient. Therefore, routine clinical evaluation of ejaculation time is not recommended, and diagnosis relies mainly on the time reported by the patient or their partner.

Male sexual dysfunction can result from physical, psychological, or a combination of both causes. Physical causes include chronic conditions such as diabetes, reproductive system tumors, hypertension, urinary infections, and surgeries affecting nerves or organs. Psychological causes may involve psychiatric disorders, the use of psychoactive drugs, anxiety, depression, and other related issues (13, 24, 25).

2. Objectives

Considering the importance of sexual health in men, particularly those with epilepsy, this study aimed to compare PE in men with focal epilepsy and generalized tonic-clonic seizures (GTCS).

3. Methods

In this cross-sectional and observational study, patients with epilepsy were included based on specific entry and exit criteria. The total sample size comprised 200 individuals, with 100 patients having focal epilepsy and 100 patients with GTCS. The patients were evaluated by psychiatrists and neurologists.

Inclusion criteria were: Males aged 18 to 40, married, with at least 6 months of sexual activity, who provided informed consent to participate, and had a diagnosis of epilepsy as determined by the physician. Patients with additional conditions such as mental retardation, a history of sexually transmitted diseases, or a history of drug use were excluded from the study.

The tools used included a demographic profile form, the Men Sexual Health Questionnaire (MSHQ), the Sexual Quality of Life-Men (SQOL-M), and the premature ejaculation diagnostic tool (PEDT).

3.1. Demographic Information Form

It included questions about age, the education level of both the patient and their spouse, smoking history, exercise history, and the number of children.

3.2. Men Sexual Health Questionnaire

This tool includes three dimensions: Erection, ejaculation, and satisfaction. The items on the instrument have a response range from zero to five, with higher scores indicating better sexual performance (26). The MSHQ instrument evaluates dimensions such as erection, which includes three questions with a score range of 0 - 15; ejaculation, which includes seven questions with a score range of 0 - 35; and satisfaction, which includes six questions with a score range of 0 - 30 (26, 27).

3.3. Sexual Quality of Life-Men

This tool consists of 11 questions with scores ranging from "completely agree" (score 1) to "completely disagree" (score 6). The total score ranges from 11 to 66, with higher scores indicating a better quality of life (28, 29).

3.4. Premature Ejaculation Diagnostic Tool

This tool consists of 5 domains, with responses scored on a Likert scale from 0 to 4. Higher scores indicate a greater level of sexual disorder. A score below 8 suggests no PE, a score between 9 and 10 indicates possible PE, and a score above 11 indicates PE (30-32).

In this study, patients with a confirmed diagnosis of epilepsy, approved by a neurologist, were included. Patients who visited specialized clinics and hospitals and met the study's entry criteria were enrolled. Since all the patients were male, the interviews were conducted by a male researcher. Informed written consent was obtained from each patient before starting the questioning, and the study procedures were

explained to them. The collected data were entered into SPSS statistical software with confidentiality maintained, and analyses were performed using independent *t*-tests, ANOVA, regression, and descriptive statistics.

4. Results

The results in Table 1 show that the mean (SD) age of patients in the focal epilepsy group was 30.09 (3.66), in the GTCS group was 30.27 (4.05), and for all patients combined was 30.18 (3.85). Additionally, 15% of all patients had experience with sports, and 15.5% had experience with smoking.

According to Table 2, the mean (SD) score for the MSHQ tool was 51.81 (11.98), for the SQOL-M tool was 34.75 (9.36), and for the PEDT tool was 8.63 (4.79). Although the mean (SD) PE score in the focal epilepsy group was 9.17 (4.49) compared to 8.09 (5.04) in the GTCS group, this difference was not statistically significant.

The findings showed a significant relationship between the PEDT score, the SQOL-M score, and the MSHQ score ($P < 0.05$). The MSHQ score and SQOL-M score were reported to be lower in patients with PE disorder (Table 3).

The results showed that 29.5% of patients were in the definitive PE status, 8.5% were in the possible PE status, and 62% were in the no PE status (Table 4).

5. Discussion

This study aimed to compare PE in men with focal epilepsy to those with GTCS. Sexual problems in both men and women with epilepsy have been investigated in various studies. The results of this study will be compared with findings from other studies across several categories: The general population, patients with epilepsy within Iran, and patients outside Iran.

In a meta-analysis conducted by Ramezani et al. in Iran, 11 articles were reviewed, covering diverse patient groups, including those with heart problems, diabetes, dialysis, neurological issues, and others. The analysis found that the prevalence of ED in men was 56.1% (33). Porst et al. reported the prevalence of PE as 24% in the United States, 20.3% in Germany, and 20% in Italy. Men with PE were also found to be at higher risk for other sexual disorders, such as ED, sexual aversion, and anorgasmia (14). Additionally, Tang and Khoo reported a 46.9% prevalence of PE in a study involving 207 men aged 18 - 70 years (34). These findings are consistent with this study's results on the prevalence of PE in patients with epilepsy.

Nikoobakht et al. included 80 male patients aged 22 to 50, all diagnosed with epilepsy. They found that 42.5% of the patients had ED, and 11.3% had PE. The study also highlighted a significant relationship between seizure type and frequency with erectile function, orgasmic function, and sexual desire (35). Mazdeh et al. examined 35 married patients and found ED in 48.1% of those on sodium valproate and 51.9% of those on carbamazepine. The study also categorized ED severity, noting 11 patients with moderate conditions, 15 with moderate to mild conditions, 16 with mild conditions, and 18 without ED (36). These previous studies, which highlight the presence of sexual disorders among patients with epilepsy in Iran, are consistent with the results of this study.

In the study by Mazdeh et al., which included 80 married women with epilepsy, disorders were reported as follows: 41% for the state of desire, 35% for the state of arousal, 48% for the state of lubrication, and 41% for the state of orgasm (37). These results are consistent with the findings of this study regarding the prevalence of sexual problems in patients with epilepsy.

Henning et al.'s study, which examined 171 patients with epilepsy, found that the most commonly reported issues were erectile problems, decreased libido, vaginal dryness, and orgasm-related issues. The prevalence of sexual problems was 63.3% in men and 75.3% in women. In men, the reported problems included sexual dysfunction (63.3%), delayed ejaculation (5.6%), reduced sexual desire (25.6%), feelings of sexual deviation (5.6%), orgasm difficulties (13.3%), and PE (15.6%) (38).

In Calabrò et al.'s study of 30 men with epilepsy, aged 20 to 53 years, the prevalence rates were as follows: Retarded ejaculation 3.8%, ED 3.3%, PE 6.7%, hyperactive desire 1.7%, hypoactive desire 8.3%, and 21.7% of patients reported no sexual disorders (39). Similarly, Sureka et al. found that 66% of patients reported sexual dysfunction, which included ED (36%), hypoactive sexual desire (0.4%), PE (26%), and 34% of patients reported no sexual dysfunction (40).

One limitation of this study is the potential influence of cultural factors. Conducted in Ilam city, some patients may have been reluctant to discuss sexual problems due to cultural reasons. Therefore, it is recommended to conduct further studies with larger sample sizes.

5.1. Conclusions

The findings indicate a significant prevalence of sexual disorders, including PE, in both groups of patients with epilepsy. Consequently, it is essential to implement both pharmacological and non-

Table 1. Demographic Characteristics of Patients with Epilepsy in the Studied Groups ^a

Variables	Focal Epilepsy	GTCS	Total Epilepsy
Education			
High school or below	49 (49)	47 (47)	96 (48)
University	51 (51)	53 (53)	104 (52)
Economic status			
Weak	65 (65)	54 (54)	119 (59.5)
Moderate	33 (33)	40 (40)	73 (36.5)
Good and excellent	2 (2)	6 (6)	8 (4)
Children			
None	57 (57)	64 (64)	121 (60.5)
One or more	43 (43)	36 (26)	79 (39.5)
Spouse education			
High school or below	38 (38)	47 (47)	85 (42.5)
University	62 (62)	53 (53)	115 (57.5)
Exercise			
Yes	18 (18)	12 (12)	30 (15)
No	82 (82)	88 (88)	170 (85)
Smoking			
Yes	14 (14)	17 (17)	31 (15.5)
No	86 (86)	83 (83)	169 (84.5)
Age	30.09 ± 3.66	30.27 ± 4.05	30.18 ± 3.85

Abbreviation: GTCS, generalized tonic-clonic seizures.

^a Values are expressed as No. (%) or mean ± SD.

Table 2. Status Mean ± SD score of Men's Sexual Health Questionnaire, Sexual Quality of Life-Men and Premature Ejaculation Diagnostic Tool Instruments of the Patients Under Study

Variables	Focal Epilepsy	GTCS	Total Epilepsy	P-Value, F
MSHQ				
Erection	6.6 ± 1.71	6.08 ± 1.27	6.7 ± 1.5	0.04, 4.25
Ejaculation	25.78 ± 4.63	21.57 ± 6.84	23.67 ± 6.19	0.000, 32.37
Satisfaction	22.47 ± 7.33	20.4 ± 7.94	21.43 ± 7.7	0.05, 3.67
MSISQ total	54.85 ± 10.38	48.77 ± 12.72	51.81 ± 11.98	0.007, 7.48
SQOL-M	36.31 ± 10.62	33.2 ± 7.66	34.75 ± 9.36	0.001, 10.47
PEDT	9.17 ± 4.49	8.09 ± 5.04	8.63 ± 4.79	0.76, 0.09

Abbreviations: MSHQ, Men's Sexual Health Questionnaire; SQOL-M, Sexual Quality of Life-Men; PEDT, premature ejaculation diagnostic tool; GTCS, generalized tonic-clonic seizures.

Table 3. Correlation Coefficient Matrix of Men's Sexual Health Questionnaire, Sexual Quality of Life-Men and Premature Ejaculation Diagnostic Tool

Variables	SQOL-M			PEDT			MSHQ		
	P	R	F	P	R	F	P	R	F
MSHQ	0.000	0.28	17.63				1	1	1
SQOL-M	1	1	1	0.03	-0.15	4.65			
PEDT				1	1	1	0.000	0.47	58.74

Abbreviations: MSHQ, Men's Sexual Health Questionnaire; SQOL-M, Sexual Quality of Life-Men; PEDT, premature ejaculation diagnostic tool.

pharmacological interventions to reduce the incidence of PE in these patients.

Acknowledgements

We thank the Ilam University of Medical sciences.

Footnotes

Authors' Contribution: SHA, MO, AD, conceived the study, performed data analysis, and wrote the manuscript; SHA, MO, AD, collected data and wrote the manuscript; SHA, MO, AD, interpreted the results and

Table 4. Investigation of Distribution of Frequency (Percentage) of Premature Ejaculation in Examined Patients^a

Variables	Focal Epilepsy	GTCS	Total Epilepsy
No PE	55 (55)	69 (69)	124 (62)
Possible PE	12 (12)	5 (5)	17 (8.5)
Definitive PE	33 (33)	26 (26)	59 (29.5)
Total	100 (100)	100 (100)	200 (100)

Abbreviation: PE, premature ejaculation; GTCS, generalized tonic-clonic seizures.

^a Values are expressed as No. (%).

wrote the manuscript; SHA, MO, AD, designed the study, wrote, and edited the manuscript.

Conflict of Interests Statement: Authors declared no conflict of interests.

Data Availability: The dataset presented in the study is available on request from the corresponding author during submission or after its publication.

Ethical Approval: The current study was conducted after obtaining approval by the Ethics Committee of Ilam University of Medical Sciences (IR.MEDILAM.REC.1403.027).

Funding/Support: No funding was received.

Informed Consent: Informed consent was obtained.

References

- Longo D, Fauci A, Kasper D, Hauser S, Jameson J, Loscalzo J. *Harrison's Principles of Internal Medicine, 18th Edition*. New York, U.S: McGraw-Hill Education; 2011.
- Kashefimehr N, Ebrahimi SF, Esmkhani AH. [Effectiveness of Acceptance & Commitment Therapy on Intensity of Pain Experience and Life Satisfaction of Patients with Epilepsy]. *J Anesthesiol Pain*. 2023;**14**(1). Persian.
- Shakhatreh L, Foster E, Siriratnam P, Neal A, Carney PW, Jackson GD, et al. Impact of epilepsy surgery on quality of life: Systematic review and meta-analysis. *Epilepsia*. 2023;**64**(7):1709-21. [PubMed ID: 37157209]. <https://doi.org/10.1111/epi.17644>.
- Fiest KM, Sauro KM, Wiebe S, Patten SB, Kwon CS, Dykeman J, et al. Prevalence and incidence of epilepsy: A systematic review and meta-analysis of international studies. *Neurol*. 2017;**88**(3):296-303. [PubMed ID: 27986877]. [PubMed Central ID: PMC5272794]. <https://doi.org/10.1212/WNL.0000000000003509>.
- Mohammadi HR, Erfani A, Jamshidbeigi Y, Rahmatian A, Otaghi M. Effect of using rituximab on disability in patients with multiple sclerosis. *J Med Pharmaceutical Chem Res*. 2024;**6**(12):1854-60. <https://doi.org/10.48309/jmpcr.2024.450019.1158>.
- Hatefi M, Abdi A, Tarjoman A, Borji M. Prevalence of depression and pain among patients with spinal cord injury in Iran: A systematic review and meta-analysis. *Trauma Monthly*. 2019;**24**(4):1-8.
- Vasigh A, Tarjoman A, Soltani B, Borji M. The relationship between mindfulness and self-compassion with perceived pain in migraine patients in Ilam, 2018. *Arch Neurosci*. 2019;**6**(3).
- Abdel-Hamid IA, Abdel-Razek MM, Anis T. Risks factors in premature ejaculation: The neurological risk factor and the local hypersensitivity. In: Jannini EA, McMahon CG, Waldinger MD, editors. *Premature ejaculation: From etiology to diagnosis and treatment*. Milano: Springer; 2013. p. 167-85.
- Yogarajah M, Mula M. Sexual Dysfunction in Epilepsy and the Role of Anti-Epileptic Drugs. *Curr Pharm Des*. 2017;**23**(37):5649-61. [PubMed ID: 28982319]. <https://doi.org/10.2174/1381612823666171004115230>.
- Svalheim S, Tauboll E, Luef G, Lossius A, Rauchenzauner M, Sandvand F, et al. Differential effects of levetiracetam, carbamazepine, and lamotrigine on reproductive endocrine function in adults. *Epilepsy Behav*. 2009;**16**(2):281-7. [PubMed ID: 19716343]. <https://doi.org/10.1016/j.yebeh.2009.07.033>.
- Tao L, Zhang X, Duan Z, Wang Y, Liu J, Hou H, et al. Sexual dysfunction and associated factors in Chinese Han women with epilepsy. *Epilepsy Behav*. 2018;**85**:150-6. [PubMed ID: 29940378]. <https://doi.org/10.1016/j.yebeh.2018.06.003>.
- Hamed SA. The effect of epilepsy and antiepileptic drugs on sexual, reproductive and gonadal health of adults with epilepsy. *Expert Rev Clin Pharmacol*. 2016;**9**(6):807-19. [PubMed ID: 26934627]. <https://doi.org/10.1586/17512433.2016.1160777>.
- Herkommer K, Meissner VH, Dinkel A, Jahnen M, Schiele S, Kron M, et al. Prevalence, lifestyle, and risk factors of erectile dysfunction, premature ejaculation, and low libido in middle-aged men: first results of the Bavarian Men's Health-Study. *Androl*. 2024;**12**(4):801-8. [PubMed ID: 37676020]. <https://doi.org/10.1111/andr.13524>.
- Porst H, Montorsi F, Rosen RC, Gaynor L, Grupe S, Alexander J. The Premature Ejaculation Prevalence and Attitudes (PEPA) survey: prevalence, comorbidities, and professional help-seeking. *Eur Urol*. 2007;**51**(3):816-23. discussion 824. [PubMed ID: 16934919]. <https://doi.org/10.1016/j.eururo.2006.07.004>.
- Verze P, Arcaniolo D, Palmieri A, Cai T, La Rocca R, Franco M, et al. Premature Ejaculation Among Italian Men: Prevalence and Clinical Correlates From an Observational, Non-Interventional, Cross-Sectional, Epidemiological Study (IPER). *Sex Med*. 2018;**6**(3):193-202. [PubMed ID: 29803639]. [PubMed Central ID: PMC6085227]. <https://doi.org/10.1016/j.esxm.2018.04.005>.
- Colonnello E, Sansone A, Zhang H, Zhang Y, Jannini EA. Towards a Universal Definition of Premature Ejaculation. *J Sex Med*. 2022;**19**(12):1717-20. [PubMed ID: 35752558]. <https://doi.org/10.1016/j.jsxm.2022.05.003>.
- Lucas Bustos P, Parron Carreno T, Moreno Gutierrez S, Fernandez Agis I. Progressive arousal: a new concept and definition for premature ejaculation. *Sex Med*. 2023;**11**(2):qfad014. [PubMed ID: 37124144]. [PubMed Central ID: PMC10136197]. <https://doi.org/10.1093/sexmed/qfad014>.
- Rowland DL, Kovi Z, Tamas S, Hevesi K. Do Ejaculation Latency and Other Sexual Measures Differ Between Men With Lifelong and Acquired Premature Ejaculation? *Urol*. 2024;**184**:112-21. [PubMed ID: 37926381]. <https://doi.org/10.1016/j.urology.2023.09.055>.
- Tadayyon F, Mellat M, Khorrami MH, Shahdoost AA, Haghdoost FS. [Comparing the Serum Level of Testosterone and the Relative

- Frequency of Premature Ejaculation in Patients with Varicocele and Normal Population]. *J Isfahan Med Sch*. 2011;**28**(124):2032-8. Persian.
20. Althof SE, McMahon CG, Waldinger MD, Serefoglu EC, Shindel AW, Adaikan PG, et al. An update of the International Society of Sexual Medicine's guidelines for the diagnosis and treatment of premature ejaculation (PE). *J Sex Med*. 2014;**11**(6):1392-422. [PubMed ID: 24848686]. <https://doi.org/10.1111/jsm.12504>.
 21. Rowland DL, Cooper SE. The Tenuous Role of Distress in the Diagnosis of Premature Ejaculation: A Narrative Review. *Sex Med*. 2022;**10**(5):100546. [PubMed ID: 35905650]. [PubMed Central ID: PMC9537272]. <https://doi.org/10.1016/j.esxm.2022.100546>.
 22. Pryor JL, Broderick GA, Ho KF, Jamieson C, Gagnon D. Comparison of estimated versus measured intravaginal ejaculatory latency time in men with and without premature ejaculation. *J Sex Med*. 2005;**3**:54.
 23. Nasiripour S, Farasatinasab M, Khodaverdi S. [Premature Ejaculation Management]. *Razi J Med Sci (RJMS)*. 2017;**23**:65-76. Persian.
 24. Irfan M, Hussain NHN, Noor NM, Mohamed M, Sidi H, Ismail SB. Epidemiology of Male Sexual Dysfunction in Asian and European Regions: A Systematic Review. *Am J Mens Health*. 2020;**14**(4):1557988320937200. [PubMed ID: 32623948]. [PubMed Central ID: PMC7338652]. <https://doi.org/10.1177/1557988320937200>.
 25. Palacios-Cena D, Carrasco-Garrido P, Hernandez-Barrera V, Alonso-Blanco C, Jimenez-Garcia R, Fernandez-de-las-Penas C. Sexual behaviors among older adults in Spain: results from a population-based national sexual health survey. *J Sex Med*. 2012;**9**(1):121-9. [PubMed ID: 22023664]. <https://doi.org/10.1111/j.1743-6109.2011.02511.x>.
 26. Rosen RC, Catania J, Pollack L, Althof S, O'Leary M, Seftel AD. Male Sexual Health Questionnaire (MSHQ): scale development and psychometric validation. *Urol*. 2004;**64**(4):777-82. [PubMed ID: 15491719]. <https://doi.org/10.1016/j.urology.2004.04.056>.
 27. Fakhri A, Morshedi H, Soleymanian A. Psychometric properties of Iranian version of male sexual function index. *Jundishapur Scientific Med J*. 2014;**12**(6):655-63.
 28. Abraham L, Symonds T, Morris MF. Psychometric validation of a sexual quality of life questionnaire for use in men with premature ejaculation or erectile dysfunction. *J Sex Med*. 2008;**5**(3):595-601. [PubMed ID: 18208501]. <https://doi.org/10.1111/j.1743-6109.2007.00749.x>.
 29. Sadat Z, Ghofranipour F, Goshtasebi A, Azin SA. [Validity and reliability of the Persian version of the Sexual Quality of Life-Male questionnaire]. *Payesh (Health Monitor)*. 2017;**16**(1):73-80. Persian.
 30. Pakpour AH, Yekaninejad MS, Nikoobakht MR, Burri A, Fridlund B. Psychometric properties of the Iranian version of the premature ejaculation diagnostic tool. *Sex Med*. 2014;**2**(1):31-40. [PubMed ID: 25356299]. [PubMed Central ID: PMC4184614]. <https://doi.org/10.1002/sm2.21>.
 31. Symonds T, Perelman MA, Althof S, Giuliano F, Martin M, May K, et al. Development and validation of a premature ejaculation diagnostic tool. *Eur Urol*. 2007;**52**(2):565-73. [PubMed ID: 17275165]. <https://doi.org/10.1016/j.eururo.2007.01.028>.
 32. Serefoglu EC, Cimen HI, Ozdemir AT, Symonds T, Bertkas M, Balbay MD. Turkish validation of the premature ejaculation diagnostic tool and its association with intravaginal ejaculatory latency time. *Int J Impot Res*. 2009;**21**(2):139-44. [PubMed ID: 19020523]. <https://doi.org/10.1038/ijir.2008.58>.
 33. Ramezani MA, Ahmadi K, Ghaemmaghami A, Marzabadi EA, Pardakhti F. Epidemiology of Sexual Dysfunction in Iran: A Systematic Review and Meta-analysis. *Int J Prev Med*. 2015;**6**:43. [PubMed ID: 26097672]. [PubMed Central ID: PMC4455123]. <https://doi.org/10.4103/2008-7802.157472>.
 34. Tang WS, Khoo EM. Prevalence and correlates of premature ejaculation in a primary care setting: a preliminary cross-sectional study. *J Sex Med*. 2011;**8**(7):2071-8. [PubMed ID: 21492404]. <https://doi.org/10.1111/j.1743-6109.2011.02280.x>.
 35. Nikoobakht M, Motamedi M, Orandi A, Meysamie A, Emamzadeh A. Sexual dysfunction in epileptic men. *Urol J*. 2007;**4**(2):111-7. [PubMed ID: 17701932].
 36. Mazdeh M, Ghiasian M, Mazaheri S, Faryadras M. [Comparing the effect of sodium valproate and carbamazepine on sexual function satisfaction in men with epilepsy in Hamadan city]. *Pajouhan Scientific J*. 2018;**16**(2):53-8. Persian. <https://doi.org/10.21859/psj.16.2.53>.
 37. Mazdeh M, Taheri M, Ghafouri-Fard S. Investigation of Sexual Satisfaction in Women with Epilepsy and Its Clinical Correlates. *J Mol Neurosci*. 2021;**71**(6):1193-6. [PubMed ID: 33180270]. <https://doi.org/10.1007/s12031-020-01744-z>.
 38. Henning OJ, Nakken KO, Traeen B, Mowinckel P, Lossius M. Sexual problems in people with refractory epilepsy. *Epilepsy Behav*. 2016;**61**:174-9. [PubMed ID: 27371882]. <https://doi.org/10.1016/j.yebeh.2016.05.038>.
 39. Calabro RS, Grisolaghi J, Quattrini F, Bramanti P, Magaouda A. Prevalence and clinical features of sexual dysfunction in male with epilepsy: the first southern Italy hospital-based study. *Int J Neurosci*. 2013;**123**(10):732-7. [PubMed ID: 23627716]. <https://doi.org/10.3109/00207454.2013.798783>.
 40. Sureka RK, Gaur V, Purohit G, Gupta M. Sexual Dysfunction in Male Patients with Idiopathic Generalized Tonic Clonic Seizures. *Ann Indian Acad Neurol*. 2021;**24**(5):726-31. [PubMed ID: 35002131]. [PubMed Central ID: PMC8680921]. https://doi.org/10.4103/aian.AIAN_1023_20.