




Prevalence of Back and Neck Pain in Iranian Dentists: A Systematic Review and Meta-Analysis

Fahimeh Feili¹, Aminollah Vasigh ², Mohamadali Roozegar^{1,*}

¹Oral and Dental Health Research Center, Ilam University of Medical Sciences, Ilam, Iran

²Department of Anesthesiology, School of Allied Medical Sciences, Ilam University of Medical Sciences, Ilam, Iran

*Corresponding Author: Oral and Dental Health Research Center, Ilam University of Medical Sciences, Ilam, Iran. Email: mohamadaliroozegar@gmail.com

Received: 20 October, 2024; Revised: 3 December, 2024; Accepted: 6 December, 2024

Abstract

Context: Musculoskeletal diseases (MSDs) and the pain caused by it are the main cause of loss of work ability, which causes people to leave their jobs and reduce their financial security.

Objectives: This meta-analysis study was conducted with the aim of determining the prevalence of back and neck pain in Iranian dentists.

Data Sources: This meta-analysis study was conducted based on the PARISMA checklist. All articles published in Farsi and English that were conducted in the group of dentists and were published between the years 2000 and 2024 were included in the study. Also, articles that were published in the group of students or assistants or had incomplete data were excluded from the study. Two of the authors of the article reviewed all the searched articles and extracted the related data. It should be noted that all the data were checked and confirmed by the third author of the articles. After entering the data into the researcher-made checklist, data analysis was performed using CMA software.

Results: Results showed, to begin with, 992 articles had been extracted, of which 46 had been removed in the preliminary search. Also, after extra reviews and evaluation of articles, forty-six articles were analyzed. Also, results showed, prevalence of neck pain in Iranian dentists was to 53.3 [CI = 48.5 - 57.9], prevalence of back pain it was to 45.2 [CI = 39.4 - 51.2], prevalence of upper back pain (UBP) in the last 12 months is equal to 39.5 [CI = 31.5 - 48.1] percent, and the prevalence of lower back pain (LBP) was equal to 34.4 [CI = 27.8 - 41.6] percent.

Conclusions: Considering the high prevalence of neck pain and back pain among dentists in Iran, it is necessary to take preventive measures to reduce neck and back pain among dentists.

Keywords: Back Pain, Neck Pain, Dentists

1. Context

Pain may be due to functional disorders or anatomical disorders. One of the causes of functional disorders is musculoskeletal disorders (MSDs) and injuries caused during work and activity. Musculoskeletal disorders are caused by physical activity that affects muscles, tendons, bones, and joints (1, 2). Musculoskeletal disorders differ in severity and include a wide range of disorders and complications, from mild to severe, chronic, and debilitating conditions (3).

Musculoskeletal diseases and the pain caused by it are the main cause of loss of work ability, which causes

people to leave their jobs and reduce their financial security. So that the reduction of work ability, employment of a new employee, hospitalization of a person, and the need to perform diagnostic and treatment measures are among the economic pressures created on dentists. On the other hand, if there is Pain in dentists, it becomes difficult to provide proper service to patients, and as a result, satisfaction with dentists also decreases (4-6).

One of the medical professions that experience MSDs is dentists. The dental profession requires high concentration and accuracy for a long time and creates continuous and significant pressures on the body's organs (7). In fact, these complications are caused by the

long-term and continuous use of dental devices in a limited environment, performing delicate tasks, and being in unfavorable body positions for long hours, which lead to the experience of pain in dentists (8, 9). Dentists experience pain in most parts of the body, including pain related to the spine (10).

Having information about spine injuries and conducting necessary investigations in this field can be effective in preventing chronic diseases (11). The cervical spine has a structure that is responsible for supporting and orienting the head in space and transmitting the force arising from the trunk (4, 12, 13). As a result of injuries caused to the spine, neck and back, it suffers damage (13, 14).

Back pain and neck pain are among the pain experienced by dentists that are caused by MSDs (15, 16). Back pain is one of the common health problems across all ages, which is the main cause of disability and absenteeism from work. Lower back pain (LBP) is one of the main reasons for patients to visit doctors, reasons for hospitalization, patient referrals, and the use of physiotherapy services for patients (5, 17, 18). Chronic neck pain has a prevalence between 16.7% and 75.1% and is considered one of the common causes of disability. On the other hand, due to the increase in the elderly population in Iran, the prevalence of neck and back pain is expected to rise (19).

2. Objectives

Considering the significance of identifying pain in patients, this meta-analysis study was conducted with the aim of determining the prevalence of back and neck pain in Iranian dentists.

3. Data Sources

This meta-analysis study was conducted based on the PRISMA checklist (20).

3.1. Inclusion and Exclusion Criteria

All articles published in Farsi and English that were conducted in the group of dentists and were published between the years 2000 and 2024 were included in the study. Articles that were published in the group of students or assistants or had incomplete data were excluded from the study.

3.2. Information Sources and Study Selection

In this meta-analysis study, which aimed to determine the prevalence of back and neck pain among dentists in Iran, all articles in the international and domestic databases of Iran, including Scopus, PubMed, EBSCO, ISI, SID, Magiran, ScienceDirect, and Iranmedx, were reviewed. The search was performed using keywords such as LBP, back pain, neck pain, pain, dentists, Iran, upper back pain (UBP), and low back pain.

3.3. Data Collection and Statistical Analysis

Two of the authors of the article (associate professor of periodontology and assistant professor of restorative dentistry) reviewed all the searched articles and extracted the related data. It should be noted that all the data were checked and confirmed by the third author of the articles (FF). After entering the data into the researcher-made checklist, data analysis was performed using CMA software.

4. Results

Results showed that initially 992 articles were extracted, of which 46 were removed during the preliminary search (Figure 1). After additional reviews and evaluations, 46 articles were analyzed (Table 1).




Figure 1. Articles included in the study

The results indicated that the prevalence of neck pain in Iranian dentists was 53.3% [CI = 48.5 - 57.9], the prevalence of back pain was 45.2% [CI = 39.4 - 51.2], the prevalence of UBP in the last 12 months was 39.5% [CI = 31.5 - 48.1], and the prevalence of LBP was 34.4% [CI = 27.8 - 41.6] (Figures 2 - 9).

Table 1. Specifications of the Articles

Authors	Years	Place	Age	Questionnaire	N	Back	Upper Back	Lower Back	Neck
Pur Abbas <i>et al.</i> (21)	2004	Tabriz	-	VAS	97	12	-	-	42.2
Ezoddini <i>et al.</i> (22)	2005	Yazd	36.61 (9.57)	VAS	70	-	-	-	47.1
Nasi Saraji <i>et al.</i> (23)	2005	Birjand	36.70	NMQ & REBA	35	60	-	-	54
Sarami <i>et al.</i> (24)	2007	Tehran	36 (5.56)	NMQ & REBA	47	46.8	51.1	-	83
Kardani <i>et al.</i> (25)	2007	Ahvaz	-	Self-administered	172	-	-	-	70
Pargali and Jowkar (26)	2010	Shiraz	37.70 (7.65)	Self-administered	82	33	-	-	28
Ahmadi Motemayel <i>et al.</i> (27)	2012	Hamedan	39.02 (7.22)	NMQ	71	-	23.9	16.9	49.7
Chamani <i>et al.</i> (28)	2012	NMQ	38.2 (7.5)	Kerman	110	-	34	35	59
Choozbineh <i>et al.</i> (29)	2012	Shiraz	39.01 (7.75)	NMQ	160	-	49.2	45.8	55.9
Rabiei <i>et al.</i> (30)	2012	Rasht	39.1 (8.73)	NMQ & RULA	92	35.8	-	-	43
Varmazyar <i>et al.</i> (31)	2012	Qazvin	28.38 (7.37)	NMQ & REBA	63	42.9	-	-	50.8
Aminian <i>et al.</i> (32)	2012	Tehran	35.75 (8.35)	-	210	-	51.3	55.5	66
Asgaripoor <i>et al.</i> (33)	2013	Semnan	38.40 (7.1)	NMQ & REBA	48	77.2	-	-	52.9
Eyvazi <i>et al.</i> (34)	2013	Tabriz	42.12 (8.99)	NMQ	100	38	-	-	48
Memarpour <i>et al.</i> (35)	2013	Shiraz	-	Painful disorders	272	-	30.7	29.9	31.8
Aminian <i>et al.</i> (36)	2013	Tehran	43.19 (9.88)	Nordic	261	54.8	-	-	-
Rahmani <i>et al.</i> (37)	2013	Tehran	41.30 (8.43)	VAS	300	-	-	-	34.7
Ebrahimian <i>et al.</i> (38)	2014	North-Khorasan	41.17	NMQ & REBA	60	46.7	-	-	27
Baroonzade <i>et al.</i> (39)	2014	Hamedan	28.74 (2.74)	NMQ & LUBA	30	43.3	-	-	63.3
Ilbeigi <i>et al.</i> (40)	2014	Mashhad	39.90 (9.7)	NMQ & RULA	80	50	-	-	56.3
Khayati <i>et al.</i> (41)	2014	Tehran	41.65 (6.23)	VAS	70	-	-	-	66.6
Shadmehr <i>et al.</i> (42)	2014	Tehran	-	Nordic	446	-	57	48.2	64.4
Rafie <i>et al.</i> (43)	2015	Kerman	-	NMQ	130	32.5	-	-	55.9
Nadri <i>et al.</i> (44)	2015	Tehran	38.2 (7.1)	VAS	150	-	-	34.5	50
Rahnamaye Tamrooyi <i>et al.</i> (45)	2015	Tehran	-	NMQ & RULA	92	55.4	-	-	78.3
Tirgar <i>et al.</i> (46)	2015	-	43.85 (6.64)	NMQ & RULA	60	-	40	-	83.3
Jalili Nasab <i>et al.</i> (47)	2016	Qazvin	38.8 (7.9)	Nordic	50	-	10	16	24
Seifi <i>et al.</i> (48)	2016	Babol	38.16 (7.43)	NMQ	64	-	-	-	41
Nokhostin and Zafarmand (49)	2016	-	42.18 (9.59)	Self-administered	600	-	-	-	51.87
Zahedpasha <i>et al.</i> (50)	2016	Isfahan	37.8 (8.3)	Nordic	116	43.1	-	-	56.9
Koosha <i>et al.</i> (51)	2016	Tehran	38.28 (6.76)	NMQ & RULA	100	58	-	-	60
Mohseni-Bandpei <i>et al.</i> (52)	2017	Tehran	41.30 (8.43)	VAS	300	31.4	-	-	-
Aliakbari <i>et al.</i> (53)	2018	Bojnourd	-	Nordic	63	50.8	-	-	47.6
Shirinbak <i>et al.</i> (54)	2018	Zanjan	37.1	Nordic	120	24.2	-	-	20.8
Yazdani <i>et al.</i> (55)	2018	Tehran	40.83 (8.2)	Nordic	66	-	49	44	66
Hosseini <i>et al.</i> (56)	2019	Shiraz	35.6 (8.7)	NMQ	136	-	61.6	35.3	37.5
Nadri <i>et al.</i> (44)	2019	Tehran	38.7 (6.6)	VAS	84	44.9	-	-	-
Khandan <i>et al.</i> (57)	2020	Qom	40.06 (9.53)	BPAI	51	-	45.5	34.1	72.7
Karampor <i>et al.</i> (58)	2020	Ilam	39.22 (5.53)	Nordic	18	77.8	-	-	66.7
Shahraki Ebrahimi <i>et al.</i> (59)	2021	Zahedan	37.6 (8.9)	Nordic	102	72.5	-	-	77.5
Hadadi <i>et al.</i> (60)	2021	Bojnourd	38.32 (6.32)	Nordic	28	50	-	-	71.4
Koochak Dezfouli <i>et al.</i> (61)	2022	Sari	43.47 (7.78)	Nordic	90	-	22.2	24.4	43.3
Hosseini and Roshani (62)	2023	Urmia	41.6 (28.51)	Nordic	76	35	-	-	53
Roozegar <i>et al.</i> (63)	2023	Ilam	26.36 (10.77)	Nordic	53	43.4	-	-	52.8
Rahbar <i>et al.</i> (64)	2023	Ahvaz	30.41 (6.22)	Nordic	152	-	59.2	53.3	71.1
Nasiri <i>et al.</i> (65)	2015	Rasht	-	Nordic	99	-	-	-	98.8




Figure 2. Prevalence of upper back pain (UBP) in Iranian dentists




Figure 3. Upper back pain (LBP) funnel plot




Figure 4. Prevalence of lower back pain (LBP) in Iranian dentists




Figure 5. Lower back pain (LBP) funnel plot




Figure 6. Prevalence of back pain in Iranian dentists




Figure 7. Back pain funnel plot




Figure 8. Prevalence of neck pain in Iranian dentists




Figure 9. Neck pain funnel plot

7. Discussion

Pain causes various complications, including disability and reduced life expectancy for patients. For this reason, it is important to identify the prevalence of pain (5, 17, 18, 66). In this study, the prevalence of neck pain in Iranian dentists was 53.3% [CI = 48.5 - 57.9]. In the study of Ohlendorf et al., the prevalence of neck pain related to the last 12 months in 204 Brazilian dentists was 55.4% (67). In the study of Rehan et al. in 270 Pakistani dentists, it was 23% (68), and in Shekhawat's study et al., neck pain was reported as 5.6% "always" and 55% "sometimes" (69). Paying attention to the prevalence of neck pain in dentists and implementing preventive and therapeutic interventions to address it is very important and necessary (70, 71).

Results showed that the prevalence of back pain was 45.2% [CI = 39.4 - 51.2]. In the study of Al-Mohrej et al., conducted on 204 Saudi Arabian dentists (103 men and 101 women), the Prevalence of LBP was 68.1% (72). In the study of Gaowgzeh et al., involving 60 dentists with an average age of 25.7 years, the prevalence of LBP was 70%, with 9.5% of dentists reporting it at an extreme level (73). Similarly, in the study by Ohlendorf et al. on qualified dental assistants, results showed that 86.9% of the surveyed individuals reported back pain (74).

Results showed that the prevalence of UBP in the last 12 months was 39.5% [CI = 31.5 - 48.1], and the prevalence of LBP was 34.4% [CI = 27.8 - 41.6]. In the study of Ohlendorf et al., conducted on 450 German dentists (163 men and 287 women), the prevalence of UBP was 33.3%, and the prevalence of LBP was 45.8% (67). In the study by Akesson et al., among dentists, the prevalence of UBP was 53%, and the prevalence of LBP was 49% (75). In the study by Kumar et al., involving 646 dentists in India, the prevalence of UBP was 18.65%, and the Prevalence of LBP was 72.01% (76).

The strengths of this study include its novelty and the reporting of the status of back pain and neck pain in dentists in Iran. One of the weaknesses of this study is the failure to report all types of pain in dentists. For this reason, it is recommended that another study be conducted to investigate the prevalence of other musculoskeletal pain in dentists.

5.1. Conclusions

Considering the high prevalence of neck pain and back pain among dentists in Iran, it is necessary to implement preventive measures to reduce neck and back pain in this professional group.

Acknowledgements

The authors would like to express their gratitude to Ilam University of Medical Sciences for its valuable contributions to this study.

Footnotes

Authors' Contribution: M. R., F. F., and A. V. conceived the study, performed data analysis, and wrote the manuscript, collected data and wrote the manuscript, interpreted the results and wrote the manuscript, designed the study, wrote, and edited the manuscript.

Conflict of Interests Statement: The authors declared that they have no conflict of interest.

Data Availability: The dataset presented in the study is available upon request from the corresponding author during submission or after publication.

Funding/Support: The authors declared that they have no funding/support.

References

- Marcum J, Adams D. Work-related musculoskeletal disorder surveillance using the Washington state workers' compensation system: Recent declines and patterns by industry, 1999-2013. *Am J Ind Med.* 2017;**60**(5):457-71. [PubMed ID: 28295479]. <https://doi.org/10.1002/ajim.22708>.
- Khalaji H, Yalfani A, Gandomi F. [Evaluation of Musculoskeletal Disorders and the Effect of Ergonomic Interventions on Pain Alleviation and Work Satisfaction among Food Factory Workers]. *J Occupational Hygiene Engin.* 2020;**7**(3):18-26. FA. <https://doi.org/10.52547/johe.7.3.18>.
- Parno A, Sayehmiri K, Nabi Amjad R, Ivanbagha R, Hosseini Ahagh MM, Hosseini Foladi S, et al. Meta-analysis Study of Work-related Musculoskeletal Disorders in Iran. *J Rehabil.* 2020;**21**(2):182-205. <https://doi.org/10.32598/rj.21.2.2444.4>.
- Maroufi N, Ahmadi A, Mousavi Khatir SR. [Comparison of Neck Muscle Activity between Healthy & Chronic Neck Pain Patients Using Electromyography]. *J Mazandaran Univ Med Sci.* 2011;**21**(85):38-46. FA.
- Fatoye F, Gebrye T, Ryan CG, Useh U, Mbada C. Global and regional estimates of clinical and economic burden of low back pain in high-income countries: a systematic review and meta-analysis. *Front Public Health.* 2023;**11**:1098100. [PubMed ID: 37383269]. [PubMed Central ID: PMC10298167]. <https://doi.org/10.3389/fpubh.2023.1098100>.
- Gorasso V, Van der Heyden J, De Pauw R, Pelgrims I, De Clercq EM, De Ridder K, et al. The health and economic burden of musculoskeletal disorders in Belgium from 2013 to 2018. *Popul Health Metr.* 2023;**21**(1):4. [PubMed ID: 37085871]. [PubMed Central ID: PMC10122398]. <https://doi.org/10.1186/s12963-023-00303-z>.
- Kawtharani AA, Chemeisani A, Salman F, Haj Younes A, Msheik A. Neck and Musculoskeletal Pain Among Dentists: A Review of the Literature. *Cureus.* 2023;**15**(1). e33609.
- Aljanakh M. Musculoskeletal disorders among dental assistants: a cross-sectional study. *BMC Musculoskelet Disord.* 2024;**25**(1):64. [PubMed ID: 38218812]. [PubMed Central ID: PMC10787391]. <https://doi.org/10.1186/s12891-024-07178-7>.

9. Mane V, Rajhans N. Assessment of ergonomic risk of work related musculoskeletal disorders among dentists in Kolhapur region. *Work*. 2024;**79**(4):2087-101. [PubMed ID: 38905078]. <https://doi.org/10.3233/WOR-240184>.
10. Pambudi ST, Adiningrat A, Prastyo AN. The Effect of Physical Activity Level on the Low Back Pain Incidence Among Dentists in Yogyakarta City. *Formosa J Sustainable Res*. 2024;**3**(1):145-56. <https://doi.org/10.55927/fjsr.v3i1.7751>.
11. Mohammadi HR, Asadoola Y, Erfani A, Ghoreishi Amin N, Karimiyarandi H, Sadeghi S, et al. Effectiveness of Pulse Intravenous Infusion of Methylprednisolone on Pain in Patients with Lumbar Disc Herniation: A Randomized Controlled Trial. *Anesthesiol Pain Med*. 2024;**14**(4). <https://doi.org/10.5812/aapm-149442>.
12. Grieve GP, Boyling JD, Jull GA. *Grieve's Modern Manual Therapy: The Vertebral Column*. Amsterdam, Netherlands: Elsevier Health Sciences; 2004.
13. Erfani A, Aghamiri SH, Karimi R. Investigating the Role of Hyper Density Signal Length in the Middle Cerebral Artery on the Degree of Disability of Arterial Ischemic Stroke Patients. *Arch Neuroscience*. 2024;**11**(4). <https://doi.org/10.5812/ans-150888>.
14. Mohammadi H, Erfani A, Sadeghi S, Komlakh K, Otaghi M. Investigating Factors Affecting Mortality Due to Spinal Cord Trauma in Patients Admitted to the Intensive Care Unit. *Bulletin of Emergency And Trauma*. 2024;**12**(3). <https://doi.org/10.30476/beat.2024.103079.1517>.
15. Alzahrani AH, Alhusayni AI, Alqahtani B, Alzahrani HG, Almalki HA, Alkhatami KM. Prevalence of Low Back Pain, Disability Among Dentists In Saudi Arabia: A Cross Sectional Study. *J Pioneering Med Sci*. 2024;**13**(2):170-4. <https://doi.org/10.61091/jpms202413225>.
16. Benfaida S, Hachami I, Chafik R, Hamza M, Bennani A. Musculoskeletal Disorders among Dentists in the Private Sector. *Europ J Med Health Sci*. 2024;**6**(1):41-4. <https://doi.org/10.24018/ejmed.2024.6.1.2041>.
17. Botelho J, Machado V, Leira Y, Proenca L, Chambrone L, Mendes JJ. Economic burden of periodontitis in the United States and Europe: An updated estimation. *J Periodontol*. 2022;**93**(3):373-9. [PubMed ID: 34053082]. <https://doi.org/10.1002/JPER.21-0111>.
18. Fatoye F, Gebrye T, Mbada CE, Useh U. Clinical and economic burden of low back pain in low- and middle-income countries: a systematic review. *BMJ Open*. 2023;**13**(4). e064119. [PubMed ID: 37185180]. [PubMed Central ID: PMC10151982]. <https://doi.org/10.1136/bmjopen-2022-064119>.
19. Fejer R, Kyvik KO, Hartvigsen J. The prevalence of neck pain in the world population: a systematic critical review of the literature. *Eur Spine J*. 2006;**15**(6):834-48. [PubMed ID: 15999284]. [PubMed Central ID: PMC3489448]. <https://doi.org/10.1007/s00586-004-0864-4>.
20. Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ*. 2021;**372**:n71. [PubMed ID: 33782057]. [PubMed Central ID: PMC8005924]. <https://doi.org/10.1136/bmj.n71>.
21. Pur Abbas R, Shakuri K, Haji Dizaji R. [Assessment of musculoskeletal pains prevalence and their risk factors in Tabriz dentists]. *Med J Tabriz Univ Med Sci*. 2005;**38**:34000000. FA.
22. Ezoddini Ardakani F, Haerian Ardakani A, AkhavanKarbasi MH, Dehghan Tezerjani K. [Assessment of musculoskeletal disorders prevalence among dentists]. *J Dent Med*. 2005;**17**(4):52-60. FA.
23. Nasl Saraji J, Hosseini MH, Shahtaheri SJ, Golbabaee F, Ghasemkhani M. [Evaluation of ergonomic postures of dental professions by Rapid Entire Body Assessment (REBA), in Birjand, Iran]. *J Dent Med*. 2005;**18**(1):61-7. FA.
24. Sarami M, Lahmi M, Faghihzadeh S. [Ergonomic intervention effect on musculoskeletal disorders Dentists]. *Med Daneshvar*. 2006;**13**:55E+63. FA.
25. Kardani M, Elahi N, Rezaei M. [Assessment of Relationship Between Ergonomic Factors And Neck Pain Among Dentist]. *Sci Med J*. 2007;**6**(1). FA.
26. Pargali N, Jowkar N. Prevalence of musculoskeletal pain among dentists in Shiraz, Southern Iran. *Int J Occup Environ Med*. 2010;**1**(2):69-74. [PubMed ID: 23022788].
27. Ahmadi Motemayel F, Abdolsamadi H, Roshanaei G, Jalilian S. [Prevalence of Musculoskeletal Disorders among Hamadan General Dental Practitioners]. *Avicenna J Clin Med*. 2012;**19**(3):61-6. FA.
28. Chamani G, Zarei MR, Momenzadeh A, Safizadeh H, Rad M, Alahyari A. Prevalence of Musculoskeletal Disorders among Dentists in Kerman, Iran. *J Musculoskeletal Pain*. 2012;**20**(3):202-7. <https://doi.org/10.3109/10582452.2012.704138>.
29. Choobineh AR, Soleimani E, Daneshmandi H, Mohamadbeigi A, Izadi K. Prevalence of musculoskeletal disorders and posture analysis using RULA method in Shiraz general dentists in 2010. *J Islamic Dent Associat Iran*. 2012;**24**(4):244-50.
30. Rabiei M, Shakiba M, Dehghan-Shahreza H, Talebzadeh M. Musculoskeletal disorders in dentists. *Int J Occupational Hygiene*. 2012;**4**(1):36-40.
31. Varmazyar S, Amini M, Kiafar M. Ergonomic Evaluation of Work Conditions in Qazvin Dentists and its Association with Musculoskeletal Disorders Using REBA Method. *J Iran Dent Assoc*. 2012;**24**(4):181-6. eng.
32. Aminian O, Alemohammad ZB, Sadeghniaat-Haghighi K. Musculoskeletal disorders in female dentists and pharmacists: a cross-sectional study. *Acta Medica Iranica*. 2012;635-40.
33. Askaripoor T, Kermani A, Jandaghi J, Farivar F. Survey of musculoskeletal disorders and ergonomic risk factors among dentists and providing control measures in Semnan. *J Health Hygiene*. 2013;**4**(3):241-8. FA.
34. Eyvazi M, Rezaei M, EterafOskoyi M, Zolghadr M, Parandavar H, Sadeghi N, et al. [Risk factors and prevalence of musculoskeletal disorders among dentists]. *Med J Tabriz Univ Med Sci*. 2013;**34**:81000000. FA.
35. Memarpour M, Badakhsh S, Khosroshahi SS, Vossoughi M. Work-related musculoskeletal disorders among Iranian dentists. *Work*. 2013;**45**(4):465-74. [PubMed ID: 22976156]. <https://doi.org/10.3233/WOR-2012-1468>.
36. Aminian O, Banafsheh Alemohammad Z, Sadeghniaat Haghighi K. [Comparative assessment of low back pain and its determinants among Iranian male general dentists and pharmacists]. *J Den Med*. 2013;**26**(2):108-14. FA.
37. Rahmani N, Amiri M, Mohseni-Bandpei MA, Mohsenifar H, Pourahmadi MR. Work related neck pain in Iranian dentists: an epidemiological study. *J Back Musculoskelet Rehabil*. 2013;**26**(1):9-15. [PubMed ID: 23411643]. <https://doi.org/10.3233/BMR-2012-0343>.
38. Ebrahimian H, Hokmabadi R, Shoja E. [Evaluation of ergonomic postures of dental professions by rapid entire body assessment (REBA) in North Khorasan, Iran]. *North Khorasan Univ Med Sci*. 2014;**5**(5):961-7. FA. <https://doi.org/10.29252/jnkums.5.5.961>.
39. Baroonyzade Z, Motamedzade M, Golmohammadi R, Kasraei S, Faradmal J. [Assessment of Postural Load Index Using LUBA Method and the Prevalence of Musculoskeletal Disorders in Dentists]. *J Occupational Hygiene Engin*. 2014;**1**(2):27-36. FA.
40. Ilbeigi S, Biglar A, Sagheb joo M, Farzaneh H. [The investigation of relationship between work-related musculoskeletal disorders and level of physical activity and body posture of dentists in Mashhad city in 2012-2013]. *J Torbat Heydariyeh Univ Med Sci*. 2015;**2**(4):37-1. FA.
41. Khayati F, Nasr Esfahani M, Firoozeh M, Kavooosi A, Saremi M. [Predictive factors of neck pain in dentists]. *Iran J Ergonomics*. 2006;**13**:55E+63. FA.

- 2014;**2**(3):24-32. FA.
42. Shadmehr A, Haddad O, Azarnia S, Sanamlo Z. Disorders of the Musculoskeletal System among Tehran, Iranian Dentists. *J Musculoskeletal Pain*. 2014;**22**(3):256-9. <https://doi.org/10.3109/10582452.2014.883022>.
 43. Rafie F, Zamani Jam A, Shahravan A, Raouf M, Eskandarizadeh A. Prevalence of Upper Extremity Musculoskeletal Disorders in Dentists: Symptoms and Risk Factors. *J Environ Public Health*. 2015;**2015**:517346. [PubMed ID: 26064141]. [PubMed Central ID: PMC4433686]. <https://doi.org/10.1155/2015/517346>.
 44. Nadri H, Rohani B, Teimori G, Vosoughi S, Fasih-Ramandi F. Thoracic Kyphosis Angle in Relation to Low Back Pain among Dentists in Iran. *Open Access Maced J Med Sci*. 2019;**7**(21):3704-9. [PubMed ID: 32010402]. [PubMed Central ID: PMC6986497]. <https://doi.org/10.3889/oamjms.2019.578>.
 45. Rahnamaye Tamrooyi F, Javar MA, Salimi S, Mohammadpour H, Avakh A, Faizollahi S. [A survey on Prevalence of Musculoskeletal Disorders in Dentists of Tehran and their posture assessment by RULA method]. *Int Res J Appl Basic Sci*. 2015;**9**(5):666-71. FA.
 46. Tirgar A, Javanshir K, Talebian A, Amini F, Parhiz A. Musculoskeletal disorders among a group of Iranian general dental practitioners. *J Back Musculoskelet Rehabil*. 2015;**28**(4):755-9. [PubMed ID: 25547232]. <https://doi.org/10.3233/BMR-140579>.
 47. Jalili Nasab A, Azmoodeh F, Alipour M, Ansari S, Varmazyar S. [Investigation the prevalence of musculoskeletal disorders and its related factors in general dentists of Qazvin in 2016]. *J Res Dent Sci*. 2018;**14**(4):220-7. FA.
 48. Seifi S, Eftekharian S, Sarrafan N, Gholinia H. [Ergonomic evaluation of frequency and risk factors of musculoskeletal disorder of specialist dentists of Babol dentistry faculty]. *Stud Med Sci*. 2016;**27**(4):330-5. FA.
 49. Nokhostin MR, Zafarmand AH. "Musculoskeletal problem": Its prevalence among Iranian dentists. *J Int Soc Prev Community Dent*. 2016;**6**(Suppl 1):S41-6. [PubMed ID: 27195226]. [PubMed Central ID: PMC4863482]. <https://doi.org/10.4103/2231-0762.181166>.
 50. Zahedpasha S, Ebrahimipour S, Sharifzadeh G, Rashed-Mohassel A, Mortaheb A. [Prevalence of musculoskeletal disorders among Iranian dentists]. *J Health Sci Techno*. 2017;**1**(2):70-4. FA.
 51. Koosha S, Bidgoli MK, Raouf A, Ezatian R. [Investigation of musculoskeletal disorders and its related factors in dentists by REBA method among dental clinics faculties in Tehran in 2014]. *J Dent Med*. 2016;**29**(2):116-28. FA.
 52. Mohseni-Bandpei MA, Rahmani N, Halimi F, Farooq MN. The prevalence of low back pain in Iranian dentists: An epidemiological study. *Pak J Med Sci*. 2017;**33**(2):280-4. [PubMed ID: 28523022]. [PubMed Central ID: PMC5432689]. <https://doi.org/10.12669/pjms.332.11519>.
 53. Aliakbari R, Vahedian-Shahroodi M, Tehrani H, Esmaeili H, Hokmabadi R. [Dentists' ergonomic assessment by RULA method and its relationship with musculoskeletal disorders]. *J Dent Med*. 2018;**31**(1):42-8. FA.
 54. Shirinbak I, Basir Shabestari S, Sefidi F. [Prevalence of Musculoskeletal pain and its Related Factors among Zanjan dentists]. *J Res Dent Sci*. 2018;**15**(1):42-8. FA. <https://doi.org/10.29252/jrds.15.1.42>.
 55. Yazdani M, Moradi K, Sobhani V, Akbari H, Kazempour M, Shamsoddini A, et al. Assessing the Prevalence of Musculoskeletal Disorders and Posture Conditions during Work Time of Dentists in a Dentistry Clinic affiliated to a Military College in Tehran in 2017. *J Military Med*. 2022;**20**(2):222-30.
 56. Hosseini A, Choobineh A, Razeghi M, Pakshir HR, Ghaem H, Vojud M. Ergonomic Assessment of Exposure to Musculoskeletal Disorders Risk Factors among Dentists of Shiraz, Iran. *J Dent (Shiraz)*. 2019;**20**(1):53-60. [PubMed ID: 30937338]. [PubMed Central ID: PMC6421327]. <https://doi.org/10.30476/DENTJODS.2019.44564>.
 57. Khandan M, Koohpaei A, Shahbazi M, Allahdadi Z, Abdi zarin S. Assessment of Individual and Occupational Risk Factors of Musculoskeletal Disorders Using BPAI among Dentists in Qom, Iran. *Arch Hygiene Sci*. 2020;**9**(3):234-45. <https://doi.org/10.29252/ArchHygSci.9.3.234>.
 58. Karapor AM, Kakaei H, Zargosh Z, Hajizadeh H, Bekri Zadeh H, Hasani AA. [Investigation of Musculoskeletal Disorders and Evaluation of Physical Condition while Working with RULA Method in Dentists of Ilam Health Centers in 2011]. *Occupational Hygiene Health Promotion*. 2020;**4**(3). FA. <https://doi.org/10.18502/ohhp.v4i3.4661>.
 59. Shahraki Ebrahimi H, Aghaei Soltani F, Hosseinfar M, Maserat V, Saeedi Robot M, Hojati J. Prevalence of Musculoskeletal Pains and Disability among Dentists in Zahedan. *J Dentomaxillofacial Radiol, Pathol Surg*. 2021;**10**(2).
 60. Hadadi J, Taghipour M, Ghorbanpour A. [Evaluation of the Prevalence of Musculoskeletal Disorders among Dentists in Bojnourd, 2018]. *J Health Res Commun*. 2021;**7**(2):52-60. FA.
 61. Koochak Dezfouli M, Bagheri B, Yazdani Charati J, Zamanzadeh M. [Prevalence of musculoskeletal disorders and related risk factors among general dentists in Sari in 2019]. *J Mashhad Dent Sch*. 2021;**45**(4):395-404. FA.
 62. Hosseini N, Roshani S. [Comparison of posture and musculoskeletal disorders of dentists with high and low work experience]. *J Dent Med*. 2023;**36**:27-36. FA.
 63. Roozegar MA, Shafiee E, Havasian MR. [Evaluation of prevalence of musculoskeletal disorders in general dentists in Ilam city in 2020]. *Daneshvar Med*. 2023;**30**(6):70-7. FA. <https://doi.org/10.22070/daneshmed.2022.16951.1290>.
 64. Rahbar P, Shirkosh P, Tehrani S. Assessment of ergonomic condition of Iranian dentists using RULA technique and its correlation with musculoskeletal disorders. *J Population Therapeutics Clin Pharmacol*. 2023;**30**(9). <https://doi.org/10.47750/jptpc.2023.30.09.009>.
 65. Nasiri E, Azarnia S, Shabani S, Kayani P. Examination of musculoskeletal disorders in dentists. *Dr. Hakimzadeh's First National Aging Conference*. Guilan, Iran. Guilan University of Medical Sciences; 2015.
 66. Mohammadi HR, Erfani A, Jamshidbeigi Y, Rahmatian A, Otaghi M. Effect of using rituximab on disability in patients with multiple sclerosis. *J Med Pharmaceutical Chem Res*. 2024;**6**(12):1854-60. <https://doi.org/10.48309/jmpcr.2024.450019.1158>.
 67. Ohlendorf D, Naser A, Haas Y, Haanel J, Fraeulin L, Holzgreve F, et al. Prevalence of Musculoskeletal Disorders among Dentists and Dental Students in Germany. *Int J Environ Res Public Health*. 2020;**17**(23). [PubMed ID: 33255491]. [PubMed Central ID: PMC7727829]. <https://doi.org/10.3390/ijerph17238740>.
 68. Rehan F, Leghari MA, Memon MS, Atif M, Masood A, Mohsin F, et al. Frequency of musculoskeletal discomfort among dentists in karachi, pakistan. *Pakistan Oral Dent J*. 2015;**35**(4).
 69. Shekhawat KS, Chauhan A, Sakthidevi S, Nimbeni B, Golai S, Stephen L. Work-related musculoskeletal pain and its self-reported impact among practicing dentists in Puducherry, India. *Indian J Dent Res*. 2020;**31**(3):354-7. [PubMed ID: 32769266]. https://doi.org/10.4103/ijdr.IJDR_352_18.
 70. Letafatkar A, Rabiei P, Alamooti G, Bertozzi L, Farivar N, Afshari M. Effect of therapeutic exercise routine on pain, disability, posture, and health status in dentists with chronic neck pain: a randomized controlled trial. *Int Arch Occup Environ Health*. 2020;**93**(3):281-90. [PubMed ID: 31654125]. <https://doi.org/10.1007/s00420-019-01480-x>.
 71. Kim ES, Jo ED, Han GS. Effects of stretching intervention on musculoskeletal pain in dental professionals. *J Occup Health*.

- 2023;**65**(1). e12413. [PubMed ID: 37347801]. [PubMed Central ID: PMC10287045]. <https://doi.org/10.1002/1348-9585.12413>.
72. Al-Mohrej OA, AlShaalan NS, Al-Bani WM, Masuadi EM, Almodaimegh HS. Prevalence of musculoskeletal pain of the neck, upper extremities and lower back among dental practitioners working in Riyadh, Saudi Arabia: a cross-sectional study. *BMJ Open*. 2016;**6**(6). e011100. [PubMed ID: 27324712]. [PubMed Central ID: PMC4916616]. <https://doi.org/10.1136/bmjopen-2016-011100>.
73. Gaowgzeh RA, Chevidikunnan MF, Al Saif A, El-Gendy S, Karrouf G, Al Senany S. Prevalence of and risk factors for low back pain among dentists. *J Phys Ther Sci*. 2015;**27**(9):2803-6. [PubMed ID: 26504297]. [PubMed Central ID: PMC4616098]. <https://doi.org/10.1589/jpts.27.2803>.
74. Ohlendorf D, Haas Y, Naser A, Haenel J, Maltry L, Holzgreve F, et al. Prevalence of Muscular Skeletal Disorders among Qualified Dental Assistants. *Int J Environ Res Public Health*. 2020;**17**(10). [PubMed ID: 32429484]. [PubMed Central ID: PMC7277800]. <https://doi.org/10.3390/ijerph17103490>.
75. Akesson I, Balogh I, Hansson GA. Physical workload in neck, shoulders and wrists/hands in dental hygienists during a work-day. *Appl Ergon*. 2012;**43**(4):803-11. [PubMed ID: 22208356]. <https://doi.org/10.1016/j.apergo.2011.12.001>.
76. Kumar VK, Kumar SP, Baliga MR. Prevalence of work-related musculoskeletal complaints among dentists in India: a national cross-sectional survey. *Indian J Dent Res*. 2013;**24**(4):428-38. [PubMed ID: 24047834]. <https://doi.org/10.4103/0970-9290.118387>.