Published Online: 2025 June 25

Research Article



The Impact of Social-Related Factors on the Severity of Prolonged Grief Symptoms: A Machine Learning Approach

Mohammad Nikbakht (b)¹, Fatemeh Serjouie (b)^{2,*}, Amir Sam Kianimoghadam (b)³, Zeinab Farjampanah⁴

¹ Faculty of Industrial Engineering, Iran University of Science and Technology, Tehran, Iran

² Department of Clinical Psychology, School of Medicine, Shahid Beheshti University of Medical Science, Tehran, Iran

³ Department of Clinical Psychology, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

⁴ Department of Clinical Psychology, School of Medicine, Kermanshah University of Medical Sciences, Kermanshah, Iran

*Corresponding Author: Department of Clinical Psychology, School of Medicine, Shahid Beheshti University of Medical Science, Tehran, Iran. Email: fatemeh.ser34@gmail.com

Received: 31 May, 2025; Accepted: 31 May, 2025

Abstract

Background: The aim of this study is to use a machine learning (ML) approach to investigate the relationship between social dimensions of childhood trauma and attachment style with the severity of prolonged grief symptoms.

Methods: This descriptive-correlational study was conducted on 516 bereaved individuals who were selected through convenience sampling. Participants completed the Prolonged Grief Scale (PG-R-13), Childhood Trauma (CTQ), and Attachment Styles (RAAS) Questionnaires with informed consent, and instead of using an independent questionnaire to measure social dimensions, items related to these dimensions were extracted from the questionnaires. Machine learning models, including linear regression, random forest, and gradient boosting, were used to analyze the data to predict the severity of prolonged grief symptoms.

Results: The results showed that social dimensions related to childhood trauma were the strongest predictors of the severity of prolonged grief symptoms. The random forest model showed the highest predictive power (60.6% accuracy). It was also found that perceived social support was associated with a reduction in the severity of grief symptoms.

Conclusions: This study demonstrated the role of social dimensions of psychological factors in prolonged grief symptoms. The use of ML provided a novel approach to uncovering hidden patterns in the grief-related process that could lead to personalized intervention.

Keywords: Prolonged Grief Disorder, Attachment Style, Childhood Trauma, Machine Learning, Social Interaction

1. Background

A significant number of bereaved adults experience intense and persistent grief that leads to functional impairment in social, occupational, and emotional domains. In the DSM-5-TR, prolonged grief disorder has been added to the category of trauma- and stress-related disorders. Adults who experience prolonged grief suffer from symptoms of intense longing for the deceased, preoccupation with memories of the deceased, emotional pain, identity disturbance, disbelief about the death, intense loneliness, emotional numbness, avoidance of recollection, and difficulty reintegrating themselves into activities (1). Given that the prevalence of prolonged grief disorder in bereaved adults is 13% (2). It is important to identify factors affecting the prolongation of grief (3).

Previous studies have shown that psychological factors, including attachment style and childhood trauma, affect grief severity. Childhood trauma involves emotional distress due to uncontrollable stressful events like death (4). It occurs when children face uncontrollable, stressful events such as physical, emotional, or sexual abuse or neglect (5). Childhood

Copyright @ 2025, Applied Psychology for Health Promotion. This open-access article is available under the Creative Commons Attribution-NonCommercial 4.0 (CC BY-NC 4.0) International License (https://creativecommons.org/licenses/by-nc/4.0/), which allows for the copying and redistribution of the material only for noncommercial purposes, provided that the original work is properly cited.

How to Cite: Nikbakht M, Serjouie F, Kianimoghadam A S, Farjampanah Z. The Impact of Social-Related Factors on the Severity of Prolonged Grief Symptoms: A Machine Learning Approach. Appl Psychol Health Promot. 2025; 2 (1): e163387. https://doi.org/10.5812/aphp-163387.

trauma disrupts parent-child bonds, leading to insecure attachment and increased vulnerability to prolonged grief in adulthood (6).

Another factor that affects social interactions related to prolonged grief disorder is attachment style. Attachment theory states children form an internal model of relationships through interactions with their primary caregivers (7). Attachment theory provides principles that allow understanding of different attachment styles and their impact on the response to grief. Individuals with various attachment styles exhibit unique grieving processes (8). Different attachment styles have different manifestations in social interactions, which makes it a risk factor for developing prolonged grief disorder when experiencing grief with an insecure attachment style. Individuals with an anxious attachment style have a high preoccupation and longing for the deceased, which leads to a lack of acceptance of the deceased's death (9). This attachment style leads to excessive reassurance in social relationships. Individuals with an avoidant attachment style suppress their feelings about their grief, which leads to prolonged and unresolved grief. They also avoid receiving social support after experiencing grief, which intensifies their grief. Therefore, insecure attachment styles lead to different manifestations of social interactions that make individuals susceptible to prolonged grief symptoms (10).

Previous studies have shown a correlation between social interactions and symptoms of prolonged grief. In grief, social interactions are associated with perceived social support, and the higher the perceived social support, the lower the symptoms of prolonged grief (11, 12). For avoidant interactions, quality matters more than quantity. A 2024 study found that the amount of eye contact was negatively correlated with symptoms of prolonged grief. Social withdrawal is a strong predictor of developing prolonged grief disorders (13, 14).

Based on prior research findings, previous studies have measured the influence of psychological factors such as childhood trauma and attachment style on prolonged grief symptoms, but few studies have examined these factors simultaneously, and it is unclear how common social dimensions of these factors are related to prolonged grief symptoms. In addition, most previous studies have used traditional statistical analysis methods to examine these relationships, which assume linearity, while psychological processes are inherently multifaceted and need to be examined using more advanced approaches. One of these approaches is machine learning)ML), which enables pattern recognition in dimensional data to identify latent patterns and complex variable relationships and helps to uncover nonlinear relationships of hidden patterns between variables. Linear regression is a supervised ML model used to predict a dependent (continuous) variable based on one or more independent variables. The model assumes that the relationship between the independent and dependent variables is linear. In other words, the dependent variable is modeled as a linear combination of the independent variables and their coefficients.

2. Objectives

The aim of the present study is to investigate the relationship between social dimensions of childhood trauma and attachment style with the severity of prolonged grief symptoms, which has been investigated through ML. Using ML models, this study seeks to uncover hidden patterns in grief-related mechanisms to provide a more accurate and data-driven understanding of the severity of grief symptoms and to provide clinical interventions. To this end, instead of employing a separate social interaction scale, relevant items were extracted from psychological measures.

3. Methods

3.1. Participants

The present study is a descriptive correlational design based on ML. The statistical population consisted of bereaved adults, from whom 516 individuals were selected through convenience sampling in Iran. Inclusion criteria included participants aged between 18 and 80 years, positive responses to questions 1 and 13 of the the Prolonged Grief Scale (PG-R-13) Questionnaire regarding functional occupation, and at least 12 months had passed since the death. Exclusion criteria included incomplete and patterned completion of the questionnaire.

3.2. Instruments

The Childhood Trauma Questionnaire (CTQ), developed by Bernstein et al., includes 28 items across five subscales: Sexual, physical, and emotional abuse, and emotional and physical neglect. Items are rated on a 5-point Likert scale. Bernstein et al. reported Cronbach's alpha values of 0.87, 0.86, 0.95, 0.89, and 0.78 (15). Concurrent validity with therapists' ratings ranged from 0.59 to 0.78. Ebrahimi et al. (16)found alpha values of 0.81 - 0.98, while the present study reported 0.74 - 0.85.

3.3. The Adult Attachment Style Questionnaire

The Adult Attachment Style Questionnaire (RAAS) by Collins and Reed comprises 18 items rated on a 5-point scale, based on three attachment styles. Reported alpha values were 0.75, 0.72, and 0.69 (17). The scale was validated in Iran by Hajializadeh et al. (18), with testretest reliabilities of 0.57, 0.75, and 0.47.

3.4. The Prolonged Grief Questionnaire

by Prigerson et al. measures prolonged grief symptoms as a unidimensional construct. Reported internal consistency values were 0.83 (Yale), 0.90 (Utrecht), and 0.93 (Oxford) (19). In Iran, Ashouri et al. (20)reported Omega reliability of 0.93 and test-retest reliability (6 weeks) of 0.89. Validity was supported through significant correlations with depression, PTSD, functional impairment, and hope.

3.5. Procedure

The authors followed the ethical considerations of the study according to the Declaration of Helsinki (1975) as revised in 2008. Informed written consent was obtained from all participants. Participants were then given a link to complete the study questionnaires via Google Forms. Participants were asked to complete the PG-13-R, CTQ, RAAS, RFQ, and SSGS questionnaires. To examine the social interactions, dimensions were estimated by identifying items related to relationships in the attachment style and childhood trauma variables, and no independent questionnaire was used to examine social interactions. Relevant items were selected from each questionnaire, which are items 2, 5, 7, 13, 19, and 28 from the Childhood Trauma Questionnaire; items 1, 12, 14, and 16 from the Attachment Styles Questionnaire; and G2, G3, G8, and G12 from Prolonged Grief Questionnaire. The criterion for selecting the social interactions variable was selected based on the number and quality of interpersonal relationships in the independent variable items. Descriptive statistics and ML models were used to analyze the data. The Pandas and NumPy libraries were used for data management and processing, Matplotlib and Seaborn for

visualization, SciPy for statistical analysis, Scikit-learn for ML modeling (including random forest and gradient boosting) and model performance evaluation, and Statsmodels for advanced statistical analysis.

4. Results

Descriptive data showed that 87.79% of the participants were female, 56.20% were single, and most had a bachelor's or master's degree. Correlation analyses showed that childhood social support variables (T2, T5, T7, T13, T19, T28) were negatively associated with symptoms of bereavement such as longing, difficulty in daily functioning, and impairment. Similarly, social skills from the Attachment Style Questionnaire (A1, A12, A14) were associated with reduced functioning problems. Conversely, distrust (A16) showed weaker but negative correlations, indicating increased vulnerability to functional impairment (Table 1 and Figure 1).

Three models were applied to predict grief severity: Linear regression to assess the impact of independent variables, a random forest for prediction, and Gradient Boosting to enhance accuracy. Among them, the most accurate model identified childhood trauma and insecure attachment as key predictors (Table 2 and Figure 2).

5. Discussion

The present study aimed to investigate the relationship between the social dimensions of childhood trauma and attachment style with the severity of prolonged grief symptoms, which are investigated through ML. The findings showed that the greatest predictor of the severity of prolonged grief symptoms was related to social interactions based on childhood trauma, and then it was related to insecure attachment of the avoidant type. The predictive power of the model was estimated at Random Forest 60.6%, which indicates a strong relationship. Also, ML methods were used to analyze the data and the factors predicting the severity of prolonged grief symptoms. As a result, this finding provided accurate insight into the factors associated with prolonged grief disorder.

5.1. Comparison with Previous Studies

The findings of the present study are consistent with previous studies and showed that social interactions significantly reduce vulnerability to prolonged grief symptoms. A study on 539 bereaved individuals showed

Table 1. Correlation Analysis				
Variables 1	Variables 2	Correlation Coefficient	P-Value	
T2	G2	-0.07	0.10375	
T2	G3	-0.05	0.22606	
T2	G12	-0.11	0.0122	
T5	G2	-0.11	0.01503	
T5	G8	-0.09	0.03667	
T5	G12	-0.10	0.02474	
T7	G2	-0.15	0.00051	
T7	G12	-0.12	0.51528	
T13	G2	-0.13	0.00304	
T13	G12	-0.15	0.05362	
T19	G2	-0.13	0.00286	
T19	G12	-0.10	0.02881	
T28	G2	-0.13	0.00222	
T28	G12	-0.15	0.00252	
A1	G3	-0.09	0.05292	
A1	G12	-0.14	0.00178	
A12	G3	-0.08	0.07653	
A12	G12	-0.13	0.00301	
A14	G3	-0.17	0.00011	
A14	G12	-0.19	1e-05	
A16	G3	-0.04	0.41647	
A16	G12	-0.14	0.0018	

that perceived social interactions are related to prolonged grief symptoms in that social interactions increase self-compassion in individuals and, as a result, reduce vulnerability to prolonged grief disorder (21). Another study showed that a decrease in social interactions, which is characterized by the frequency and negative perception of social relationships, causes social isolation (22). Therefore, the bereaved individual does not receive social support after grief, and grief symptoms become more severe and prolonged. Studies have shown inconsistent results with the hypothesis of the present study and demonstrated that in bereaved individuals who experience anxiety, social interactions do not play a reducing role in the anxiety caused by their grief symptoms (21). Another study also considered the role of social interactions conditional on emotional engagement with a supportive source in order to be able to reduce the severity of grief symptoms, and if social interactions are not based on emotional experience, they do not have a reducing effect on prolonged grief symptoms (22).

The findings of the present study are consistent with previous studies, showing that the social aspect of childhood trauma leads to an exacerbation of symptoms of prolonged grief (23). A meta-analysis showed that childhood trauma is associated with lifelong psychological disorders. In particular, disadvantages socioeconomic and inequalities predispose individuals to experience psychological disorders with grief (24). A study on grief due to the death of a spouse showed that childhood maltreatment, combined with poor social status, predicts symptoms of prolonged grief. Another study showed that bereaved individuals due to the death of a parent in childhood become socially withdrawn and exacerbate prolonged grief (25).

The hypothesis of the present study can be explained as follows: The aspect of social interactions resulting from childhood trauma causes individuals to cognitively perceive others as untrustworthy or rejecting, and as a result, to have a negative interpretation of social interactions. This cognitive bias causes social isolation, which intensifies the symptoms of prolonged grief (26, 27). Childhood trauma also leads to emotional dysregulation, and as a result, the grieving person has a disorder in integrating his loss into life and therefore does not process the loss and does not experience his emotions properly, which leads to an



Table 2. Results of Models					
Model	Accuracy	Diagnosis of Severe Grief (Recall - 1)	F1-Scores	Precision	
Linear regression (%)	57.7	75	62	53	
Random forest (%)	60.6	69	62	56	
Gradient boosting (%)	54.8	65	57	51	

exacerbation of the symptoms of prolonged grief (28, 29).

The findings of the present study are consistent with those from previous studies. A study of 157 bereaved individuals showed that avoidant attachment style



Comparison of percision, recall, and F1-Score

Figure 2. Comparison of precision, recall, and F1-score

moderates the effect of perceived social support on grief with individuals symptoms because avoidant attachment style avoid social support due to distress from emotional intimacy and therefore do not experience natural grief, and on the contrary, individuals with secure attachment style use social interactions as their support network to process grief (30). A study inconsistent with the findings of the present study showed that insecure attachment style predicts prolonged grief symptoms cross-sectionally and does not predict grief symptoms longitudinally and 12 months after death. A comparison of these two studies shows differences in social context (10). The attachment model of complex grief suggests that internalized working models regulate grief by integrating the loss into the individual's life narrative, and social interactions facilitate the integration of the loss by examining the internal representation of the

deceased (31). Secure attachment allows the grieving person to reactivate their exploration system and go through a natural grieving process. However, individuals with an avoidant attachment style have internalized emotional distance from important individuals, and after the loss, they use emotional suppression as an emotion regulation strategy to stay away from social support and experience emotions related to grief (32). These cases lead to unfinished grief in them. In summary, secure attachment helps social interactions in grief processing, and individuals with an insecure-avoidant attachment style do not trust others enough for social support, and this intensifies their grief (33).

5.2. Clinical Implications

The findings offer significant clinical value for psychologists, psychiatrists, and social workers dealing

with grief and crisis intervention. Children and adults bereaved by suicide are vulnerable to prolonged grief, and social interaction factors — particularly perceived social support — play a critical role. This study highlights the impact of social history on grief symptom severity.

5.3. Limitations and Future Directions

The cross-sectional design limits causal interpretation; longitudinal studies are recommended. Data were gathered via retrospective self-reports, making recall bias likely; future research should include interviews. The study measured limited dimensions of social interaction; broader aspects should be addressed in future work. It is also suggested to use deep neural models to analyze data and assess the effect of treatment on bereaved individuals' interpersonal relationships.

5.4. Conclusions

Overall, the present study measured the important role of the social interaction dimensions of childhood trauma and insecure attachment style in the severity of prolonged grief symptoms using a ML approach. The use of ML provides new insights into these complex interactions and paves the way for more personalized grief interventions. Future research could address psychological interventions based on this method to improve prolonged grief disorder.

Footnotes

Authors' Contribution: Study concept and design: M. N.; Acquisition of data: Z. F.; Analysis and interpretation of data: M. N.; Drafting of the manuscript: F. S.; Critical revision of the manuscript for important intellectual content: M. N.; Statistical analysis: M. N.; Administrative, technical, and material support: F. S.; Study supervision: A. K.

Conflict of Interests Statement: All authors declare that they have no conflict of interests.

Data Availability: The dataset utilized in the study can be obtained upon request from the corresponding author during submission or following publication. The data is not publicly accessible due to privacy and ethical. **Ethical Approval:** Ethics committee review board license number: Ethical approval was also obtained from the ethics committee of Vice-Chancellor in Research Affairs - Shahid Beheshti University of Medical Sciences (IR.SBMU.MSP.REC.1403.124).

Funding/Support: This article is financially supported by Research Department of the School of Medicine Shahid Beheshti University of Medical Sciences (NO: 66019595).

Informed Consent: Informed written consent was obtained from all participants.

References

- Eisma MC. Prolonged grief disorder in ICD-11 and DSM-5-TR: Challenges and controversies. *Aust N Z J Psychiatry*. 2023;57(7):944-51. [PubMed ID: 36748103]. [PubMed Central ID: PMC10291380]. https://doi.org/10.1177/00048674231154206.
- Comtesse H, Smid GE, Rummel AM, Spreeuwenberg P, Lundorff M, Duckers MLA. Cross-national analysis of the prevalence of prolonged grief disorder. *J Affect Disord*. 2024;**350**:359-65. [PubMed ID: 38220101]. https://doi.org/10.1016/j.jad.2024.01.094.
- Szuhany KL, Malgaroli M, Miron CD, Simon NM. Prolonged Grief Disorder: Course, Diagnosis, Assessment, and Treatment. Focus (Am Psychiatr Publ). 2021;19(2):161-72. [PubMed ID: 34690579]. [PubMed Central ID: PMC8475918]. https://doi.org/10.1176/appi.focus.20200052.
- Silverman GK, Johnson JG, Prigerson HG. Preliminary explorations of the effects of prior trauma and loss on risk for psychiatric disorders in recently widowed people. *Isr J Psychiatry Relat Sci.* 2001;**38**(3-4):202-15. [PubMed ID: 11725418].
- Gilbert R, Widom CS, Browne K, Fergusson D, Webb E, Janson S. Burden and consequences of child maltreatment in high-income countries. *Lancet.* 2009;373(9657):68-81. [PubMed ID: 19056114]. https://doi.org/10.1016/S0140-6736(08)61706-7.
- Alvis L, Zhang N, Sandler IN, Kaplow JB. Developmental Manifestations of Grief in Children and Adolescents: Caregivers as Key Grief Facilitators. J Child Adolesc Trauma. 2023;16(2):447-57. [PubMed ID: 35106114]. [PubMed Central ID: PMC8794619]. https://doi.org/10.1007/s40653-021-00435-0.
- Ainsworth MD. The Bowlby-Ainsworth attachment theory. *Behav* Brain Sci. 2010;1(3):436-8. https://doi.org/10.1017/s0140525x00075828.
- Shaver PR, Tancredy CM. Emotion, attachment, and bereavement: A conceptual commentary. In: Stroebe MS, editor. *Handbook of bereavement research: Consequences, coping, and care.* Washington, DC, US: American Psychological Association; 2001. p. 63-88. https://doi.org/10.1037/10436-003.
- 9. Russ V, Stopa L, Sivyer K, Jane H, Tess M. The Relationship Between Adult Attachment and Complicated Grief: A Systematic Review. *Omega* (Westport). 2024;**89**(4):1293-319. [PubMed ID: 35635029]. [PubMed Central ID: PMC11423550]. https://doi.org/10.1177/00302228221083110.
- 10. Janshen A, Boelen PA, Schut HAW, Eisma MC. Do insecure attachment styles predict prolonged grief symptoms? Significant null findings.

Death Stud. 2024;**48**(10):1076-84. [PubMed ID: 38180083]. https://doi.org/10.1080/07481187.2023.2300063.

- Reime MA, O'Connor M, Hystad SW, Dyregrov K. Perceived social support and symptoms of prolonged grief after a drug-related death. *Death Stud.* 2024:1-10. [PubMed ID: 38970782]. https://doi.org/10.1080/07481187.2024.2376037.
- Maccallum F, Malgaroli M, Bonanno GA. Networks of loss: Relationships among symptoms of prolonged grief following spousal and parental loss. J Abnorm Psychol. 2017;126(5):652-62. [PubMed ID: 28594192]. [PubMed Central ID: PMC5523866]. https://doi.org/10.1037/abn0000287.
- 13. Treur A. Prolonged Grief Disorder and the Association with Social Contact: An Experience Sampling Study. Twente, Netherlands: University of Twente; 2024.
- Huda NU, Malik K, Ateeq A. Prolonged grief and social support among bereaved young adults. Int E-J Adv Soc Sci. 2024;10(28):74-84.
- Bernstein DP, Stein JA, Newcomb MD, Walker E, Pogge D, Ahluvalia T, et al. Development and validation of a brief screening version of the Childhood Trauma Questionnaire. *Child Abuse Neglect*. 2003;27(2):169-90. [PubMed ID: 12615092]. https://doi.org/10.1016/S0145-2134(02)00541-0.
- Ebrahimi H, Dejkam M, Seghatoleslam T. [Childhood Traumas and Suicide Attempt in adulthood]. *Iran J Psychiatry Clin Psychol.* 2014;19(4):275-282. FA.
- Collins NL, Read SJ. Adult attachment, working models, and relationship quality in dating couples. J Pers Soc Psychol. 1990;58(4):644. [PubMed ID: 14570079]. https://doi.org/10.1037/0022-3514.58.4.644.
- Hajializadeh K, Ahadi H, Jomehri F, Rahgozar M. Psychosocial predictors of barriers to cervical cancer screening among Iranian women: the role of attachment style and social demographic factors. *J Prev Med Hyg.* 2013;54(4):218-22. [PubMed ID: 24779284]. [PubMed Central ID: PMC4718322].
- Prigerson HG, Boelen PA, Xu J, Smith KV, Maciejewski PK. Validation of the new DSM-5-TR criteria for prolonged grief disorder and the PG-13-Revised (PG-13-R) scale. *World Psychiatry*. 2021;**20**(1):96-106. [PubMed ID: 33432758]. [PubMed Central ID: PMC7801836]. https://doi.org/10.1002/wps.20823.
- Ashouri A, Yousefi S, Prigerson HG. Psychometric properties of the PG-13-R scale to assess prolonged grief disorder among bereaved Iranian adults. *Palliat Support Care*. 2024;22(1):174-181. [PubMed ID: 37052287]. https://doi.org/10.1017/S1478951523000202.
- 21. Sarper E, Rodrigues DL. The role of perceived social support in the grief experiences of more anxious and self-compassionate people. *OMEGA-J Death Dying*. 2024:302228241229484. [PubMed ID: 38265046]. https://doi.org/10.1177/00302228241229484.
- 22. Smith KV, Wild J, Ehlers A. The Masking of Mourning: Social Disconnection After Bereavement and Its Role in Psychological Distress. *Clin Psychol Sci.* 2020;8(3):464-76. [PubMed ID: 32550046]. https://doi.org/10.1177/2167702620902748.
- 23. Kirkbride JB, Anglin DM, Colman I, Dykxhoorn J, Jones PB, Patalay P, et al. The social determinants of mental health and disorder: evidence,

prevention and recommendations. *World Psychiatry*. 2024;**23**(1):58-90. [PubMed ID: 38214615]. [PubMed Central ID: PMC10786006]. https://doi.org/10.1002/wps.21160.

- Chen MA, Brown RL, Chen JY, de Dios MA, Green CE, Heijnen CJ, et al. Childhood maltreatment, subjective social status, and health disparities in bereavement. *Psychoneuroendocrinology*. 2022;**135**. [PubMed ID: 34837775]. [PubMed Central ID: PMC8702370]. https://doi.org/10.1016/j.psyneuen.2021.105595.
- Dvilansky AS, Zadok H, Shoshani A, Samra NN, Verbeke W, Vrticka P, et al. The long-term associations of childhood parental loss with attachment, creativity, and epigenetic regulation. *Sc Rep.* 2025;15(1). [PubMed ID: 39924594]. [PubMed Central ID: PMC11808068]. https://doi.org/10.1038/s41598-025-89467-2.
- Calhoun CD, Stone KJ, Cobb AR, Patterson MW, Danielson CK, Bendezú JJ. The Role of Social Support in Coping with Psychological Trauma: An Integrated Biopsychosocial Model for Posttraumatic Stress Recovery. Psychiatric Quarterly. 2022;93(4):949-70. [PubMed ID: 36199000]. [PubMed Central ID: PMC9534006]. https://doi.org/10.1007/s11126-022-10003-w.
- 27. Krieger CLLV. Relationship Between Social Support and Childhood Trauma on Resilience. Minneapolis, USA: Walden University; 2021.
- Dvir Y, Ford JD, Hill M, Frazier JA. Childhood Maltreatment, Emotional Dysregulation, and Psychiatric Comorbidities. *Harvard Rev Psychiatry*. 2014;22(3):149-61. [PubMed ID: 24704784]. [PubMed Central ID: PMC4091823]. https://doi.org/10.1097/hrp.000000000000014.
- Boelen PA, Olff M, Smid GE. Traumatic loss: Mental health consequences and implications for treatment and prevention. *Europ J Psychotraumatol*. 2019;10(1). [PubMed ID: 33178405]. [PubMed Central ID: PMC7593703]. https://doi.org/10.1080/20008198.2019.1591331.
- Luckett T, Lenzo V, Sardella A, Musetti A, Faraone C, Quattropani MC. Toward a preventive approach to prolonged grief disorder in palliative care: Insecure attachment moderates the impact of perceived support on the severity of symptoms. *Plos One*. 2022;**17**(12). [PubMed ID: 36454987]. [PubMed Central ID: PMC9714949]. https://doi.org/10.1371/journal.pone.0266289.
- Shear K, Monk T, Houck P, Melhem N, Frank E, Reynolds C, et al. An attachment-based model of complicated grief including the role of avoidance. *Europ Arch Psychiatr Clin Neuroscience*. 2007;**257**(8):453-61. [PubMed ID: 17629727]. [PubMed Central ID: PMC2806638]. https://doi.org/10.1007/s00406-007-0745-z.
- Huh HJ, Kim KH, Lee H, Chae J. Attachment styles, grief responses, and the moderating role of coping strategies in parents bereaved by the Sewol ferry accident. *Europ J Psychotraumatol.* 2018;8(sup6). [PubMed ID: 29372009]. [PubMed Central ID: PMC5774413]. https://doi.org/10.1080/20008198.2018.1424446.
- Baker AW, Keshaviah A, Horenstein A, Goetter EM, Mauro C, Reynolds CF, et al. The Role of Avoidance in Complicated Grief: A Detailed Examination of the Grief-Related Avoidance Questionnaire (GRAQ) in a Large Sample of Individuals with Complicated Grief. *J Loss Trauma*. 2016;21(6):533-47. [PubMed ID: 28649184]. [PubMed Central ID: PMC5482544]. https://doi.org/10.1080/15325024.2016.1157412.