



# Cardiac Manifestations of Multisystem Inflammatory Syndrome in Children (MIS-C) Associated with SARS-CoV-2 Infection

Alireza Ghodsi <sup>1</sup>, Elnaz Mahmoudabadi <sup>1</sup>, Sara Ghahremani <sup>2</sup> and Abdolreza Malek <sup>2,\*</sup>

<sup>1</sup>Student Research Committee, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

<sup>2</sup>Department of Pediatrics, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

\*Corresponding author: Department of Pediatrics, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran. Email: malekar@mums.ac.ir

Received 2020 October 04; Revised 2020 December 24; Accepted 2020 December 28.

## Abstract

**Context:** Multisystem inflammatory syndrome in children (MIS-C) is an emerging condition after the spread of the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) pandemic, termed COVID-19. This study aimed to describe the cardiac manifestations of children diagnosed with MIS-C.

**Evidence Acquisition:** This narrative review was conducted by searching the PubMed, Scopus, and Google Scholar databases to review MIS-C cardiac manifestations up to September 30, 2020. The demographic features, past medical history, clinical signs and symptoms, cardiac involvement, and the type of COVID-19 diagnosis confirmation were extracted.

**Results:** In many children, MIS-C seems to be a post-infectious complication of the COVID-19 infection. This syndrome affects multiple organs and has various clinical manifestations mimicking Kawasaki disease. Patients frequently present with persistent fever, kidney injury, gastrointestinal (GI) problems, neurologic symptoms, mucosal changes, conjunctivitis, and cardiac involvement. Children with MIS are more likely to present with hypotension, shock, and cardiac dysfunction, rather than coronary artery abnormalities and arrhythmia. Children with MIS need close observation; some need to be hospitalized, and a few may need a Pediatric Intensive Care Unit (PICU) admission. Treatment currently includes anticoagulants, IV immunoglobulin, and anti-inflammatory drugs.

**Conclusions:** As a novel syndrome associated with SARS-CoV-2 infection, MIS-C is potentially lethal. Cardiac manifestations, including coronary and myocardial involvement, are common and should be carefully identified. With prompt diagnosis and proper treatment, most children will survive, but the outcomes of the disease are unknown, so long-term follow-ups are required.

**Keywords:** Multisystem Inflammatory Syndrome in Children, COVID-19, SARS-CoV-2, Cardiac Involvement

## 1. Context

The SARS-CoV-2 infection, termed COVID-19, has spread very quickly and affected all ages, even newborns (1, 2). Initially, there was a misconception among the researchers believing that the virus does not affect children. However, it was then proposed that children are usually asymptomatic or present mild symptoms, but a complete immunity cannot be proposed for this age group (3). Since late April 2020, multiple studies from Europe and the United States showed that 2% - 6% of SARS-CoV-2-infected children exhibited a severe multi-system inflammatory syndrome with similarities to Kawasaki disease (4-6).

As there is no comprehensive testing, the true incidence of this severe condition in pediatrics has remained unclear. This significant hyper-inflammatory response can cause cardiovascular disorders, and some of these children

may deteriorate faster and need admission to the Pediatric Intensive Care Unit (PICU) due to cardiogenic shock or acute left ventricular dysfunction (7). Apart from cardiac manifestations, various clinical symptoms have also been reported, including neurological, renal, significant gastrointestinal (GI), and mild respiratory symptoms, as well as rashes and stomatitis (8). Although it is still not fully understood whether this multisystem inflammatory syndrome in children (MIS-C) is a primary complication of COVID-19 infection or a post-infectious complication, a correlation is highly suggestive based on epidemiologic data (7).

There is little information about the cardiac involvement associated with COVID-19 in pediatric cases, and most of the studies on cardiac involvement are case reports or case series. It is also possible that children with an underlying cardiac disease are at a higher risk of experiencing

severe cardiac complications following the COVID-19 infection (9). Reports on MIS-C have shown coronary artery involvement, myocarditis, ventricular dysfunction, hemodynamic instability, and PICU admission, all of which suggest that cardiac dysfunction might be a notable risk factor for severe SARS-CoV-2 infection in pediatrics (10). Therefore, a literature review in this regard would be beneficial. This study aimed to review and summarize the available evidence on the potential cardiac clinical presentations in children with MIS to give a better perspective on management and care for these patients.

It should be noted that different terms have been used to refer to this novel condition, such as multisystem inflammatory syndrome in children (MIS-C), hyperinflammatory shock in children with COVID-19, "Coronavirus disease-19", "COVID19", "SARS-CoV-2", "systemic inflammatory response syndrome", "multisystem inflammatory disorder in children and adolescents", "MISC", "PMIS", "PCAI", "Coronasacki", and "Kawashocky". In this review article, we continue to use the term MIS-C.

## 2. Methods

A literature review was conducted by searching PubMed, Scopus, and Google Scholar. The following keywords were used for the search: "Pediatrics", "children", "heart", "cardiac", "Coronavirus disease-19", "COVID19", "SARS-CoV-2", "systemic inflammatory response syndrome", "multisystem inflammatory disorder in children and adolescents", "MISC", "PMIS", "PCAI", "Coronasacki", and "Kawashocky". We included case reports and case series related to MIS-C associated with SARS-CoV-2 infection. The inclusion criterion was defined as being a case report or case series reporting some findings of cardiac involvement in pediatrics with COVID-19. Articles published in languages other than English and those irrelevant to our objectives, like commentaries, other reviews, and letters to editors, were excluded. The last search was performed on September 30, 2020. We used the Newcastle-Ottawa scale (NOS) (11) for the quality assessment of the studies, and those with top quality were enrolled. The baseline data including the first author's name of the article, number of cases, age, sex, past medical history, ethnicity, clinical signs and symptoms, and the type of COVID-19 diagnosis confirmation (PCR or serology) were extracted. Moreover, the laboratory, electrocardiography, and echocardiography findings were gathered.

## 3. Definition of MIS-C

The three definitions of MIS-C, by the World Health Organization (WHO) (12), the Centers for disease control and prevention (CDC) (13), and the Royal College of Pediatrics and Child Health (RCPCH) (14), are presented in Table 1. The presence of fever, multisystem organ involvement without alternative plausible diagnoses, laboratory evidence of inflammation, and recent exposure to a COVID-19 case or evidence of COVID-19 infection are the key elements in all MIS-C cases. But, some signs, including fever duration and organ involvement, vary among these criteria.

## 4. Clinical Manifestation

The available information regarding the syndrome shows that the age of patients ranged from two months to 20 years, and the majority of cases were previously healthy (15, 16). Almost all affected children had a persistent fever for  $\geq 4$  days and GI symptoms, including abdominal pain, diarrhea, and vomit. Other common clinical manifestations were mucocutaneous changes resembling Kawasaki disease (skin rash and conjunctivitis), extremity edema, lymphadenopathy, headache, mild respiratory distress, myalgia, fatigue, and cardiac symptoms (Table 2). Some patients presented with shock and hypotension requiring PICU admission (17, 18). The cardiac findings in MIS-C patients are divergent from Kawasaki disease's manifestations. Children with this syndrome were more likely to present with hypotension, shock, and cardiac dysfunction, rather than coronary artery abnormalities (Figure 1) (19).

## 5. COVID-19 Infection

Epidemiological information indicates that SARS-CoV-2 is the possible cause of the syndrome, but the causality is unknown (1). Based on the studies mentioned in Table 2, the positivity percentage of the COVID-19 reverse-transcriptase protein chain reaction (RT-PCR) test varies from 0% to 100%. In most reports, it was positive in less than 50% of cases. On the other hand, the majority of the studies had evidence of positive immunoglobulin G (IgG) antibodies. These data suggest that a post-infectious disease is more likely to be responsible for this condition than an active infection (44).

## 6. Cardiac Involvement

### 6.1. Cardiac Dysfunction

In most cases diagnosed with MIS-C, left ventricular systolic dysfunction has been reported (Table 2). In the first

**Table 1.** Case Definition of Multisystem Inflammatory Syndrome in Children


	<b>WHO</b>	<b>CDC</b>	<b>NHS-the Royal College of Pediatrics and Child Health</b>
<b>Age (years)</b>	0 - 19	< 21	Children
<b>Fever</b>	Fever $\geq$ 3 days	Subjective persistent fever $\geq$ 24 hours or documented fever $>$ 38.0°C for $\geq$ 24 hours	Persistent fever $>$ 38.5°C
<b>Clinical findings</b>	Mucocutaneous inflammation signs (e.g., stomatitis), bilateral non-purulent conjunctivitis or rash, shock or hypotension, ventricular dysfunction, valvulitis, pericarditis, or coronary involvement (including an increased level of NT-proBNP /troponin or echo findings), acute GI manifestations, including abdominal pain, diarrhea, or vomiting	Evidence of clinical deterioration requiring hospital admission, in addition to multiple ( $\geq$ 2) organ dysfunction (renal, dermatologic, cardiovascular, respiratory, GI, hematologic, or neurologic)	Single or multi-organ dysfunction (cardiovascular, respiratory, renal, GI, hematologic, dermatologic, or neurologic). Oxygen requirement and hypotension have been reported in most kids. Other features, including abdominal pain, diarrhea, vomiting, conjunctivitis, rash, sore throat, cough, respiratory symptoms, headache, confusion, lymphadenopathy, stomatitis, neck swelling, hands and feet edema, syncope have been seen in some children.
<b>Laboratory findings</b>	Elevated inflammatory factors (e.g., CRP, ESR, or procalcitonin) evidence of coagulopathy in laboratory data (PT, PTT, INR, and D-dimer)	Elevated ESR, CRP, LDH, procalcitonin, fibrinogen, D-dimer, ferritin, IL-6, hypoalbuminemia neutrophilia; lymphocytopenia	Abnormal fibrinogen level, hypoalbuminemia, high D-dimer and ferritin amount Some: acute kidney injury, anemia, coagulopathy, thrombocytopenia, elevated IL-6, elevated IL-10, hypertransaminasemia, proteinuria, high troponin, increased creatine kinase level, elevated triglycerides, high LDH
<b>Evidence of COVID-19 infection</b>	Positive for SARS-CoV-2 infection by serology, antigen test, or RT-PCR or exposure to patients with COVID-19 infection	RT-PCR, or antigen test, or serology positive for COVID-19 or possible contact with COVID-19 patients within a month before the initiation of clinical features	SARS-CoV-2 RT-PCR testing may be positive or negative
<b>Exclusion of other microbial causes</b>	Exclusion of any other infectious causes of inflammation, including toxic shock syndrome, bacterial sepsis, staphylococcal or streptococcal infections	Exclusion of alternative plausible diagnoses	Exclusion of any other microbial cause, including infectious myocarditis, bacterial sepsis, and staphylococcal/streptococcal toxic shock syndromes
<b>Additional comments</b>	MIS-C must be considered in children with characteristics of toxic shock syndrome or typical or atypical Kawasaki disease	Consider MIS-C in any pediatric death with evidence of COVID-19 infection. It should be considered in children with features of typical or atypical Kawasaki disease who meet the case definition for MIS-C	Children may fulfill full or partial criteria for Kawasaki disease

Abbreviations: APTT, activated partial thromboplastin time; COVID-19, coronavirus disease 2019; CRP, C-reactive protein, echo, echocardiography; CXR, chest X-ray; ESR, erythrocyte sedimentation rate; GI, gastrointestinal; IL, interleukin; LDH, lactic acid dehydrogenase; MIS-C, multisystem inflammatory syndrome in children; NT-proBNP, N-terminal pro-B-type natriuretic peptide; PT, prothrombin time; PTT, partial thromboplastin time; 2, RT-PCR, reverse transcriptase-polymerase chain reaction; SARS-CoV-2, severe acute respiratory syndrome coronavirus.

report of MIS-C, Riphagen et al. (24) found ventricular dysfunction in six out of eight cases. The majority of available studies in the literature also reported that ventricular dysfunction happens in up to 50% of MIS-C patients (Table 2). Two case series mentioned cardiac dysfunction as an inclusion criterion (20, 38). Belhadjer et al. (20) considered an acute LV failure (LV ejection fraction (LVEF)  $<$  50%), and Grimaud et al. considered cardiogenic/vasogenic shock as an inclusion criterion (38). The incidence rate of cardiogenic shock is much greater in this syndrome than in the Kawasaki disease (60% vs. 7%) (45). Matsubara et al. (46)

found severe LV dysfunction in MIS-C patients. They suggested that the subclinical myocarditis might lead to ventricular dysfunction.

The pathophysiology of cardiac dysfunction in patients with this syndrome is not known. The possible causes of acute myocardial dysfunction following a COVID-19 infection in adults include hypoxic injury, myocarditis, ischemic heart injury due to coronary artery disease, stress cardiomyopathy (Takotsubo), acute cor pulmonale, and systemic inflammatory response syndrome (SIRS) (47-50).



**Figure 1.** Schematic representation of clinical signs of MIS-C patients (both CDC and WHO criteria are shown)

## 6.2. Laboratory Findings

The elevated troponin and B-type natriuretic peptide (BNP)/pro-BNP levels have been reported in many patients with MIS-C. In most studies, the elevated BNP and troponin levels have been assessed as biomarkers to diagnose myocardial injury in the absence of cardiac magnetic resonance imaging (CMRI) or myocardial biopsies (15).

## 6.3. Coronary Involvement

Most reports have mentioned coronary involvement in 0-40% of cases (Table 2) (16, 21, 24, 26, 41). Mild coronary artery dilation with z-scores of 2 - 2.5 has been reported in most cases. However, large coronary artery aneurysm cases have also been mentioned. Some studies found the late development of coronary artery dilation, which necessitates an ongoing follow-up of MIS-C patients.

## 6.4. ECG Findings

Some case series reported rhythm abnormalities with variable severity in 4% - 58% of patients (Table 2) (18, 22). The most frequently reported arrhythmic manifestations were QTc prolongation, ST-segment changes, and premature atrial or ventricular beats, which all are non-specific. Whittaker et al. (18) reported first- and second-degree atrioventricular blocks, and two other studies mentioned atrial fibrillation (18, 33). Hemodynamic collapse and the need for extracorporeal membrane oxygenation (ECMO)

support due to sustained dysrhythmias have also been reported in these patients (18, 24).

## 6.5. Management

This syndrome is a newly reported condition, and only a few studies have addressed it so far. Our knowledge about it is, therefore, limited, and the treatment of children with MIS has been based on the experts' advice and the management of Kawasaki disease. Due to the similarity of the symptoms, the treatment method for adults with COVID-19 and other systemic inflammatory diseases can also be used in pediatrics. Management of these children requires a multidisciplinary care team comprising pediatric specialists in cardiology, infectious disease, critical care, and rheumatology.

Generally, the management is decided based on symptoms and their severity. Due to the potential shortage of drug supplies in a pandemic and considering side effects, pharmacotherapy is not recommended for non-hospitalized children. For children who present with mild symptoms, supportive care, including respiratory support and fluid resuscitation, is, therefore, recommended. However, children with hemodynamic instability and severe illness require PICU admission, mostly for inotropic support, which was reported in 20-100% of the cases (15-18, 20-22, 24-27, 30-35, 37, 38). Some of the PICU-admitted children required a veno-arterial (V-A) support (0% - 28%) (15, 17, 18, 20, 24, 30, 33).

Various treatments have been suggested, but their effectiveness is still questionable. Furthermore, these treatments are based on experts' opinions with no evidence to affirm them.

#### 6.6. Cardiac Support

As mentioned above, a large proportion of children presenting with hemodynamic instability required acute resuscitation. Therefore, it is necessary to follow the pediatric resuscitation guidelines (51). Children suspicious of ventricular dysfunction and cardiogenic shock should receive smaller fluid bounces (e.g., 10 mg/kg), with an evaluation of the signs of fluid overload before each administration.

#### 6.7. Immunomodulatory Therapy

The advantages of using immunomodulatory therapy in the treatment of Kawasaki disease, as well as other systemic inflammatory disorders, are well established (52, 53). An anti-inflammatory therapy, including intravenous immunoglobulins (IVIGs) and corticosteroids, was used in most patients, and a few cases also received an anti-inflammatory dosage of aspirin (15, 16, 18, 20-22, 24, 26, 31, 32, 34-38). It is critical to remember that the administration of IVIG in patients with cardiac dysfunction must be slower to reduce the risk of fluid overload. The dosage of corticosteroids is based on clinical judgment, but in more severe patients, using a low dosage is recommended.

The assessment of the pattern of cytokine storm in patients with MIS-C showed that an important component of this disorder is macrophage activation, as it is also observed in Kawasaki disease and other autoimmune disorders such as systemic lupus erythematosus (35, 54). Therefore, corticosteroids are another option in the treatment of MIS-C patients, as they can modulate this condition. However, corticosteroids may cause hypertension that can further exacerbate the underlying cardiac problem (45).

Two treatment protocols with corticosteroids have been proposed. The first method involves an intravenous injection of 0.8 mg/kg methylprednisolone, twice a day for 5 - 7 days or until achieving a normal CRP level and then continuing with oral treatment with 2 mg/kg/day for 2 - 3 weeks. The second protocol includes intravenous methylprednisolone 10 - 30 mg/kg/day for three days, followed by oral prednisone/prednisolone 2 mg/kg/day for four days or until achieving a normal CRP level and then tapering the treatment over 2 - 3 weeks. It is important to know that corticosteroids should not be administered in an active infection phase (1).

In some studies, cytokine blockers have been used as a supplemental therapy, for example, interleukin 1 receptor antagonist (e.g., anakinra), interleukin 6 (IL-6) inhibitors (e.g., tocilizumab), and tumor necrosis factor (TNF)- $\alpha$  inhibitors (e.g., infliximab) (16, 18, 20, 22-25, 27, 29-31, 34). These drugs can be prescribed for children who do not respond to routine treatments.

#### 6.8. Antiplatelet Treatment and Anticoagulation

Hypercoagulable state, blood stasis due to immobilization, possible endothelial injury, and ventricular dysfunction are the proposed reasons for the increased risk of thrombotic complications. As a result, anticoagulant therapy should be considered based on coagulation tests and symptoms (55, 56).

#### 6.9. Antiviral Therapy

The benefits of antiviral therapy, such as remdesivir, for children with this syndrome are still unknown (57, 58). The reports suggest that MIS-C is more likely a post-infectious complication in children rather than an active infection. Nonetheless, antiviral drugs could be considered in patients with a positive RT-PCR test, after consulting an infectious disease specialist.

### 7. Conclusions

In conclusion, children seem to proceed better with the novel coronavirus infection than adults. On the other hand, some children show signs and symptoms of MIS-C, which is a severe complication of the disease. Pediatricians should be aware of this syndrome and differentiate it from other differential diagnoses, including the Kawasaki disease. These children can quickly deteriorate and should closely be observed. The etiology of MIS-C is not yet fully understood, and treatment is mostly based on experts' opinions. More studies are, therefore, required to define evidence-based management for this new syndrome, and our study played a part in this literature contribution.

### Footnotes

**Authors' Contribution:** All authors contributed to the study design, data collection, writing, and preparing this article.

**Conflict of Interests:** The authors declare no conflict of interest.

**Funding/Support:** The authors received no specific funding for this work.

## References

- Nakra NA, Blumberg DA, Herrera-Guerra A, Lakshminrusimha S. Multi-System Inflammatory Syndrome in Children (MIS-C) Following SARS-CoV-2 Infection: Review of Clinical Presentation, Hypothetical Pathogenesis, and Proposed Management. *Children (Basel)*. 2020;7(7). doi: [10.3390/children7070069](https://doi.org/10.3390/children7070069). [PubMed: 32630212]. [PubMed Central: PMC7401880].
- Farhat AS, Sayedi SJ, Akhlaghi F, Hamed A, Ghodsi A. Coronavirus (COVID-19) Infection in Newborns. *Int J Pediatr*. 2020;8(6):ii1513-7.
- Ludvigsson JF. Systematic review of COVID-19 in children shows milder cases and a better prognosis than adults. *Acta Paediatr*. 2020;109(6):1088-95. doi: [10.1111/apa.15270](https://doi.org/10.1111/apa.15270). [PubMed: 32202343]. [PubMed Central: PMC7228328].
- Dong Y, Mo X, Hu Y, Qi X, Jiang F, Jiang Z, et al. Epidemiology of COVID-19 Among Children in China. *Pediatrics*. 2020;145(6). doi: [10.1542/peds.2020-0702](https://doi.org/10.1542/peds.2020-0702). [PubMed: 32179660].
- Liguoro I, Pilotto C, Bonanni M, Ferrari ME, Pusiol A, Nocerino A, et al. SARS-CoV-2 infection in children and newborns: a systematic review. *Eur J Pediatr*. 2020;179(7):1029-46. doi: [10.1007/s00431-020-03684-7](https://doi.org/10.1007/s00431-020-03684-7). [PubMed: 32424745]. [PubMed Central: PMC7234446].
- Sanna G, Serrau G, Bassareo PP, Neroni P, Fanos V, Marcialis MA. Children's heart and COVID-19: Up-to-date evidence in the form of a systematic review. *Eur J Pediatr*. 2020;179(7):1079-87. doi: [10.1007/s00431-020-03699-0](https://doi.org/10.1007/s00431-020-03699-0). [PubMed: 32474800]. [PubMed Central: PMC7261213].
- Hennon TR, Penque MD, Abdul-Aziz R, Alibrahim OS, McGreevy MB, Prout AJ, et al. COVID-19 associated Multisystem Inflammatory Syndrome in Children (MIS-C) guidelines; a Western New York approach. *Prog Pediatr Cardiol*. 2020;101232. doi: [10.1016/j.ppedcard.2020.101232](https://doi.org/10.1016/j.ppedcard.2020.101232). [PubMed: 32837142]. [PubMed Central: PMC7244417].
- Ghodsi A, Azarfar A, Ghahremani S. A Review of Coronavirus Disease (COVID-19) in Children. *J Pediatr Nephrol*. 2020;8(3).
- Castagnoli R, Votto M, Licari A, Brambilla I, Bruno R, Perlini S, et al. Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) Infection in Children and Adolescents: A Systematic Review. *JAMA Pediatr*. 2020;174(9):882-9. doi: [10.1001/jamapediatrics.2020.1467](https://doi.org/10.1001/jamapediatrics.2020.1467). [PubMed: 32320004].
- Ghodsi A, Malek A, Ghahremani S. A Review of Multisystem Inflammatory Syndrome in Children (MIS-C) Associated with COVID-19. *Hormozgan Med J*. 2020;24(4). doi: [10.5812/hmj.107048](https://doi.org/10.5812/hmj.107048).
- Wells GA, Shea B, O'Connell DA, Peterson J, Welch V, Losos M, et al. *The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandomised studies in meta-analyses*. Oxford; 2000.
- Labe P, Ly A, Sin C, Nasser M, Chapelon-Fromont E, Ben Said P, et al. Erythema multiforme and Kawasaki disease associated with COVID-19 infection in children. *J Eur Acad Dermatol Venereol*. 2020;34(10):e539-41. doi: [10.1111/jdv.16666](https://doi.org/10.1111/jdv.16666). [PubMed: 32455505]. [PubMed Central: PMC7283825].
- Rivera-Figueroa EI, Santos R, Simpson S, Garg P. Incomplete Kawasaki Disease in a Child with Covid-19. *Indian Pediatr*. 2020;57(7):680-1. [PubMed: 32393680]. [PubMed Central: PMC7387257].
- Raymond TT, Das A, Manzuri S, Ehrett S, Guleserian K, Brenes J. Pediatric COVID-19 and Pericarditis Presenting With Acute Pericardial Tamponade. *World J Pediatr Congenit Heart Surg*. 2020;11(6):802-4. doi: [10.1177/2150135120949455](https://doi.org/10.1177/2150135120949455). [PubMed: 32909890]. [PubMed Central: PMC7484599].
- Dufort EM, Koumans EH, Chow Ej, Rosenthal EM, Muse A, Rowlands J, et al. Multisystem Inflammatory Syndrome in Children in New York State. *N Engl J Med*. 2020;383(4):347-58. doi: [10.1056/NEJMoa2021756](https://doi.org/10.1056/NEJMoa2021756). [PubMed: 32598830]. [PubMed Central: PMC7346766].
- Wolfson A, Mannarino S, Giacomet V, Camporesi A, Zuccotti G. Acute myocardial injury: a novel clinical pattern in children with COVID-
- Lancet Child Adolesc Health*. 2020;4(8):e26-7. doi: [10.1016/S2352-4642\(20\)30168-1](https://doi.org/10.1016/S2352-4642(20)30168-1). [PubMed: 32497521]. [PubMed Central: PMC7263818].
- Feldstein LR, Rose EB, Horwitz SM, Collins JP, Newhams MM, Son MBF, et al. Multisystem Inflammatory Syndrome in U.S. Children and Adolescents. *N Engl J Med*. 2020;383(4):334-46. doi: [10.1056/NEJMoa2021680](https://doi.org/10.1056/NEJMoa2021680). [PubMed: 32598831]. [PubMed Central: PMC7346765].
- Whittaker E, Bamford A, Kenny J, Kafourou M, Jones CE, Shah P, et al. Clinical Characteristics of 58 Children With a Pediatric Inflammatory Multisystem Syndrome Temporally Associated With SARS-CoV-2. *JAMA*. 2020;324(3):259-69. doi: [10.1001/jama.2020.10369](https://doi.org/10.1001/jama.2020.10369). [PubMed: 32511692]. [PubMed Central: PMC7281356].
- Shulman ST. Pediatric COVID-associated Multi-system Inflammatory Syndrome (PMIS). *J Pediatr Infect Dis Soc*. 2020.
- Belhadjer Z, Meot M, Bajolle F, Khraiche D, Legembre A, Abakka S, et al. Acute Heart Failure in Multisystem Inflammatory Syndrome in Children in the Context of Global SARS-CoV-2 Pandemic. *Circulation*. 2020;142(5):429-36. doi: [10.1161/CIRCULATIONAHA.120.048360](https://doi.org/10.1161/CIRCULATIONAHA.120.048360). [PubMed: 32418446].
- Toubiana J, Poirault C, Corsia A, Bajolle F, Fourgeaud J, Angoulvant F, et al. Kawasaki-like multisystem inflammatory syndrome in children during the covid-19 pandemic in Paris, France: prospective observational study. *BMJ*. 2020;369:m2094. doi: [10.1136/bmj.m2094](https://doi.org/10.1136/bmj.m2094). [PubMed: 32493739]. [PubMed Central: PMC7500538].
- Cheung EW, Zachariah P, Gorelik M, Boneparth A, Kernie SG, Orange JS, et al. Multisystem Inflammatory Syndrome Related to COVID-19 in Previously Healthy Children and Adolescents in New York City. *JAMA*. 2020;324(3):294-6. doi: [10.1001/jama.2020.10374](https://doi.org/10.1001/jama.2020.10374). [PubMed: 32511676]. [PubMed Central: PMC7281352].
- Dolinger MT, Person H, Smith R, Jarchin L, Pittman N, Dubinsky MC, et al. Pediatric Crohn Disease and Multisystem Inflammatory Syndrome in Children (MIS-C) and COVID-19 Treated With Infliximab. *J Pediatr Gastroenterol Nutr*. 2020;71(2):153-5. doi: [10.1097/MPG.0000000000002809](https://doi.org/10.1097/MPG.0000000000002809). [PubMed: 32452979]. [PubMed Central: PMC7268863].
- Riphagen S, Gomez X, Gonzalez-Martinez C, Wilkinson N, Theocharis P. Hyperinflammatory shock in children during COVID-19 pandemic. *Lancet*. 2020;395(10237):1607-8. doi: [10.1016/S0140-6736\(20\)31094-1](https://doi.org/10.1016/S0140-6736(20)31094-1). [PubMed: 32386565]. [PubMed Central: PMC7204765].
- Waltuch T, Gill P, Zinns LE, Whitney R, Tokarski J, Tsung JW, et al. Features of COVID-19 post-infectious cytokine release syndrome in children presenting to the emergency department. *Am J Emerg Med*. 2020;38(10):2246 e3-6. doi: [10.1016/j.ajem.2020.05.058](https://doi.org/10.1016/j.ajem.2020.05.058). [PubMed: 32471782]. [PubMed Central: PMC7255141].
- Liacciardi F, Pruccoli G, Denina M, Parodi E, Taglietto M, Rosati S, et al. SARS-CoV-2-Induced Kawasaki-Like Hyperinflammatory Syndrome: A Novel COVID Phenotype in Children. *Pediatrics*. 2020;146(2). doi: [10.1542/peds.2020-1711](https://doi.org/10.1542/peds.2020-1711). [PubMed: 32439816].
- Borocco C, Pouletty M, Galeotti C, Meinzer U, Faye A, Kone-Paut I, et al. Response to 'Correspondence on 'Paediatric multisystem inflammatory syndrome temporally associated with SARS-CoV-2 mimicking Kawasaki disease (Kawa-COVID19): a multicentre cohort' by Mastrolia et al. *Ann Rheum Dis*. 2020. doi: [10.1136/annrheumdis-2020-218814](https://doi.org/10.1136/annrheumdis-2020-218814). [PubMed: 33023959].
- Jones VG, Mills M, Suarez D, Hogan CA, Yeh D, Segal JB, et al. COVID-19 and Kawasaki Disease: Novel Virus and Novel Case. *Hosp Pediatr*. 2020;10(6):537-40. doi: [10.1542/hpeds.2020-0123](https://doi.org/10.1542/hpeds.2020-0123). [PubMed: 32265235].
- Balasubramanian S, Nagendran TM, Ramachandran B, Ramanan AV. Hyper-inflammatory Syndrome in a Child With COVID-19 Treated Successfully With Intravenous Immunoglobulin and Tocilizumab. *Indian Pediatr*. 2020;57(7):681-3. [PubMed: 32393681]. [PubMed Central: PMC7387261].
- Kaushik S, Aydin SI, Derespina KR, Bansal PB, Kowalsky S, Trachtman R, et al. Multisystem Inflammatory Syndrome in Children Assos-

- ciated with Severe Acute Respiratory Syndrome Coronavirus 2 Infection (MIS-C): A Multi-institutional Study from New York City. *J Pediatr*. 2020;224:24–9. doi: 10.1016/j.jpeds.2020.06.045. [PubMed: 32553861]. [PubMed Central: PMC7293760].
31. Greene AG, Saleh M, Roseman E, Sinert R. Toxic shock-like syndrome and COVID-19: Multisystem inflammatory syndrome in children (MIS-C). *Am J Emerg Med*. 2020;38(11):2492 e5–6. doi: 10.1016/j.ajem.2020.05.117. [PubMed: 32532619]. [PubMed Central: PMC7274960].
  32. Rauf A, Vijayan A, John ST, Krishnan R, Latheef A. Multisystem Inflammatory Syndrome with Features of Atypical Kawasaki Disease during COVID-19 Pandemic. *Indian J Pediatr*. 2020;87(9):745–7. doi: 10.1007/s12098-020-03357-1. [PubMed: 32462354].
  33. Deza Leon MP, Redzepi A, McGrath E, Abdel-Haq N, Shawaqfeh A, Sethuraman U, et al. COVID-19-Associated Pediatric Multisystem Inflammatory Syndrome. *J Pediatric Infect Dis Soc*. 2020;9(3):407–8. doi: 10.1093/jpids/piaa061. [PubMed: 32441749]. [PubMed Central: PMC7313914].
  34. Chiotos K, Bassiri H, Behrens EM, Blatz AM, Chang J, Diorio C, et al. Multisystem Inflammatory Syndrome in Children During the Coronavirus 2019 Pandemic: A Case Series. *J Pediatric Infect Dis Soc*. 2020;9(3):393–8. doi: 10.1093/jpids/piaa069. [PubMed: 32463092]. [PubMed Central: PMC7313950].
  35. Verdoni L, Mazza A, Gervasoni A, Martelli L, Ruggeri M, Ciuffreda M, et al. An outbreak of severe Kawasaki-like disease at the Italian epicentre of the SARS-CoV-2 epidemic: an observational cohort study. *Lancet*. 2020;395(10239):1771–8. doi: 10.1016/S0140-6736(20)31103-X. [PubMed: 32410760]. [PubMed Central: PMC7220177].
  36. Blondiaux E, Parisot P, Redheuil A, Tzaroukian L, Levy Y, Sileo C, et al. Cardiac MRI in Children with Multisystem Inflammatory Syndrome Associated with COVID-19. *Radiology*. 2020;297(3):E283–8. doi: 10.1148/radiol.2020202288. [PubMed: 32515676]. [PubMed Central: PMC7294821].
  37. Ramcharan T, Nolan O, Lai CY, Prabhu N, Krishnamurthy R, Richter AG, et al. Paediatric Inflammatory Multisystem Syndrome: Temporally Associated with SARS-CoV-2 (PIMS-TS): Cardiac Features, Management and Short-Term Outcomes at a UK Tertiary Paediatric Hospital. *Pediatr Cardiol*. 2020;41(7):1391–401. doi: 10.1007/s00246-020-02391-2. [PubMed: 32529358]. [PubMed Central: PMC7289638].
  38. Grimaud M, Starck J, Levy M, Marais C, Chareyre J, Khraiche D, et al. Acute myocarditis and multisystem inflammatory emerging disease following SARS-CoV-2 infection in critically ill children. *Ann Intensive Care*. 2020;10(1):69. doi: 10.1186/s13613-020-00690-8. [PubMed: 32488505]. [PubMed Central: PMC7266128].
  39. Jain MK, Sahu SK, Behera JR, Patnaik S. Multisystem Inflammatory Syndrome in Children Associated with COVID-19 Treated with Oral Steroid. *Indian J Pediatr*. 2021;88(1):106. doi: 10.1007/s12098-020-03497-4. [PubMed: 32930973]. [PubMed Central: PMC7490472].
  40. Heidemann SM, Tilford B, Bauerfeld C, Martin A, Garcia RU, Yagiela L, et al. Three Cases of Pediatric Multisystem Inflammatory Syndrome Associated with COVID-19 Due to SARS-CoV-2. *Am J Case Rep*. 2020;21:e925779. doi: 10.12659/AJCR.925779. [PubMed: 32790652]. [PubMed Central: PMC7447294].
  41. Alnashri H, Aljohani N, Tayeb S, Rabie N, AlBenayan E, Alharthi A, et al. A challenging case of multisystem inflammatory syndrome in children related to coronavirus Disease-19 hospitalized under adult medical service. *IDCases*. 2020;22: e00957. doi: 10.1016/j.idcr.2020.e00957. [PubMed: 32953455]. [PubMed Central: PMC7489252].
  42. Torres JP, Izquierdo G, Acuna M, Pavez D, Reyes F, Fritis A, et al. Multisystem inflammatory syndrome in children (MIS-C): Report of the clinical and epidemiological characteristics of cases in Santiago de Chile during the SARS-CoV-2 pandemic. *Int J Infect Dis*. 2020;100:75–81. doi: 10.1016/j.ijid.2020.08.062. [PubMed: 32861823]. [PubMed Central: PMC7452906].
  43. Dolhnikoff M, Ferreira Ferranti J, de Almeida Monteiro RA, Duarte-Neto AN, Soares Gomes-Gouveia M, Viu Degasparre N, et al. SARS-CoV-2 in cardiac tissue of a child with COVID-19-related multisystem inflammatory syndrome. *Lancet Child Adolesc Health*. 2020;4(10):790–4. doi: 10.1016/S2352-4642(20)30257-1. [PubMed: 32828177]. [PubMed Central: PMC7440866].
  44. McCrindle BW, Manlhiot C. SARS-CoV-2-Related Inflammatory Multisystem Syndrome in Children: Different or Shared Etiology and Pathophysiology as Kawasaki Disease? *JAMA*. 2020;324(3):246–8. doi: 10.1001/jama.2020.10370. [PubMed: 32511667].
  45. Rodriguez-Gonzalez M, Castellano-Martinez A, Cascales-Poyatos HM, Perez-Reviriego AA. Cardiovascular impact of COVID-19 with a focus on children: A systematic review. *World J Clin Cases*. 2020;8(21):5250–83. doi: 10.12998/wjcc.v8.i21.5250. [PubMed: 33269260]. [PubMed Central: PMC7674714].
  46. Matsubara D, Kauffman HI, Wang Y, Calderon-Anyosa R, Nadaraj S, Elias MD, et al. Echocardiographic Findings in Pediatric Multisystem Inflammatory Syndrome Associated With COVID-19 in the United States. *J Am Coll Cardiol*. 2020;76(17):1947–61. doi: 10.1016/j.jacc.2020.08.056. [PubMed: 32890666]. [PubMed Central: PMC7467656].
  47. Shi S, Qin M, Shen B, Cai Y, Liu T, Yang F, et al. Association of Cardiac Injury With Mortality in Hospitalized Patients With COVID-19 in Wuhan, China. *JAMA Cardiol*. 2020;5(7):802–10. doi: 10.1001/jamacardio.2020.0950. [PubMed: 32211816]. [PubMed Central: PMC7097841].
  48. Creel-Bulos C, Hockstein M, Amin N, Melhem S, Truong A, Sharifpour M. Acute Cor Pulmonale in Critically Ill Patients with Covid-19. *N Engl J Med*. 2020;382(21):e70. doi: 10.1056/NEJM2010459. [PubMed: 32374956]. [PubMed Central: PMC7281714].
  49. Zheng YY, Ma YT, Zhang JY, Xie X. COVID-19 and the cardiovascular system. *Nat Rev Cardiol*. 2020;17(5):259–60. doi: 10.1038/s41569-020-0360-5. [PubMed: 32139904]. [PubMed Central: PMC7095524].
  50. Tersalvi G, Vicenzi M, Calabretta D, Biasco I, Pedrazzini G, Winterton D. Elevated Troponin in Patients With Coronavirus Disease 2019: Possible Mechanisms. *J Card Fail*. 2020;26(6):470–5. doi: 10.1016/j.cardfail.2020.04.009. [PubMed: 32315733]. [PubMed Central: PMC7166030].
  51. Edelson DP, Sasson C, Chan PS, Atkins DL, Aziz K, Becker LB, et al. Interim Guidance for Basic and Advanced Life Support in Adults, Children, and Neonates With Suspected or Confirmed COVID-19: From the Emergency Cardiovascular Care Committee and Get With The Guidelines-Resuscitation Adult and Pediatric Task Forces of the American Heart Association. *Circulation*. 2020;141(25):e933–43. doi: 10.1161/CIRCULATIONAHA.120.047463. [PubMed: 32270695]. [PubMed Central: PMC7302067].
  52. Pineton de Chambrun M, Luyt CE, Beloncle F, Gousseff M, Mauhin W, Argaud L, et al. The Clinical Picture of Severe Systemic Capillary-Leak Syndrome Episodes Requiring ICU Admission. *Crit Care Med*. 2017;45(7):1216–23. doi: 10.1097/CCM.0000000000002496. [PubMed: 28622216].
  53. Thompson PD, Myerburg RJ, Levine BD, Udelson JE, Kovacs RJ. Eligibility and disqualification recommendations for competitive athletes with cardiovascular abnormalities: task force 8: coronary artery disease: a scientific statement from the American Heart Association and American College of Cardiology. *Circulation*. 2015;132(22):e310–e4. doi: 10.1161/CIR.000000000000244. [PubMed: 26401111]. [PubMed Central: PMC4500000].
  54. Ravelli A, Minoia F, Davi S, Horne A, Bovis F, Pistorio A, et al. 2016 Classification Criteria for Macrophage Activation Syndrome Complicating Systemic Juvenile Idiopathic Arthritis: A European League Against Rheumatism/American College of Rheumatology/Paediatric Rheumatology International Trials Organisation Collaborative Initia-

- tive. *Ann Rheum Dis*. 2016;75(3):481-9. doi: [10.1136/annrheumdis-2015-208982](https://doi.org/10.1136/annrheumdis-2015-208982). [PubMed: [26865703](https://pubmed.ncbi.nlm.nih.gov/26865703/)].
55. Wright FL, Vogler TO, Moore EE, Moore HB, Wohlauer MV, Urban S, et al. Fibrinolysis Shutdown Correlation with Thromboembolic Events in Severe COVID-19 Infection. *J Am Coll Surg*. 2020;231(2):193-203. doi: [10.1016/j.jamcollsurg.2020.05.007](https://doi.org/10.1016/j.jamcollsurg.2020.05.007). [PubMed: [32422349](https://pubmed.ncbi.nlm.nih.gov/32422349/)]. [PubMed Central: [PMC7227511](https://pubmed.ncbi.nlm.nih.gov/PMC7227511/)].
56. Raval JS, Burnett AE, Rollins-Raval MA, Griggs JR, Rosenbaum L, Nielsen ND, et al. Viscoelastic testing in COVID-19: a possible screening tool for severe disease? *Transfusion*. 2020;60(6):1131-2. doi: [10.1111/trf.15847](https://doi.org/10.1111/trf.15847). [PubMed: [32374920](https://pubmed.ncbi.nlm.nih.gov/32374920/)]. [PubMed Central: [PMC7267656](https://pubmed.ncbi.nlm.nih.gov/PMC7267656/)].
57. Beigel JH, Tomashek KM, Dodd LE, Mehta AK, Zingman BS, Kalil AC, et al. Remdesivir for the Treatment of Covid-19 - Final Report. *N Engl J Med*. 2020;383(19):1813-26. doi: [10.1056/NEJMoa2007764](https://doi.org/10.1056/NEJMoa2007764). [PubMed: [32445440](https://pubmed.ncbi.nlm.nih.gov/32445440/)]. [PubMed Central: [PMC7262788](https://pubmed.ncbi.nlm.nih.gov/PMC7262788/)].
58. Goldman JD, Lye DCB, Hui DS, Marks KM, Bruno R, Montejano R, et al. Remdesivir for 5 or 10 Days in Patients with Severe Covid-19. *N Engl J Med*. 2020;383(19):1827-37. doi: [10.1056/NEJMoa2015301](https://doi.org/10.1056/NEJMoa2015301). [PubMed: [32459919](https://pubmed.ncbi.nlm.nih.gov/32459919/)]. [PubMed Central: [PMC7377062](https://pubmed.ncbi.nlm.nih.gov/PMC7377062/)].

**Table 2.** Demographics, Clinical Manifestations, and Cardiac Involvement in Reports of Children WITH Multisystem Inflammatory Syndrome

Author	Number of Cases	Age, y	Sex	Past Medical History	Rhythm	Symptoms	Cardiac involvement		ProBNP/bNP	RT-PCR	SARS-CoV-2 test Serology		
							Coronary involvement	Ventricular Function					
Belladher et al. (20)	35	1-16	18 M, 17 F	Previously healthy (3); asthma (3); STE (1); overweight (BMI > 25) (6)	-	Fever (35); GI symptoms (29); asthenia (3); respiratory distress (23); rash (20); lymphadenopathy (21); meningitis (1); chest pain (6); cardiac shock (28); pericardial effusion (3)	Mild coronary dilation ( $\geq$ score > 2) (6/35); no aneurysms	LVEF < 30% (16); LVEF 30 to 50% (25); LV < 50% (inclusion criteria); takotsubo (1); segmental wall hypokinesis (3); global LV hypokinesis (31)	Elevated (mild to moderate)	Elevated proBNP or BNP (35)	14 cases were positive	30/35; IgG+ (28); IgM+ (2)	
Toubiana et al. (21)	21	4-17	9 M, 12 F	-	African ancestry (12)	Fever (21); GI symptoms (21); rash (16); conjunctivitis (5); hypotension/shock (7); lips and oral cavity changes (6); changes to extremities (10); serous effusion (12); irritability (12); cervical lymphadenopathy (12); neurological features (6)	Dilated coronary arteries ( $\geq$ score = 2.5) (5); echo-bright coronary arteries (3)	Ventricular arrhythmia (1); ST elevation at onset (1)	Elevated (7)	Elevated (14/18)	8 cases were positive	IgG+ (19)	
Cheung et al. (22)	17	2-16	11 M, 4 F	Mild asthma (3)	Ashkenazi; Jewish (3); Black (4); White (6); Asian (1)	Fever (17); skin rash (12); GI symptoms (5); conjunctivitis (1); mucosal changes (9); neurological problems (6); respiratory (7); shock (3); myalgia (5); cervical lymphadenopathy (6); skin desquamation (3)	Echobright coronaries (7); medium-sized aneurysm ( $\geq$ score = 2.5) (1)	Normal-mild LV dysfunction (1); moderate-severe LV dysfunction (6)	Elevated (14)	Elevated (15)	8 cases were positive	IgM/IgG+ (9)	
Whittaker et al. (23)	58	6-14	25 M, 33 F	Comorbid (7); asthma (3); idiopathic (1); sickle cell trait (1); epilepsy (1); neuro-disability (1)	White (12); Black (22); Asian (18); other (6)	Fever; GI symptoms; confusion (5); lymphadenopathy (9); skin rash (10); respiratory symptoms (12); conjunctivitis (26); peripheral edema (9); mucosal changes (7); sore throat (6); shock (29)	Dilated coronary arteries ( $\geq$ score > 2) (8/29); 2.5 (7) giant aneurysm (2)	IV dysfunction (18/29)	4/58; atrial fibrillation (1); first-degree AV block (1); second-degree AV block (1); intractable bradycardia (1)	Elevated	Elevated	15 cases were positive	IgG+ (40)/46)
Dollinger et al. (24)	1	14	M	Crohn disease	-	Persistent fever; GI symptoms; rash; myalgia	-	-	-	-	Positive	-	
Feldstein et al. (27)	186	3-12	115 M, 71 F	Previously healthy (95); 51/88 had an underlying disease other than obesity (autoimmune or immunocompromise (10); respiratory (3); cardiac (5); BMI-based obesity (45)	White (35); Black (46); Latino or Hispanic (57)	Persistent fever; GI symptoms (17); conjunctivitis (10); skin rash (10); lymphadenopathy (18); peripheral edema (69); respiratory symptoms (11); oropharyngeal changes (7); cardiovascular symptoms (49)	Coronary artery aneurysm ( $\geq$ score $\geq$ 2.5) (15/70)	Myocardial dysfunction (90)	Arrhythmia (12)	Elevated (50/128)	Elevated (73/73)	73 cases were positive	IgM/IgG+ (85)
Dufort et al. (25)	99	0-20	53 M, 46 F	Pre-existing condition (36/99); 29 of them had obesity	White (29); Black (31); Hispanic (31)	Persistent fever or chills; GI symptoms (79); rash (59); swollen hands or feet (9); hypotension (6); mucosal change (6); neurologic symptoms (30); shock (10); lower respiratory symptoms (40); upper respiratory (27); chest pain (11)	9 had coronary artery aneurysm ( $\geq$ score > 2.5)	Some degree of ventricular dysfunction (51); myocarditis (52)	-	Elevated (63/89)	Elevated (74/82)	50 cases were positive	IgG+ (76/77); IgM+ (37/77)
Birnboim et al. (24)	8	4-14	3 F, 5 M	Previously healthy (6); autism (1); atopica areata and allergic rhinitis (1)	Afro-Caribbean (6)	Persistent fever; GI symptoms (7); myalgia (1); rash (4); odynophagia (3); conjunctivitis (5); headache (2); hypotension (2)	Echobright coronary vessels; (8) giant aneurysm (1)	Mild to severe IV dysfunction; (8) giant aneurysm (1)	-	-	-	-	
Waluch et al. (26)	4	5-13	1 F, 3 M	Previously healthy (2); hypothyroidism (1); asthma (1)	-	Persistent fever; GI symptoms; rash (2); myalgia (1); cough (2); conjunctivitis (3); fatigue (2); hypotension	Slight ectasia (1); Dilated coronary arteries (1); mildly dilated coronary artery (1)	Moderately depressed IV function	Elevated (1)	Elevated	-	IgG+ (4)	

Licciardet et al. (26)	2	7,12	2 M	Previously healthy pRAPA syndrome (1)	-	Persistent fever; rash; GI symptoms; hypotension; shock; peripheral edema; conjunctivitis	No	Ventricular dysfunction (2)	Elevated (2)	Elevated (1)	Negative	IgM+; IgG+	
Borocco et al. (27)	16	5-12	8 M, 8 F	Previously healthy (10); asthma (2); overweight (4)	-	Persistent fever; GI symptoms (13); mucosal changes (4); rash (13); respiratory symptoms (2); anosmia (1); neurological symptoms (9); shock (1); conjunctivitis (5); lymphadenopathy (6); arthritis (7); pericarditis (4)	Coronary dilation (3) median z score 2.6	Myocarditis (7); LVEF 35%	-	Elevated (11)	II cases were positive	IgG+ (7/8)	
Jones et al. (28)	1	6 months	F	None	-	Persistent fever; mucosal changes; conjunctivitis; peripheral edema; rash; irritability; respiratory symptoms	No	Normal	-	-	Positive	-	
Balasubramanian et al. (29)	1	8	M	None	Indian	Persistent fever; rash; odynophagia; mild respiratory distress; mucosal changes; conjunctivitis; hypotension; peripheral edema	No	Normal	-	-	Positive	-	
Label et al. (12)	2	3,6	2 M	None	-	Rash; conjunctivitis; mucosal changes; persistent fever (1); cervical lymphadenopathy	-	-	-	-	One case was positive	-	
Woller et al. (16)	5	2-168 months	2 M, 3 F	None	-	Persistent fever; GI symptoms; hypotension; shock (5); conjunctivitis (1); rash (3); respiratory distress (1)	No	Mild moderate heart dysfunction (5); LVEF < 50% (3)	Elevated	Elevated	Positive	-	
Kanshiket et al. (30)	33	6-13	20 M, 13 F	Previously healthy (7); overweight (4); asthma (5)	Latino or Hispanic (15); Black (13)	Persistent fever; GI symptoms (23); conjunctivitis (2); rash (14); neurologic involvement (4); respiratory symptoms (1); hypotension (2); mucosal changes (7)	Prominent coronary arteries (6); coronary ectasia (2)	LVEF < 50% (21); LVEF < 30% (4)	Elevated	Elevated	II cases were positive	IgM/IgG+ (27)	
Greene et al. (31)	1	11	F	None	-	Persistent fever; rash; GI symptoms; shock; malaise; leg pain; some throat pain	No	Decreased LV function	-	Elevated	Elevated	Positive	
Rauf et al. (32)	1	5	M	None	-	Persistent fever; conjunctivitis; GI symptoms; hypotension; peripheral edema; purpura	No	Moderate LV dysfunction (Fl=35%); LV global hypokinesia	-	Elevated	Elevated	Negative	
Deza Leon et al. (33)	1	6	F	None	-	Fever; conjunctivitis; respiratory distress; rash; edema; dryophagia	No	Mildly diminished; LVEF at onset severe dysfunction; requiring ECMO	Elevated	-	Positive	-	
Chiotos et al. (34)	6	5-14	1 M, 5 F	None	African-American (2); Caucasian (2)	Persistent fever (6); rash (2); conjunctivitis (2); GI symptoms (5); shock (6); peripheral edema (2); mucosal changes (3); respiratory failure (4); irritability (2); headache (1)	Echobright coronaries (1); diffuse dilation right coronary artery (z score 3.5) (1)	Mild-Moderate LV dysfunction (4)	-	Elevated (3/5)	Elevated	3 cases were positive	
Verdoni et al. (35)	10	3-16	7 M, 3 F	None	Caucasian	Persistent fever; Peripheral edema; hypotension; conjunctivitis; rash; GI symptoms (6); meningeal signs (4); mucosal changes (4)	Coronary aneurysms (> 4 mm) (2)	LVEF < 50% (5)	Elevated (5)	Elevated	Two cases were positive	IgM+ (3); IgG+ (8)	
Blondiaux et al. (36)	4	6-12	3 F; 1 M	None	-	Persistent fever; conjunctivitis; rash; GI symptoms	No	Transient systolic dysfunction (4); LVEF < 30% (1)	ST depression (1); T waves abnormalities (1)	Elevated	Elevated	Negative	IgG+ (4); IgM+ (1)
Rambaran et al. (37)	15	7-11	11 M, 4 F	-	African/Afro-Caribbean (6); Asian (6); mixed (2)	Persistent fever (15); GI symptoms (13); lethargy (4); myalgia (4)	Coronary artery abnormalities (14); 1 aneurysm; ectasia (7) prominent	Reduced LV fractional shortening (8); LVEF < 55% (12)	Abnormal PR interval (9) abnormal I waves	Elevated (15)	Two cases were positive	IgM/IgG/IgA+ (22/12)	

Rivera-Figueroa et al. (3)	1	5	M	None	African-American	Persistent fever; rash; GI symptoms; shock; conjunctivitis; peripheral edema; mucosal changes	No	Normal	-	Elevated	-	Positive	
Grimaudet al. (38)	20	3-15	10 M, 10 F	-	-	Fever (20); abdominal pain (20); rash (10); conjunctivitis (6); mucosal changes (5); lymphadenopathy (2)	No	Cardiogenic/haemoplegic shock (inclusion criteria) (20)	-	Elevated	Elevated	10 cases were positive	IgG+ (15)
Raymond et al. (44)	1	7	F	None	-	Cough; chest pain; orthopnoea; tachycardia; pericarditis	-	-	Low voltage QRS with electrical alternans T-wave inversion in inferior and lateral leads Sinus tachycardia	Not elevated	Elevated	Positive	-
Jain et al. (39)	1	3	M	-	-	Fever; urticarial rashes; diarrhea; facial puffiness; conjunctivitis; edema of hands and feet	-	Normal	-	-	-	Positive	-
Heidenmann et al. (40)	3	5-7	2 M, 1 F	-	-	Fever; conjunctivitis; dry and cracked lips; rash; cervical lymphadenopathy; GI symptoms; Shock	Dilation of the proximal left arterial descending coronary artery (z-score: 2.1) and the main left coronary artery (z-score: 2.3) seen in one case. Others were normal.	Mildly decreased LV function mid/mtrial insufficiency (2)	Early repolarization in the lateral precordial leads low voltages in the limb leads Sinus tachycardia PR interval prolongation	Elevated	-	Positive (1)	IgG+ (3)
Anashiri et al. (41)	1	16	M	-	-	Fever; diarrhea; vomiting; generalized abdominal pain; pleuritic chest; fissuring of the lower lip; conjunctivitis; bilateral elbow and knee effusion; hypopigmented macular rash on the chest	-	Hypokinesia of inferior wall with an ejection fraction of 45%	Normal	Elevated	-	-	IgG+
Torres et al. (42)	27	0-14	14 M, 13 F	-	-	Fever (27); abdominal pain (13); diarrhea (7); vomiting (13); rash (14); conjunctivitis (13); oral/mucosal changes (13); Cough (7); peripheral extremity changes (7)	Coronary artery abnormalities were in one patient	Myocardial dysfunction 4/26	-	-	-	14	10
Dollnikoff et al. (43)	1	11	F	None	African	Cardiovascular shock; persistent fever;odynophagia; myalgia; abdominal pain	No	Diffuse LV hypokinesia with decreased LVEF (3%) no segmental wall motion abnormalities	Sinus tachycardia	Elevated	Positive	-	

Abbreviations: AV, atrio-ventricular; BV, biventricular; ECG, electrocardiogram; F, female; GI, gastrointestinal; IQF, interquartile range; LVEF, left ventricular ejection fraction; LV, left ventricle; M, male; RV, right ventricle; SD, standard deviation; SARS-CoV-2; severe acute respiratory syndrome coronavirus 2; VR, ventricular tachycardia.