








# Factors Associated with Intra Hospital Neonatal Mortality: A Case Study of the Neonatology Department at the Regional Hospital Center of Agadir (Morocco)

Hanane El Ghazouani <sup>1,\*</sup>, Laila Lahlou <sup>2</sup>, Souad Yakini <sup>1</sup>, Oussama El Amraoui<sup>3</sup>, Yassine Ziani<sup>3</sup>, El Arbi Bouaiti <sup>4</sup>, Amina Barkat <sup>1</sup>

<sup>1</sup> Department of Clinical Epidemiology and Medico-Surgical Sciences, Health and Nutrition Research Team of the Mother-Child Couple, Faculty of Medicine and Pharmacy, University Mohammed V, Rabat, Morocco

<sup>2</sup> Laboratory of Health and Science, Therapeutic Innovation, Translational Research and Epidemiology, Faculty of Medicine and Pharmacy, Ibn Zohr University, Agadir, Morocco

<sup>3</sup> Higher Institute of Nursing Professions and Health Techniques of Agadir, Agadir, Morocco

<sup>4</sup> Biostatistics Research Team (BRT), Clinical Epidemiology and Medico-Surgical Sciences, Faculty of Medicine and Pharmacy, Mohamed V University, Rabat, Morocco

\*Corresponding Author: Department of Clinical Epidemiology and Medico-Surgical Sciences, Health and Nutrition Research Team of the Mother-Child Couple, Faculty of Medicine and Pharmacy, University Mohammed V, Rabat, Morocco. Email: elghazouani.hanane@gmail.com

Received 2023 December 27; Revised 2024 July 24; Accepted 2024 August 13

## Abstract

**Background:** Neonatal mortality (NNM) has significantly decreased in developed countries. However, it remains a major public health issue and a particular challenge in developing countries, with Morocco being no exception, as its NNM rate remains high.

**Objectives:** This study aims to identify factors associated with NNM within the Neonatology Department at the Regional Hospital Center of Agadir.

**Methods:** An analytical cross-sectional study design was chosen. To determine factors associated with early NNM, we employed binary logistic regression in both univariate and multivariate analyses, including all statistically significant variables in the multivariate model.

**Results:** Three hundred and eight deceased neonates were included. The intra-hospital NNM rate was 17.4%, 95% CI (13.2 - 21.7), during the period from 2018 to 2021. A predominance of early NNM (81.8%). 52.3% were females. 47.7% of the deceased had low birth weight. 65.9% of the cases had an Apgar score < 7. 44.2% were resuscitated. The factors that were statistically associated with the risk of early NNM were as follows: Admission age  $\leq 1$  day (aOR: 17.87; 95% CI: 5.78 - 55.29), neonatal admission age 2 - 7 days (aOR: 6.64; 95% CI: 1.68 - 26.30). Intra-hospital referral (aOR: 4.49; 95% CI: 1.59 - 12.65); emergency consultations (aOR: 4.25; 95% CI: 1.28 - 14.01).

**Conclusions:** Neonatal mortality remains a major public health challenge in Morocco. Different risk factors associated with NNM have been identified. Addressing these factors could reduce neonatal deaths in low-resource countries. For this reason, we need to intensify our efforts to develop reliable and precise approaches.

**Keywords:** Neonatal Mortality, Morocco, Factors Associated, Intra Hospital

## 1. Background

The arrival of a newborn child is an expected moment, and in the great majority of cases, it is presented as a source of happiness and hope for the future, both for individual families and for the community as a whole (1). However, the time of delivery

and the first month are critical for child survival, as the risk of morbidity and mortality can increase considerably if complications emerge (2, 3). The first 28 days of life—the neonatal period—represent the most vulnerable time for a child's survival. Children face the highest risk of dying during their first month of life, with a global rate of 17 deaths per 1,000 live births. Globally, approximately 2.4 million newborns died

during the first month of life in 2020, which is about 6,500 per day (4). In 2020, the United States experienced a neonatal mortality (NNM) rate of 3.4 per 1,000 live births. Europe and Central Asia recorded a mortality rate of 4.1 per 1,000 live births, Tunisia registered a rate of 11.8 per 1,000 live births, and Libya had a rate of six per 1,000 live births. However, in Morocco, NNM has decreased from 19.3 deaths per 1,000 live births for newborns in 2010 to 11.5 deaths per 1,000 live births in 2020 (5). Children who die within the first 28 days of life exhibit conditions and illnesses linked to the lack of quality care at birth or the absence of care or treatment provided by skilled personnel immediately after birth and in the early days of life (6). Furthermore, approximately one-third of all neonatal deaths occur on the first day after birth, and nearly three-quarters occur during the first week of life (4). In 2018, Northern Africa had the highest NNM rate at 28 deaths per 1,000 live births (7). Moreover, the NNM rate in Morocco was 13.6% in 2018 (8) and 11.1% in 2021 (9). For in-hospital NNM, a rate of 19% was reported (10), and early NNM was estimated at 21.4% (11). Prematurity, low birth weight, neonatal infections, and neonatal distress are the main factors of morbidity and the leading causes of death among newborns, with neonatal asphyxia being the primary cause (12). Regarding in-hospital NNM, neonatal infections are the leading cause of death, followed by neonatal respiratory distress and perinatal asphyxia. Prematurity emerges as the primary predisposing factor (13). The burden of NNM and morbidity remains high, and the opportunities to enhance women's well-being and provide attentive care to newborns have not been fully exploited (14). In summary, each newborn is a roadmap that outlines the path to ending preventable NNM and reducing disabilities (15).

## 2. Objectives

The aim of this study was to identify factors associated with NNM in the Neonatology Unit at the Regional Center of Agadir and to estimate its prevalence.

## 3. Methods

This was an analytical cross-sectional study focused on deceased newborns recorded at the Neonatology and Intensive Care Unit of Hassan II Regional Hospital in Agadir. The study was conducted over the period from January 1, 2018, to July 31, 2021. It is a level II reference neonatal intensive care center in the Souss-Massa region. It receives and provides care for newborns who have health problems after birth and who are referred from the regional maternity ward, provincial hospitals,

birthing centers, and clinics in the city of Agadir and neighboring regions. The facility has a capacity of 18 beds, along with 10 warming tables and five self-priming syringes.

The study population consists of newborns who were hospitalized and deceased at the neonatology unit of the Agadir Regional Hospital. The sample size was estimated using the sample size estimation formula for prevalence studies (16, 17). The required sample size calculation was based on the actual prevalence of in-hospital NNM (19%) (10) and the national prevalence of NNM of 11.1 per 1,000 live births (9), using a significance level of 5% (i.e.,  $\alpha = 0.05$ ) with a study power of 80%. Subsequently, a minimum of 108 deceased newborns were required to obtain statistically representative data. The selection of records to be included was done randomly from the death registers, applying the simple random sampling method. All newborns who died in the Neonatology Unit at the Agadir Regional Hospital during the study period were included. The study excluded any newborns who were discharged alive from the Neonatology Unit and children who were hospitalized and died after 28 days of age. To avoid bias, the measuring instrument was tested on a small sample of files excluded following random sampling.

The criteria for judgment were NNM, which is defined as death occurring within the first 28 complete days of life (9, 15, 18). It was categorized as early deaths during the first week and late deaths during the subsequent three weeks (19). The modes of use of hospital services and benefits were: (1) ordinary admission—any scheduled hospitalization outside of emergencies; (2) emergency admissions (emergency consultations, transfers, and intra-hospital referrals) (20).

The tool used for data collection was an exploitation form of medical records and hospitalization registers, which was designed and selected based on a literature review. This exploitation form was developed using Sphinx software and included open-ended and multiple-choice questions aimed at exploring several aspects: (a) neonatal (age, gender, weight, hospitalization diagnosis, etc.); (b) maternal (mother's age, place of origin, etc.); and (c) obstetric (place and method of delivery, etc.).

The collected data were analyzed using Jamovi V21 software. Firstly, a descriptive analysis was performed, and the qualitative variables were expressed as frequency and percentages; the quantitative variables were expressed as medians and interquartile ranges. The distribution of quantitative variables was assessed based on graphical representations, including histograms, Q-Q plots, and the Shapiro-Wilk test. The

assessment of associations was conducted using the chi2 test or Fisher's exact test, according to the conditions of application for each test. To determine factors associated with early NNM, binary logistic regression was employed in both univariate and multivariate analyses; all variables with  $P \leq 0.05$  in the univariate analysis were included in the multivariate model. The odds ratios (OR) were calculated with a 95% confidence interval (95% CI). The significance level was set at  $P \leq 0.05$ .

The Ethics Committee of the Faculty of Medicine and Pharmacy of Rabat approved this study with reference number (C64-20). The study strictly adhered to ethical considerations, including (a) obtaining authorization for access to records from the responsible administration; (b) preserving anonymity; (c) confidentiality; and (d) professional secrecy.

## 4. Results

### 4.1. Neonatal Mortality Rate

The NNM rate at the Hassan II Regional Hospital in Agadir was 17.44%, 95% CI (13.2-21.7), for the period from 2018 to 2021. Among the newborns who died in the Neonatology Unit at CHR Hassan II in Agadir during the study duration, there were 1,535 cases, while the hospitalized newborns totaled 8,802.

### 4.2. Demographic and Clinical Characteristics

A total of 308 deceased neonates were included in the study; early NNM accounted for 81.81%. The median length of hospitalization was 1 [1-1] day, with 95.45% being hospitalized during their first week of life, of which 97% died during the early neonatal period. 52.3% of the deceased were from rural areas. There was a predominance of females, with a percentage of 52.3% compared to 47.7% for males. As for the median weight, it was 2200 (1450 - 3200) grams (g), with 47.7% of cases having low birth weight, and 50% of them lost their lives during the early neonatal period. 43.7% of the deaths during the early neonatal period were from intra-hospital referrals, 31.7% were transfers, and 17.2% were admitted through the emergency department.

The median Apgar score was 7 [6-8]. 65.9% of cases had an Apgar score less than 7 at the 5th minute after birth, of which nearly 64% were classified as early NNM. The main diagnoses for hospitalization of newborns who died in the neonatal unit were prematurity (59.7%), neonatal distress (14.3%), convulsions (5.8%), malformation (4.3%), jaundice (2.9%), and neonatal infection (1%). (Table 1). The types of malformations were

distributed as follows: 8 (32%) neural tube defects, 4 (16%) multiple malformations, 2 (8%) cardiac malformations, and abdominal, ocular, and genital malformations, each recorded at a frequency of 1 (4%). However, 8 (32%) of malformation types were not specified. 85.7% were born to women aged between 20 and 35 years. 72.4% of cases were born in a hospital maternity ward, followed by 25.6% born in a private clinic. Only 1.3% and 0.6% were born at home and in a birthing center, respectively. 65.3% of cases were from singleton pregnancies, 30.8% from twin pregnancies, and only 3.9% from multiple pregnancies (Table 1).

The results from Table 1 indicated a statistically significant difference in the frequency of neonatal admission age ( $P < 0.001$ ), mode of admission ( $P < 0.001$ ), and prematurity ( $P = 0.052$ ) between early NNM and late NNM.

### 4.3. Factors Associated with Early Neonatal Mortality

In univariate analysis, the factors associated with early NNM were APGAR (cOR: 0; 95% CI: 6.62 - 57.96); admission age  $\leq 1$  day (cOR: 19.6; 95% CI: 6.62 - 57.96); neonatal admission age 2 - 7 days (cOR: 4.94; 95% CI: 1.34 - 18.40); emergency consultations (cOR: 3.64; 95% CI: 1.25 - 10.59); intra-hospital referral (cOR: 5.93; 95% CI: 2.26 - 15.56). However, the other factors studied in the univariate analysis (Table 2) were not significantly associated with early NNM ( $P > 0.05$ ).

After adjusting for the following factors: Neonatal age, mode of admission, jaundice, and length of hospitalization, the factors that were statistically associated with the risk of early NNM were as follows: Newborns admitted during the ultra-early neonatal period (age  $\leq 1$  day) had a risk multiplied by 17 of death (aOR: 17.87; 95% CI: 5.78 - 55.29). In addition, those admitted during the early neonatal period (2 - 7 days) had a risk 6 times higher of death (aOR: 6.64; 95% CI: 1.68 - 26.30). Intra-hospital referral increases the risk of death by 4 times (aOR: 4.49; 95% CI: 1.59 - 12.65). Newborns admitted via the emergency department had a 4-fold increased risk of death (aOR: 4.25; 95% CI: 1.28 - 14.01) (Table 2).

## 5. Discussion

Through the reported results, the present study attempted to identify the factors associated with NNM and to estimate its prevalence. Newborns are the most vulnerable population group in the world due to their specific needs; consequently, they serve as a sensitive indicator of the quality of services provided, as they can die within a few days (21). The NNM rate at Hassan II

**Table 1.** Distribution of Deceased Newborns According to Demographic and Clinical Characteristics and Comparison between Early and Late Neonatal mortality

Variables	Total <sup>a</sup> ; N = 308	Early NNM; n = 252	Late NNM; n = 56	P-Value
<b>Sex</b>				0.501 <sup>b</sup>
Female	161 (52.3)	134 (53.2)	27 (48.2)	
Male	147 (47.7)	118 (46.8)	29 (51.8)	
<b>Apgar 7 [6-8] <sup>c</sup></b>				0.113 <sup>b</sup>
<7	203 (65.9)	161 (63.9)	42 (75)	
≥7	105 (34.1)	91 (36.1)	14 (25)	
<b>Birth weight 2200 [1450 - 3200] <sup>c</sup></b>				0.198 <sup>b</sup>
Normal	123 (40)	95 (37.2)	28 (50)	
Low birth weight	147 (47.7)	126 (50)	21 (37.5)	
Overweight	38 (12.3)	31 (12.3)	7 (12.5)	
<b>Neonatal age at admission [1-1] <sup>c</sup></b>				< 0.001 <sup>d</sup>
≤1 jr	268 (87.01)	231 (91.7)	33 (58.9)	
2-7	26 (8.44)	16 (6.3)	9 (16.1)	
>7	14 (4.55)	5 (2)	14 (25)	
<b>Mode of admission</b>				< 0.001 <sup>b</sup>
Ordinary	28 (9.1)	17 (6.7)	11 (19.6)	
Emergency consultations	53 (17.2)	45 (17.9)	8 (14.3)	
Intra-hospital referral	122 (39.6)	110 (43.7)	12 (21.4)	
Transfer	105 (34.1)	80 (31.7)	25 (44.6)	
<b>Origin</b>				0.206 <sup>b</sup>
Rural	161 (52.3)	136 (54)	25 (44.6)	
Urban	147 (47.7)	116 (46)	31 (55.4)	
<b>Type of pregnancy</b>				0.536 <sup>b</sup>
Singleton	201 (65.3)	161 (63.9)	40 (71.4)	
Twin	95 (30.8)	81 (32.1)	14 (25)	
Multiple	12 (3.9)	10 (4)	2 (3.6)	
<b>Place of delivery</b>				0.717 <sup>d</sup>
Private clinic	79 (25.6)	62 (24.6)	17 (30.4)	
Hospital maternity ward	223 (72.4)	184 (73)	39 (69.6)	
Birth center	2 (0.6)	2 (0.8)	0	
At home	4 (1.3)	4 (1.6)	0	
<b>Maternal age</b>				0.555 <sup>d</sup>
≤ 20	17 (5.5)	13 (5.2)	4 (7.1)	
> 20 - 35	264 (85.7)	215 (85.3)	49 (87.5)	
> 35	27 (8.8)	24 (9.5)	3 (5.4)	
<b>Prematurity</b>				0.05 <sup>b</sup>
Yes	184 (59.7)	157 (62.3)	27 (48.2)	
No	124 (40.3)	95 (37.7)	29 (51.8)	
<b>Neonatal infection</b>				0.086 <sup>d</sup>
Yes	3 (1)	1 (0.4)	2 (3.6)	
No	305 (99)	251 (99.6)	54 (96.4)	
<b>Neonatal distress</b>				0.398 <sup>b</sup>
Yes	44 (14.3)	38 (15.1)	6 (10.7)	
No	264 (85.7)	214 (84.9)	50 (89.3)	
<b>Seizure</b>				0.752 <sup>d</sup>
Yes	18 (5.8)	14 (5.6)	4 (7.1)	
No	290 (94.2)	238 (94.4)	52 (92.9)	
<b>Malformation</b>				0.759 <sup>d</sup>
Yes	19 (6.2)	15 (6)	4 (7.1)	
No	289 (93.8)	237 (94)	52 (92.9)	

Abbreviations: NNM, neonatal mortality; ds, days.

<sup>a</sup> Expressed in numbers (percentage).

<sup>b</sup> Chi2 test.

<sup>c</sup> Expressed in median and interquartile ranges.

<sup>d</sup> Fisher's exact test.

Regional Hospital in Agadir is 17.44%, 17.44% (95% CI: 13.2 - 21.7) for the period from 2018 to 2021. This rate remains significant compared to that recorded at the reference hospital in Kindu, where the hospital NNM rate was 15% (22). However, it is lower than the NNM rates reported by studies conducted in Meknes and in the South Kivu province of Congo, 19% and 26.6%, respectively (10, 23). The main diagnoses for hospitalization of newborns who died in the neonatal unit were prematurity (59.7%), neonatal distress (14.3%), convulsions (5.8%), malformation (4.3%), jaundice (2.9%), and neonatal

infection (1%). WHO declared that premature birth, birth asphyxia, infections, and birth defects are the main causes of neonatal death (6).

Birth weight is a good indicator to enhance the chances of a newborn's survival. Low birth weight and overweight are considered risk markers, with low birth weight predominating in our sample (47.7%), and 50% of them lost their lives during the early neonatal period. In this regard, Ilunga et al. state that there is a relationship between birth weight and mortality. The lower the birth weight, the higher the mortality rate (24). The findings

**Table 2.** Factors Associated with Early Neonatal Mortality

Variables	Analyze Univariate			Analyze Multivariate		
	cOR	IC 95%	P-Value	aOR	IC 95%	P-Value
<b>Sex</b>						
Female	1.22	0.68 - 2.18	0.502	-	-	-
Male	1	1	-	-	-	-
<b>Apgar</b>						
<7	0.590	0.30 - 1.14	0.115	-	-	-
≥7	1	1	-	-	-	-
<b>Birth weight</b>						
Low birth weight	1.77	0.94 - 3.30	0.074	-	-	-
Overweight	1.31	0.51 - 3.28	0.571	-	-	-
Normal	1	1	-	-	-	-
<b>Neonatal age at admission</b>						
≤ 1jrs	19.6	6.62 - 57.96	< 0.001	17.87	5.78 - 55.29	< 0.001
2 - 7 jrs	4.978	1.34 - 18.40	0.016	6.64	1.68 - 26.30	0.007
> 7jrs	1	1	-	-	-	-
<b>Origin</b>						
Rural	1.45	0.81 - 2.60	0.208	-	-	-
Urban	1	1	-	-	-	-
<b>Mode of admission</b>						
Emergency consultations	3.64	1.25 - 10.59	0.018	4.25	1.28 - 14.01	0.017
Intra hospital referral	5.93	2.26 - 15.56	< 0.001	4.49	1.59 - 12.65	0.004
Transfer	2.07	0.20 - 1.16	0.106	-	-	-
Ordinary	1	1	-	-	-	-
<b>Type of pregnancy</b>						
Twuin	1.44	0.74 - 2.79	0.285	-	-	-
Multiple	1.24	0.26 - 5.89	0.785	-	-	-
Singleton	1	1	-	-	-	-
<b>Place of delivery</b>						
Private clinic	0.773	0.40 - 1.46	0.429	-	-	-
Birth center	3.32e+6	0.00 - inf	0.993	-	-	-
At home	3.32e+6	0.00 - inf	0.990	-	-	-
Hospital maternity ward	1	1	-	-	-	-
<b>Maternal age</b>						
≤ 20	0.741	0.23 - 2.37	0.613	-	-	-
> 35	1.823	0.52 - 6.30	0.342	-	-	-
20 - 35	1	1	-	-	-	-
<b>Prematurity</b>	1.78	0.99 - 3.18	0.054	-	-	-
<b>Neonatal infection</b>	0.108	0.009 - 1.21	0.071	-	-	-
<b>Neonatal distress</b>	1.48	0.59 - 3.69	0.401	-	-	-
<b>Seizure</b>	0.765	0.24 - 2.42	0.648	-	-	-
<b>Malformation</b>	0.823	0.26 - 2.58	0.738	-	-	-

Abbreviations: cOR, crude odds ratio; aOR, adjusted odds ratio; CI, confidence interval.

of Nouria et al. revealed that 84.27% of deceased newborns had low birth weight (25). In light of these results, new studies on all parameters associated with low birth weight are necessary to monitor the ongoing evolution of this issue and its recurrent associated factors (26). The Apgar score is the first test that a newborn undergoes at birth. It provides an idea of the newborn's adaptation to extrauterine life, its circulatory functions, as well as its neurological condition. In

France, a study on NNM reported that newborns with a very low Apgar score had an earlier age of death (27). This is consistent with the data from our series. The median Apgar score was 7 [6-8], with 65.9% of cases having an Apgar score less than 7 at the 5th minute of birth, of which nearly 64% were classified as early NNM.

The results of our study identified four factors significantly associated with the risk of early NNM: Admission age ≤ 1 day (aOR: 17.87; 95% CI: 5.78 - 55.29);



neonatal admission age 2 - 7 days (aOR: 6.64; 95% CI: 1.68-26.30); emergency consultations (aOR: 4.25; 95% CI: 1.28 - 14.01); intra-hospital referral (aOR: 4.49; 95% CI: 1.59 - 12.65).

The age of hospitalization provides insight into the alarming age of newborns as well as the delicate stage after birth and reflects the early onset of a serious medical condition. In our series, 95.45% were hospitalized during the early neonatal phase, with 98% of them dying during the early neonatal period. Additionally, the multivariate analysis is statistically significant, showing an association between the age of hospitalization and early NNM for admission age  $\leq$  1 day (aOR: 17.87; 95% CI: 5.78 - 55.29), and neonatal admission age 2 - 7 days (aOR: 6.64; 95% CI: 1.68 - 26.30). This is consistent with the study by Feirouz et al., which reported that the NNM rate is very high during the early neonatal period, with 15% occurring on the first day of life and 60% dying beyond 48 hours of life (28). Similarly, Kedy Koum et al. and Garba et al. reported hospital NNM rates of 97.1% and 63.21%, respectively, during the early neonatal period (29, 30). More than half of the births, even more, conclude with death within the first 24 hours of life, and 75% within the first week of life (31). However, the study by Zoungrana-Yameogo et al. reported that the median age of death was 2 [1-5] days and the probability of survival at the seventh day of birth was 77% (32). This similarity with other studies could be explained by the fact that the early neonatal period is very critical, as almost two-thirds of these deaths occur in this period, and to reduce early NNM, it is imperative to reduce NNM.

The probability of death occurrence was multiplied by 2.1 among newborns born to referred mothers [aOR = 2.11; (1.12 - 3.99)] (33); this implies that each level of care facility should be able to immediately provide essential care for survival, with the required skills and equipment to stabilize a newborn and safely facilitate transfer if necessary (21). In this context, the present study found a statistically significant association between mode of admission and the risk of early NNM: Emergency consultations (aOR: 4.25; 95% CI: 1.28 - 14.01); intra-hospital referral (aOR: 4.49; 95% CI: 1.59 - 12.65). In comparison with a study at the Douala referral hospital, where NNM was higher among referrals from healthcare facilities other than the maternity unit, compared to those referred from the maternity unit (29). In contrast, in the study by Mouahid et al., one-fifth of the deceased were transferred from another facility (10). This suggests a qualitative approach to understanding why these preventable deaths occur and how they are related to social determinants on one hand, hospital care provided

on the other, and the skill and motivation of health staff on the third (33).

This study reported ultra-early age, early age, intra-hospital referral, and transfer as factors associated with the risk of early NNM. Nevertheless, this study had certain limitations. The retrospective nature of our study, based on data from hospital records and registers, meant that a large amount of data was missing due to the under-notification of data, in particular socio-demographic, clinical, medical, and obstetrical data on the mothers of newborn babies. Given the high proportion of missing data for these variables, they were not investigated in our study. Otherwise, the dearth of information was also a limitation in this study, resulting in a difference between our results and those of other studies. To strengthen our findings, we recommend similar prospective research involving all the obstetric and neonatal teams over a longer period, with a comparison of the results with some regional and international studies.

### 5.1. Conclusions

Neonatal mortality is an indicator of population health. It represents a significant health challenge worldwide, particularly in developing countries where rates of NNM are predominant, such as the NNM rate at Agadir hospital, which remains high. The results of the present study highlight the need to strengthen efforts to eliminate preventable deaths and bring about significant changes in the field of maternal and child health. This can be achieved through better monitoring of pregnancy and ensuring early diagnosis, monitoring and surveillance of pregnancy, screening and management of high-risk pregnancies, better monitoring of delivery, good management of childbirth, promoting intrauterine transfer, and improving transfer conditions. Additionally, quality care for small and sick newborns is essential.

### Footnotes

**Authors' Contribution:** All authors contributed to data collection, analysis, writing of the article, and proofreading for evaluation, correction, and validation. Each author declares substantial contributions through the following: (1) the conception and design of the study, or acquisition of data, or analysis and interpretation of data; and (2) drafting the article or revising it critically for important intellectual content.

**Conflict of Interests Statement:** The authors do not declare any conflict of interest, any potential competing

or non-financial interests on behalf of all authors of the manuscript.

**Data Availability:** The dataset presented in the study is available on request from the corresponding author during submission or after publication.

**Ethical Approval:** The Ethics Committee of the Faculty of Medicine and Pharmacy of Rabat has approved this study with reference number (C64-20 ). The study strictly adhered to ethical considerations, including (a) obtaining authorization for access to records from the responsible administration; (b) preserving anonymity; (c) confidentiality; and (d) professional secrecy.

**Funding/Support:** The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

## References

- Clavandier G, Charrier P. La naissance en mutation. Un enjeu pour la sociologie ? *Recherches familiales*. 2015;(12):165-74. EN.
- OMS. *Recommandations de l'OMS Sur les soins intrapartum pour une expérience positive de l'accouchement Transformer le soins des femmes et des nourrissons pour améliorer leur santé et leur bien-être*. 2018. Available from: <https://www.who.int/fr/publications/i/item/WHO-RHR-18.12>.
- OMS. *Les femmes et les enfants sont aujourd'hui plus nombreux que jamais à avoir la vie sauve, selon un rapport des Nations Unies*. 2022. Available from: <https://www.who.int/fr/news/item/19-09-2019-more-women-and-children-survive-today-than-ever-before-un-report>.
- United Nations International Children's Emergency Fund. *Neonatal mortality*. 2021, [cited 21 August 2021]. Available from: <https://data.unicef.org/topic/child-survival/neonatal-mortality/>.
- Banque mondiale des données. *Taux de mortalité néonatale (pour 1000 naissances vivantes)*. 2022, [cited 2 October 2022]. Available from: <https://donnees.banquemondiale.org/indicateur/SH.DYN.NMRT>.
- OMS. *Nouveau-nés: améliorer leur survie et leur bien-être*. 2020, [cited 2 October 2022]. Available from: <https://www.who.int/fr/news-room/fact-sheets/detail/newborns-reducing-mortality>.
- Unicef; World Health Organization; World Bank Group. *Levels and Trends in Child Mortality Report 2019*. 2019, [cited 18 August 2021]. Available from: <https://documents1.worldbank.org/curated/en/105841568905930695/pdf/Levels-and-Trends-in-Child-Mortality-Report-2019.pdf>.
- Sante. *Rapport ENPSF 2018 2ième*. 2019, [cited 31 January 2024]. Available from: <https://www.sante.gov.ma/Documents/2020/03/Rapport%20ENPSF%202018%202i%C3%A8me%20C3%A9dition.pdf>.
- World Health Organisation. *Neonatal mortality rate(per1000 live birth)*. World Health Organisation; 2023. Available from: <https://data.who.int/indicators/i/A4C49D3>.
- Mouahid S, Chafii H, Mouhaddach O, Jaouhar S, Bouzid J. Study of risk factors of neonatal mortality in the neonatology unit at Pagnon mother-child hospital in Meknes (Morocco). *Environmental and Water Sciences, public Health and Territorial Intelligence Journal*. 2019;3(1):75-80.
- Sellouti M, Kaswati M, Barkat A. risk factors for early neonatal mortality at souissi maternity hospital rabat morocco. *Adv Pediatr Neonatal Care*. 2020;2(1).
- Doukkali L, Barkat A. *Etude de la prise en charge du Nouveau-né dans un Centre Hospitalier Provincial*. 2020. Available from: <http://ap-hm.fr/service/neonatalogie-hopital-nord>.
- Mekaoui N. *Mortalité néonatale intra hospitalière Experience de l'hôpital d'enfants de Rabat [Thesis]*. Rabat, Morocco: Université Mohammed V de Rabat; 2011.
- OMS. *Recommandations de l'OMS concernant les soins maternels et néonataux pour une expérience positive de la période postnatale: résumé analytique*. Geneva, Switzerland: World Health Organization; 2022, [cited 5 June 2022]. Available from: <https://iris.who.int/bitstream/handle/10665/354559/9789240048478-fre.pdf?sequence=1>.
- Oms; Unicef. *Mettre Fin Aux Décès De Nouveau-Nés Et Aux Mortinances Évitable D'ici 2030, Accélérer La Mise En Place D'une Couverture Sanitaire Universelle De Qualité En 2020-2025*. 2020. Available from: <https://www.who.int/fr/publications/m/item/every-newborn-coverage-target-milestones-to-2025>.
- Fleiss JL, Tytun A, Ury HK. A simple approximation for calculating sample sizes for comparing independent proportions. *Biometrics*. 1980;36(2):343-6. [PubMed ID: 26625475].
- Woodward M. Formulae for sample size, power and minimum detectable relative risk in medical studies. *Journal of the Royal Statistical Society Series D: The Statistician*. 1992;41(2):185-96.
- Unicef; World Health Organization; World Bank Group. *Levels & Trends in Child Mortality Estimates developed by the UN Inter-agency Group for Child Mortality Estimation Report 2019*. 2019, [cited 5 June 2022]. Available from: <https://www.unicef.org/media/60561/file/UN-IGME-child-mortality-report-2019.pdf>.
- Ined. *Institut national d'études démographiques*. 2022, [cited 31 August 2022]. Available from: <https://www.ined.fr/fr/lexique/mortalite-neonatale/>.
- Hôpitaux des Intérieur Règlement. *reglement intérieur des hôpitaux*. 2010, [cited 21 April 2024].
- Organisation mondiale de la Santé. *Stratégies en ressources humaines visant à améliorer la prestation des soins néonataux dans les établissements de santé des pays à revenu faible ou intermédiaire*. Geneva, Switzerland: Organisation mondiale de la Santé; 2021, [cited 28 September 2021]. Available from: <https://apps.who.int/iris/handle/10665/345348>.
- Kingwengwe Aimé A, Mwamba E, Mishika P, Nyenga Muganza A, Katamea T, Assumani N, et al. Morbi-mortalité néonatale à l'hôpital général de référence de Kindu Etat de lieux et perspectives. *Médecine d'Afrique noire*. 2019;66:569-78.
- Kambale R, Maseka A, Bwija J, Bapolisi W, Bashi J, Masumbuko B, et al. Facteurs de risque associés à la mortalité néonatale dans un hôpital de niveau tertiaire en République Démocratique du Congo. *Med Afr noire*. 2016;63(7):401-8.
- Ilunga PM, Mukuku O, Mawaw PM, Mutombo AM, Lubala TK, Wembonyama SO, et al. Fréquence et pronostic néonatal précoce de faible poids de naissance à Lubumbashi, République Démocratique du Congo. *Pan African Medical Journal*. 2016;23(1).
- Noria H, Sarah O, Asmaa O. Facteurs de risques de mortalité néonatale dans l'hôpital de gynécologie-obstétrique de la wilaya de Sidi Bel Abbes, Algérie. *The Pan African Medical Journal*. 2015;20.
- Tshinzobe JCK, Ngaya DK. Etude cas-témoins des facteurs associés à l'insuffisance pondérale à la naissance au Centre Hospitalier de Kingasani, Kinshasa (République Démocratique Congo). *Pan African Medical Journal*. 2021;38(1).
- Blondel B, Eb M, Matet N, Bréart G, Jouglu E. La mortalité néonatale en France :: bilan et apport du certificat de décès néonatal. *Archives de*

- Pédiatrie.* 2005;**12**(10):1448-55. <https://doi.org/10.1016/j.arcped.2005.05.009>.
28. Feirouz A, Sdiri Y, Cherifi E, Chourou H, Cheour M, Bel Haj Ammar W, et al. Déterminants de la mortalité néonatale précoce dans une maternité de type 3 : à propos de 150 cas. *Périnatalité.* 2021;**13**(1):26-36. <https://doi.org/10.3166/rmp-2020-0110>.
  29. Kedy Koum DC, Essomba NE, Ngaba GP, Sintat S, Ndombo PK, Coppieters Y. Morbidité et facteurs de risque de mortalité néonatale dans un hôpital de référence de Douala. *Pan African Medical Journal.* 2015;**20**. <https://doi.org/10.11604/pamj.2015.20.258.5648>.
  30. Garba M, Kamaye M, Alido S, Zoubeirou H, Oumarou Z, Amadou A. Les déterminants de la mortalité néonatale précoce à la maternité Issaka-Gazobi de Niamey. *J de Pédiatrie et de Puériculture.* 2017;**30**(4):156-61. <https://doi.org/10.1016/j.jpp.2017.01.005>.
  31. Unfpa. *Soins Obstétricaux Et Neonataux Durgence Guide Pour La Réalisation Des Enquetes Rapides.* 2022, [cited 9 October 2022]. Available from: <https://wcaro.unfpa.org/sites/default/files/pub-pdf/UNFPA-WCARO-ER-SONU-F5-2%20%282%29.pdf>.
  32. Zoungrana-Yameogo WN, Dahourou DL, Diallo AH, Sangho O, Nikiema E, Tougouma S, et al. Mortalité néonatale au centre hospitalier universitaire de Tengandogo, Ouagadougou, Burkina Faso: une étude de cohorte rétrospective. *Journal of Interventional Epidemiology and Public Health.* 2021;**4**(3). <https://doi.org/10.37432/jieph.suppl.2021.4.3.03.4>.
  33. Barkat Radouani MA, Barkat A. Social Determinants of Perinatal Health in Morocco; In: Laher I, editor. *Handbook of Healthcare in the Arab World.* Cham: Springer; 2021.