

Antimicrobial Effects of Folk Medicinal Plants From the North of Iran Against *Mycobacterium tuberculosis*

Shadi Jahanpour¹; Kiumars Ghazisaidi¹; Homa Davoodi¹; Masoumeh Mazandarani²; Motahare Samet¹; Nadia Jahanpour³; Ezzat Allah Ghaemi^{1,*}

¹Infectious Disease Research Center, Golestan University of Medical Sciences, Gorgan, IR Iran

²Islamic Azad University, Gorgan Branch, Gorgan, IR Iran

³Islamic Azad University, Lahijan Branch, Lahijan, IR Iran

*Corresponding author: Ezzat Allah Ghaemi, Infectious Disease Research Center, Golestan University of Medical Sciences, Gorgan, IR Iran. Tel: +98-9113711770, Fax: +98-1732245597, E-mail: eghaemi@yahoo.com

Received: February 9, 2014; Revised: May 18, 2014; Accepted: August 23, 2014

Background: Medicinal plants have been used traditionally in Golestan province (north of Iran), against *Mycobacterium tuberculosis* or the clinical signs of tuberculosis (TB).

Objectives: This study aimed to define the inhibitory effects of ethanolic extracts of six of these medicinal plants against *Mycobacterium tuberculosis*.

Materials and Methods: *Peganum harmala* (seed extract), *Punica granatum* (peel extract), *Digitalis* sp. (leaf extract), fruit extract of *Citrus lemon*, *Rosa canina* and *Berberis vulgaris* were extracted in ethanol and their activity against *M. tuberculosis* isolates were determined by the agar diffusion method. The zone of inhibition (at 200 to 1.6 mg/mL) was measured and the results were compared with isoniazid and rifampin as standard positive controls. Also the concentration of vitamin C of each the extracts was evaluated.

Results: The ethanolic extract of *Peganum harmala* seed and *Punica granatum* peel exhibited potential activity against all *M. tuberculosis* isolates with mean inhibitory zone of 18.7 and 18.8 mm, at 200 mg/mL concentration. The mean inhibitory zone around isoniazid and rifampin were 19.2 and 18.8 mm. Ethanolic extract of *Citrus lemon* showed moderate inhibitory activity only against sensitive (non MDR; non multi drug resistant) strains of *M. tuberculosis*, and *Digitalis* sp. showed inhibitory effects on five isolates. Ascorbic acid content was 43.3 mg/dL in *Punica granatum* and *Digitalis* sp. and only 9.1 mg/dL in ethanolic extract of *Peganum harmala*.

Conclusions: The highest content of vitamin C was observed in the extract of *Punica granatum*, which was observed to be highly active against *Mycobacterium tuberculosis*, while the *P. harmala* must have contained other phytochemical constituents that contributed to the anti-tuberculosis effects of this plant. Our findings showed that ethanolic extracts of *P. granatum* and *P. harmala* had anti-TB effects comparable to isoniazid and rifampin and can be good candidates for novel and safe natural products against tuberculosis.

Keywords: Medicinal Plants; *Mycobacterium tuberculosis*; Vitamin C; In Vitro

1. Background

Tuberculosis (TB) is one of the major fatal infectious diseases in the world (1). It has infected one-third of the world's population and is responsible for approximately 2 million deaths each year, mostly in developing countries (2). In 2010, approximately 8.8 million new cases of active TB was reported and 1.1 million people died from the disease, most of which happened in developing countries (3). Tuberculosis typically affects the lungs and spreads from an infected person to a susceptible person (4). This infection requires a long duration of therapy, which is six months for first-line drugs, including isoniazid, rifampin, ethambutol and pyrazinamide (5). In certain situations (for example, bacterial drug resistance and intolerance), second-line drugs are used, mostly based on high-risk medications, such as fluoroquinolones, capreomycin, ethionamide and cycloserine, that are usually less effective and more toxic with side ef-

fects for the patient (6). The improper use of these drugs results in increasing prevalence of multi-drug resistant (MDR) and extensively drug resistant (XDR) strains of *M. tuberculosis* that are resistant to both first and second-line drugs (7). Therefore there is an urgent need to search for complementary medicines or new alternatives of antibiotics, especially from natural sources, which work more effectively with fewer side effects (8, 9). Medicinal plants have been used for this purpose in many countries to prevent and treat infectious diseases (10). Antioxidants such as vitamin C, tocopherols, carotenoids, polyphenols and flavonoids may be responsible for the antibacterial activity of medicinal plants. Many studies have found an association between anti-tuberculosis activity of medicinal plant extracts and their vitamin C content, which can be an important finding (11). Tuberculosis has been a threatening disease in Iran from the distant past. According to

the world health organization, the estimated incidence rate of TB in the Islamic Republic of Iran was 21 cases per 100000 people in 2011. The Golestan province, located in the south-east of the Caspian sea has the highest incidence of TB in Iran (12). This province due to its diverse climatic and heterogeneous ecological condition is able to grow many plants, especially medicinal plants, and the search for finding medicinal plants that inhibit the growth of *Mycobacterium tuberculosis* is of great importance in this region.

2. Objectives

The present study aimed to investigate the anti-*M. tuberculosis* activity of six medicinal plants in the region against *Mycobacterium tuberculosis* for the management of the tuberculosis disease.

3. Materials and Methods

3.1. Preparation of Plant Samples

Six medicinal plants were selected for the present study. The selected plants namely *Citrus lemon* (Rutaceae), *Peganum harmala* (Zygophyllaceae), *Punica granatum* (Lythraceae), *Rosa canina* (Rosaceae), *Berberis vulgaris* (Berberidaceae), *Digitalis* sp. (Scrophulariaceae) had shown anti-mycobacterial activity against non-pathogenic mycobacteria in a previous study (13). The selected plants were collected from ranges as high as 650-2250 m and from three regions Ziaret, Deraznoo and Charbagh, located south of Golestan province from July to September 2012. Table 1 presents the scientific name, local names and their uses in traditional medicine.

The method of Alade and Irobi (14) was used for the extraction of plant extracts with minor modifications. Briefly the plants were air-dried in the dark at room temperature and then ground into fine powder using an electric blender. Plant hydro-alcoholic extracts were prepared by the maceration method; 30 g of each dried powder was soaked in 100 mL of 70% ethanol for 72 hours. A sterile glass rod was used for stirring each mixture every 24 hours. The plant extracts were filtered twice through Whatman filter paper no. 1 (Whatman, UK). The obtained

filtrates were evaporated using a rotary evaporator. The final products were sticky-brown substances and stored in a laboratory refrigerator prior to the test. The concentrated extract was freshly dissolved in dimethyl sulfoxide (DMSO) before use.

3.2. Preparation of Inoculums

In the present study a total of seven isolates of *Mycobacterium tuberculosis* were used; six clinical isolates were identified by conventional methods including two MDR and four sensitive isolates to rifampin and isoniazid, which were isolated from sputum of tuberculosis patients in Gorgan. Also a drug-susceptible strain of *M. tuberculosis* (H37Rv), which was sensitive to rifampin and isoniazid, was kindly provided by Dr. Hashemi (Ahvaz University of Medical Sciences, Ahvaz, Iran). The bacterial cultures, which were used to prepare the standard inoculums, were maintained on Lowenstein-Jensen medium (Merck). A suspension of each bacterium was prepared in normal saline containing 0.05% Tween-80 and adjusted approximately to McFarland 0.5 standard (15).

3.3. Anti *M. tuberculosis* Activity

Antibacterial activity was determined by the disk diffusion method (16, 17). One hundred microliters of mycobacterial suspension was spread onto the surface of Lowenstein Jensen media, using a sterile glass spreader. The extracts were dissolved in 2% DMSO (Sigma) to obtain a concentration of 200 mg/mL and further diluted to their final concentrations. The final concentration of DMSO in all assays was 2% or less, which is nontoxic for mycobacteria (18). Sterile paper disks (6 mm diameter) were loaded with 40 µL of extract (varying concentrations: 200, 100, 50, 25, 12.5, 6.25, 3.12 and 1.6 mg/mL) and then left to dry. The impregnated-disks were then placed on the surface of plates on which the microorganisms were cultured. Rifampin (MAST) and Isoniazid (SIGMA-ALDRICH) was used as the positive control and disks treated with 2% DMSO were used as the negative control. The culture plates were sealed in plastic bags and incubated at 37°C for three weeks, after which culture-growth was clearly visible on the agar. The clear zone visible all around the

Table 1. Ethnobotanical Information of Selected Medicinal Plants

Plants	Local Name	Family	Part(s) Used	Preparation Used	Therapeutic Use
<i>Citrus lemon</i>	Limoo	Rutaceae	fruit pulp	decoction, powder, fresh juice	cold, fever, flu, bronchitis
<i>Peganum harmala</i>	Esphand	zygophyllaceae	seed	soaked eaten	respiratory and nervous system diseases, cough, stomachache
<i>Punica granatum</i>	Anar	Punicaceae	peel	cooked raw	antibiotic, urinary tract infections, anti-inflammation, cold, flu
<i>Rosa canina</i>	Nastaran	Rosaceae	fruit	decoction	preventive for flu
<i>Berberis vulgaris</i>	Zereshk	Berberidaceae	fruit	decoction	dysentery, sedative, malaria, anti-inflammatory
<i>Digitalis</i> sp.	Angoshtane	Scrophulariaceae	Leaf	fresh water of Leaf, liquid extract	anti-inflammatory, anti-oxidant, antipyretic, antirheumatic, stimulatory for the immune system

impregnated-disk was indicative of the inhibitory effects. The diameter of the zone of inhibition around each of the disks was measured and recorded. Each experiment was performed in triplicates.

3.4. Measurement of Vitamin C

The concentration of vitamin C in each extract was measured using the colorimetric method (19). In short, 0.2 g of vitamin C powder as the standard and 500 µL of the stock solutions of each plant extract were mixed with metaphosphoric acid 6g/dL and centrifuged at 3000 rpm for 10 minutes. Next, 1.2 mL of the supernatant from each of the tubes were combined with a mixture containing copper sulphate 0.6%, thiourea 5% and diphenyl hydrazine 2%. After three hours of incubation in a water bath at 37°C, 2 mL of sulphuric acid 12 M was added to each tube. The content of vitamin C in each sample was measured by reading the absorbance in a spectrophotometer at OD of 520 nm. This experiment was carried out in duplicates.

4. Results

4.1. Anti *M. tuberculosis* Activity

Amongst the hydro-alcoholic extracts of the six selected medicinal plants, fruit extracts of *Berberis vulgaris* and *Rosa canina* showed no inhibitory activity against *Myco-*

bacterium tuberculosis. Extract of *Punica granatum* even at concentration of 25 mg/mL could prevent the growth of all *M. tuberculosis* isolates while the extract of *Peganum harmala* had this property at a concentration of 50 mg/mL (Table 2). Ethanolic extract of *Citrus lemon* could prevent the growth of non-MDR *Mycobacterium tuberculosis* isolates while the ethanolic extract of *Digitalis sp.* showed a mild anti-*M. tuberculosis* activity at concentrations of 200 and 100 mg/mL. Isoniazid and rifampin showed inhibitory effects only against non-MDR *M. tuberculosis* isolates. The zone of growth inhibition and the number of isolates which showed inhibitory effects were decreased with decreasing concentrations of ethanolic extracts, so that *Citrus lemon* extract at concentration of 200 and 25 mg/mL inhibited the growth of four and one *M. tuberculosis* isolates, respectively. Although, ethanolic extract of *Digitalis sp.* at a concentration of 200 mg/mL could inhibit the growth of five *M. tuberculosis* isolates, none of isolates were inhibited at a concentration of 50 mg/mL (Table 2).

Anti-TB activity was shown by extracts of *Punica granatum* and *Peganum harmala* on sensitive and resistant strains of *Mycobacterium tuberculosis*. The maximum zone of inhibition against an MDR strain was observed with extract of *Punica granatum* (19.5 ± 3.5 mm) and against a non-MDR strain by extract of *Peganum harmala* (19 ± 2.6 mm) as compared with the standard rifampin (19.5 ± 2.4 mm) and isoniazid (19.5 ± 2.1 mm).

Table 2. Anti-Tuberculosis Activity of Selected Medicinal Plants Against Multi-Drug Resistant and Non-Multi-Drug Resistant *M. tuberculosis*^a

Concentrations, mg/mL	H37RV	MDR ₁	MDR ₂	Non MDR ₁	Non MDR ₂	Non MDR ₃	Non MDR ₄	Mean	
								MDR	Non MDR
Citrus lemon									
200	10	R	R	20	20	R	10	R	12.50 ± 9.6
100	R	R	R	15	15	R	R	R	7.50 ± 8.7
50	R	R	R	8	10	R	R	R	4.50 ± 5.3
25	R	R	R	R	8	R	R	R	2.00 ± 4.0
Peganumharmala									
200	18	20	17	18	22	20	16	18.50 ± 2.1	19.00 ± 2.6
100	15	16	15	15	15	15	12	15.50 ± 1	14.25 ± 1.5
50	8	12	10	14	12	12	10	11.00 ± 1.4	12.00 ± 1.6
25	R	10	8	12	8	10	9	9.00 ± 1.4	9.75 ± 1.7
Punicagaranatium									
200	20	22	17	18	15	18	20	19.50 ± 3.5	17.75 ± 2.1
100	17	18	13	15	13	15	16	15.50 ± 3.5	14.75 ± 1.3
50	15	15	10	13	10	14	13	12.50 ± 3.5	12.50 ± 1.7
25	13	10	8	10	8	10	8	9.00 ± 1.4	9.00 ± 1.2
Digitalis Sp.									
200	R	R	12	12	12	12	14	6.00 ± 6	12.50 ± 1
100	R	R	R	10	10	10	10	R	10
50	R	R	R	R	R	R	R	R	R
25	R	R	R	R	R	R	R	R	R
Isoniazid, µg/mL									
2	18	R	R	17	22	20	19	R	19.50 ± 2.1
Rifampin, µg/mL									
5	16	R	R	18	23	19	18	R	19.50 ± 2.4

^a Abbreviations: MDR, multiple-drug resistant; Non MDR, non-multiple-drug resistant.

Table 3. Level of Vitamin C of Selected Medicinal Plants

Name of the Plant	OD	Concentration(mg/dL)
<i>Punica granatum</i>	0.300	43.3
<i>Digitalis sp.</i>	0.300	43.3
<i>Berberis vulgaris</i>	0.285	41.1
<i>Rosa canina</i>	0.145	20.9
<i>Citrus lemon</i>	0.078	11.3
<i>Peganum harmala</i>	0.063	9.1

4.2. Measurement of Vitamin C Level

The level of vitamin C in ethanolic extracts of the selected plants are presented in Table 3. Extract of *Punica granatum* and *Digitalis sp.* with vitamin C amount 43.3 mg/dL showed the maximum level of vitamin C.

5. Discussion

In the present study, selection of medicinal plants was based on a previous study, conducted by Ghaemi et al. (13). The plants were used for the treatment of tuberculosis, chronic respiratory diseases or symptoms of diseases, by local healers in this area. Ghaemi et al. reported that among the 52 herbal plants examined, only six extracts had good anti-mycobacterial activity against two non-pathogenic strains of *M. bovis* BCG and *M. smegmatis* (13). Therefore, the present study aimed to evaluate the inhibitory effects of these six medicinal plants on the growth of *Mycobacterium tuberculosis*, as the most important pathogenic mycobacteria. Among these plants only the ethanolic extracts of *Peganum harmala* and *Punica granatum* showed potential inhibitory effects against *M. tuberculosis*. Similarly, Gautam et al. (20) found anti-mycobacterial activity by ethanolic extracts of *P. harmala* and *P. granatum*. Also Ghaemi et al. showed that ethanolic extracts of *P. granatum* have greater effects on *M. bovis* compared to *M. smegmatis*. Pomegranate is an edible fruit, which natively grows in Iran and many studies have been done on its biological properties that confirm the therapeutic properties of this plant. Pomegranate has a wide range of bioactive compounds such as: alkaloids, ellagic acid, Punicalagin, ellagitannins, anthocyanin, flavonoids and tannins. Due to presence of these compounds it has various pharmacological activities such as, antioxidant, anti-microbial and anti-virus (21) and has been used to treat respiratory diseases (22). In addition to its antibacterial activities against *Escherichia coli* (23) and *Staphylococcus aureus* (24), we found that the peel extracts of pomegranate have a great potential to be used as anti-TB agents. The ethanolic extract of *Peganum harmala* followed by the extract of *Punica granatum* showed strong anti-TB activity against *M. tuberculosis*. In two studies, conducted by Gautam et al. and Ghaemi et al. it was demonstrated that the ethanolic extract of this plant was active against *M. smegmatis* (13, 20) and *M. bovis* (13). *Peganum harmala* is known as “Espand” in Iran (25) and has been used traditionally in Turkey, Iran and China to treat coughs, rheu-

matism, high blood pressure, diabetes and asthma (26). The seed and root extracts of *P. harmala* have broad antibacterial activity against gram-positive and negative-bacteria (25). The most pharmacological active compounds of *P. harmala* are alkaloids (27). Ghaemi et al. reported that ethanolic extracts of *Citrus lemon* have strong inhibitory activity against *M. smegmatis* but limited inhibitory activity on *M. bovis*. In another study, Kirbaslar et al. (28) showed that *Citrus lemon* peel oil was active against *M. phlei* and *M. smegmatis*, as indicated by the disk diffusion assay with an inhibitory zone equal to 13 mm. Saeed et al. (29) reported that *Citrus lemon* juice had antibacterial activity against a range of gram-positive and negative bacteria. *Citrus lemon* has been used traditionally as an anti-inflammatory, antiseptic, expectorant and relaxing agent (13). In the present study we found that this extract has moderate activity against *M. tuberculosis*. Thus, high concentrations of this extract may be effective in controlling the growth of *M. tuberculosis*. It is interesting to note that the extract of *Citrus lemon* only showed inhibitory activity against isoniazid and rifampin-sensitive isolates. *Citrus lemon* is a native plant of India that belongs to the Rutaceae family and is widely cultivated in Iran including the Golestan province, located north of Iran. Anti-bacterial, anti-fungal, anti-viral, anti-cancer and anti-diabetic activity of this plant is due to the presence of flavonoids and alkaloids in its crude extracts (30). Similar to the results obtained by Ghaemi et al. (13), in this study, we found weak activity for ethanolic extracts of *Digitalis sp.* against *M. tuberculosis*, therefore in vitro anti-tuberculosis activity of this plant is not remarkable. This plant may be effective in other ways, such as boosting the immune system to inhibit the growth of TB in the body. Recently, potential activity of vitamin C has been shown to be part of the dramatic killing of sensitive and resistant strains of *Mycobacterium tuberculosis* in vitro (11). Narwadiya et al. (31) supports the fact that as the concentration of vitamin C increases in plant extracts there is evidence of decrease in the number of tubercle bacilli; a claim that is somewhat confirmed by our result. Our findings demonstrated that *Punica granatum* (peel) has a high concentration of vitamin C and showed maximum activity against *M. tuberculosis*, yet *Digitalis sp.* (leaf) although it has a high concentration of vitamin C, it did not show anti-tuberculosis activity. On the other hand, *Peganum harmala* (seed) had the minimum concentration of Vitamin C among all examined plants of this study, yet it showed potential activity against the tubercle bacilli. Therefore, vitamin C by itself is not sufficient for the inhibitory effect of extracts of medicinal plants against *M. tuberculosis*. The reason behind these observations needs further investigations. The peel of ethanolic extracts of *Punica granatum* and the seed extracts of *Peganum harmala* exhibited significant inhibitory activity in vitro. According to our findings, we can conclude that the high concentration of vitamin C in the *P. granatum* extract can justify its anti-tuberculosis effect. On the other hand, *P. harmala* extract does not

contain enough vitamin C, thus it may contain other phytochemical constituents that contribute to the anti-tuberculosis effects in this plant. These medicinal plants as promising candidates and may be used as novel natural products without any side effects for new anti-TB drug discovery. The application of these two extracts either in cell culture or animal models is recommend, to evaluate their growth inhibitory effects on *M. tuberculosis*.

Acknowledgements

We thank the management of the Medicinal Plants Department of Islamic Azad University of Gorgan. Also, we are grateful for the assistance of Mrs M. Tazike and Mrs S. Rahimialang, the experts at the mycobacteriology laboratory of Golestan University of Medical Sciences and the Deputy of Health for providing the necessary facilities to carry out this research. Furthermore great thanks to Mis Fatemeh Shakeri and Naemeh Javid for their kindly cooperation.

Authors' Contributions

Shadi Jahanpour: MSc student, doing experimental work and manuscript preparation; Kiumars Ghazisaidi: scientific advisor; Homa Davoodi; technical advisor; Maesoumeh Mazandarani, cooperation in collecting and identifying the herbal medicine and extracts preparation; Motahare Samet, cooperation in extract preparation; Nadia Jahanpour, cooperation in collecting herbal medicine; Ezzat Allah Ghaemi, director of the research project and corresponding author.

Financial Disclosure

This proposal was confirmed by Deputy of Research and Technology with this letter number; 35/1925, 28/09/2011.

Funding/Support

This study was financially supported by Infectious Disease Research Center, Deputy of Research and Technology, Golestan University of Medical Sciences Gorgan, Iran.

References

1. Grange JM, Zumla A. The global emergency of tuberculosis: what is the cause? *J R Soc Promot Health*. 2002;**122**(2):78–81.
2. Crane LR. *History of Tuberculosis*.; 2013.
3. World Health Organization. *Global tuberculosis programme. Global tuberculosis control*. WHO; 2011. Available from: <http://www.who.int/gtb/publications>.
4. Dye C, Scheele S, Dolin P, Pathania V, Raviglione MC. Consensus statement. Global burden of tuberculosis: estimated incidence, prevalence, and mortality by country. WHO Global Surveillance and Monitoring Project. *JAMA*. 1999;**282**(7):677–86.
5. Zhang Y, Yew WW. Mechanisms of drug resistance in Mycobacterium tuberculosis. *Int J Tuberc Lung Dis*. 2009;**13**(11):1320–30.
6. Blumberg HM, Burman WJ, Chaisson RE, Daley CL, Etkind SC, Friedman LN, et al. American Thoracic Society/Centers for Disease Control and Prevention/Infectious Diseases Society of America: treatment of tuberculosis. *Am J Respir Crit Care Med*. 2003;**167**(4):603–62.
7. World Health Organization. *Global tuberculosis control: epidemiology, strategy, financing*. WHO; 2009. Available from: http://www.who.int/tb/publications/global_report/2009/pdf/full_report.pdf.
8. Newman DJ, Cragg GM, Snader KM. Natural products as sources of new drugs over the period 1981–2002. *J Nat Prod*. 2003;**66**(7):1022–37.
9. Newman DJ, Cragg GM, Snader KM. The influence of natural products upon drug discovery. *Nat Prod Rep*. 2000;**17**(3):215–34.
10. Windisch W, Schedle K, Plitzner C, Kroismayr A. Use of phyto-genic products as feed additives for swine and poultry. *J Anim Sci*. 2008;**86**(14 Suppl):E140–8.
11. Vilcheze C, Hartman T, Weinrick B, Jacobs WR, Jr. Mycobacterium tuberculosis is extraordinarily sensitive to killing by a vitamin C-induced Fenton reaction. *Nat Commun*. 2013;**4**:1881.
12. World Health Organization. *Global tuberculosis programme Global tuberculosis control*. WHO; 2011. Available from: <http://www.who.int/gtb/publications>.
13. Antimycobacterial activity of some medicinal plants used in traditional medicine in North of Iran. In: Ghaemi EA, Mazandarani M, Mansourian AR, Babaii Kochaksaraei M editors. *Proceedings of International Conference on Life Science and Technology*. 2011 Iran.
14. Alade PI, Irobi ON. Antimicrobial activities of crude leaf extracts of *Acalypha wilkesiana*. *J Ethnopharmacol*. 1993;**39**(3):171–4.
15. Chaturvedi V, Dwivedi N, Tripathi RP, Sinha S. Evaluation of Mycobacterium smegmatis as a possible surrogate screen for selecting molecules active against multi-drug resistant Mycobacterium tuberculosis. *J Gen Appl Microbiol*. 2007;**53**(6):333–7.
16. Ahmad I, Beg AZ. Antimicrobial and phytochemical studies on 45 Indian medicinal plants against multi-drug resistant human pathogens. *J Ethnopharmacol*. 2001;**74**(2):113–23.
17. Japoni Nejad A, Sofian M, van Belkum A, Ghaznavi Rad E. Nosocomial Outbreak of Extensively and Pan Drug-Resistant Acinetobacter baumannii in Tertiary Hospital in Central Part of Iran. *Jundishapur J Microbiol*. 2013;**6**(8).
18. Molina-Salinas GM, Ramos-Guerra MC, Vargas-Villarreal J, Mata-Cardenas BD, Becerril-Montes P, Said-Fernandez S. Bactericidal activity of organic extracts from *Flourensia cernua* DC against strains of Mycobacterium tuberculosis. *Arch Med Res*. 2006;**37**(1):45–9.
19. Tietz NW, Burtis CA, Ashwood ER. *Tietz textbook of clinical chemistry*. Philadelphia: Saunders; 1994.
20. Gautam R, Saklani A, Jachak SM. Indian medicinal plants as a source of antimycobacterial agents. *J Ethnopharmacol*. 2007;**110**(2):200–34.
21. Jasuja ND, Saxena R, Chandra S, Sharma R. Pharmacological Characterization and Beneficial Uses of *Punica granatum*. *Asian J Plant Sci*. 2012;**11**(6):251–67.
22. Khan AJ, Haneef S. Antibacterial properties of *Punica granatum* peels. *Int J Appl Biol Pharm Technol*. 2011;**2**(3):23–7.
23. Voravuthikunchai SP, Sririrak T, Limsuwan S, Supawita T, Iida T, Honda T. Inhibitory effects of active compounds from *Punica granatum* pericarp on verocytotoxin production by enterohemorrhagic *Escherichia coli* O157: H7. *J Health sci*. 2005;**51**(5):590–6.
24. Machado TB, Pinto AV, Pinto MC, Leal IC, Silva MG, Amaral AC, et al. In vitro activity of Brazilian medicinal plants, naturally occurring naphthoquinones and their analogues, against methicillin-resistant *Staphylococcus aureus*. *Int J Antimicrob Agents*. 2003;**21**(3):279–84.
25. Darabpour E, Motamedi H, Poshtkouhian Bavi A, Seyyed Nejad SM. *Eldorado Resources for and from Research Teaching and Studying*.; 2011.
26. Shao H, Huang X, Zhang Y, Zhang C. Main alkaloids of *Peganum harmala* L. and their different effects on dicot and monocot crops. *Molecules*. 2013;**18**(3):2623–34.
27. Asgarpanah J, Ramezanloo F. Chemistry, pharmacology and medicinal properties of *Peganum harmala* L. *Afr J Pharm Pharmacol*. 2012;**6**(22):1573–80.
28. Kirbaslar GF, Tavman A, Dulger B, Turker G. Antimicrobial activity of Turkish citrus peel oils. *Pak J Bot*. 2009;**41**(6):3207–12.
29. Saeed S, Tariq P. Effects of some seasonal vegetables and fruits on the growth of bacteria. *Pak J Biol Sci*. 2006;**9**(8):1547–51.

30. Dhanavade MJ, Jalkute CB, Ghosh JS, Sonawane KD. Study antimicrobial activity of Lemon (*Citrus lemon L.*) peel extract. *Br J Pharmacol Toxicol.* 2011;**2**(3):119–22.
31. Narwadiya SC, Sahare KN, Tumane PM, Dhumne UL, Meshram VG. In vitro anti-tuberculosis effect of vitamin C contents of medicinal plants. *Asian J Exp Biol Sci.* 2011;**2**(1):151–4.