



Depression and Anxiety Among Children and Adolescents During the COVID-19 Outbreak: A Systematic Review

Maryam Seyedtabib ¹, Masoomah Asadi ², Fatemeh Maghsoodi ³ and Naser Kamyari ^{4,*}

¹Department of Biostatistics and Epidemiology, School of Health, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

²Department of Operating Room Nursing, Abadan University of Medical Sciences, Abadan, Iran

³Department of Public Health, School of Health, Abadan University of Medical Sciences, Abadan, Iran

⁴Department of Biostatistics and Epidemiology, School of Health, Abadan University of Medical Sciences, Abadan, Iran

*Corresponding author: Department of Biostatistics and Epidemiology, School of Health, Abadan University of Medical Sciences, Abadan, Iran. Email: n.kamyari@abadanums.ac.ir

Received 2023 January 05; Revised 2023 March 06; Accepted 2023 July 22.

Abstract

Context: During the outbreak of COVID-19, mental health problems were increasingly observed in children and adolescents, leading to negative physical and psychological consequences. This pandemic provided a unique opportunity for conducting survey studies on this topic. As there is no comprehensive research in this area, our aim was to provide a concise overview of previous studies on the effects of the COVID-19 pandemic on the mental well-being of young people.

Methods: We performed an electronic search in multiple databases for this systematic review. The quality of the studies was assessed on the basis of a 14-item quality assessment tool proposed by the National Heart, Lung, and Blood Institutes (NHLBI) for observational cohort and cross-sectional studies. The time span encompassed December 2019 to December 2021, and the databases included PubMed, Scopus, ScienceDirect, and Google Scholar. The search was based on the keywords: COVID-19, depression, anxiety, children, and adolescents. Based on our inclusion criteria, out of the 4516 articles reviewed, only eight papers were included in the final review, and their results were summarized and reported.

Results: The negative mental health consequences of children during the coronavirus outbreak included increased anxiety and depression, increased behavioral and emotional problems, experiencing misbehaviors and emotional bullying, and increased anxiety and stress following harm, especially in girls.

Conclusions: Considering the negative impact of the COVID-19 pandemic on children's mental health, it seems necessary to conduct interventional studies to improve and reduce stress and anxiety in children and adolescents during future disease outbreaks.

Keywords: Depression, Anxiety, Children, Adolescent, COVID-19

1. Context

COVID-19 is a systemic respiratory disorder caused by the SARS-CoV-2 virus. The acute respiratory syndrome of COVID-19 may manifest with mild to severe respiratory symptoms, multiple organ failure, and death (1). This disease started in China at the end of 2019, spread rapidly worldwide, and became a major health crisis (1). The WHO (World Health Organization) declared the COVID-19 disease a significant threat to physical and mental health (2). COVID-19 has caused various problems, such as fear of infection (3-5) and death anxiety (6). As a result, it has affected various physical, psychological, and even economic aspects of life (7). Fear, anxiety, mood changes, and depression are common in this disease, and depression (8) and mood disorders (9)

are also prevalent. Anxiety is a very unpleasant and vague feeling accompanied by one or more physical symptoms such as shortness of breath, palpitations, sweating, headache, restlessness, and the urge to move. According to diagnostic guidelines and mental disorder statistics, anxiety is one of the most common mental health problems (10, 11). Anxiety disorder is also the most common clinical problem during childhood and adolescence, and approximately 5% of children and adolescents fulfill the diagnostic criteria for an anxiety disorder during childhood. More than 20% of people experience anxiety problems before the age of 16 years (12, 13). Therefore, childhood and adolescence are high-risk periods for developing anxiety. In addition, spontaneous recovery from anxiety symptoms is low in children

and adolescents, presenting a continuous problem throughout the growth period (14-16).

During the COVID-19 pandemic, children and teenagers were assumed to be one of the most vulnerable groups in society (17). Quarantine inflicted many physical and psychological effects on children (18). In this context, several studies have investigated the psychological disorders of people during quarantine. The results of a study demonstrated that during the COVID-19 pandemic and after the closure of schools in China, more than 180 million students were quarantined at home. This restriction during the outbreak of COVID-19 caused symptoms such as anxiety and depression in children (19, 20). Ghosh et al.'s study showed that during the outbreak of COVID-19, children suffered from different psychological problems (21). Zhao et al. reported that children often had more behavioral and emotional problems during the COVID-19 outbreak due to forced isolation and closure of schools, reduced activities, and reduced communication with their friends and classmates (22). Also, during the pandemic, quarantine at home, exposure to domestic violence, and reduced opportunities for playing and entertainment severely affected children's mental health (23, 24). Compared to adults, children and adolescents are less severely affected by COVID-19, and most of them have mild symptoms and a good prognosis. Nevertheless, the disruption of family support, increased fear of losing dependents, and the death of a family member can lead to severe mental health problems in children, leading to changes in their adaptation capabilities and mood. This phenomenon provides the basis for stress, anxiety, depression, and even suicidal thoughts in children (25).

Children are one of the vulnerable groups in society, and their mental health problems, besides tourniquet complications and costs, can threaten the functioning of the family and society (26-28). Therefore, it is important to timely recognize stress and anxiety in children during disease outbreaks, such as the COVID-19 pandemic, so that families, professionals, and psychologists can implement effective preventive measures, educational plans, and therapeutic interventions. Already, various studies have examined the consequences of the COVID-19 disease and its mental health consequences in children. Therefore, the present systematic review aims to investigate the severity of depression, anxiety, and stress in children and adolescents during the COVID-19 pandemic.

2. Methods

This systematic review was conducted and organized according to the PRISMA (Preferred Reporting Items for

Systematic reviews and Meta-Analyses) guideline (29). We focused on quantitative studies published during the COVID-19 pandemic on changes in anxiety and depression in children and adolescents.

2.1. Data Sources and Search Strategy

A comprehensive electronic search was conducted in multiple databases from December 2019 to December 2021. The language was limited to English. The databases included Embase, Cochrane, PubMed, Web of Science, Scopus, and Google Scholar. We also applied the technique of "backward snowballing" to find other relevant articles (30) by reviewing the reference lists of the selected articles. Controlled vocabularies and keywords describing mental health problems of children and adolescents in the context of the COVID-19 pandemic were used for the search. In accordance with the topic of our study, the keywords that were searched included "depression", "anxiety", "children", "adolescents", and "COVID-19". Each keyword was searched individually. In similar reviews or other meta-analyses containing these keywords, we checked if there were synonymous terms with different meanings. We then integrated these words into the search formula to extend the search scope. The following search syntax was used: ("mental health" OR "mental illness*" OR "mental disorder*" OR "psychological illness*" OR "psychiatric disorder*" OR "depress*" OR "depressive disorder*" OR "depressive symptom*" OR "emotional depression" OR "affective disorder*" OR "mood disorder*" OR "anxiety" OR "angst" OR "nervousness" OR "hypervigilance" OR "social anxiety" OR "anxiety disorder*" OR "anxious" OR "overanxious disorder*") AND ("adolescent*" OR "teenager*" OR "child*" OR "student*" OR "juvenile*" OR "school*" OR "teen*" OR "young" OR "youth*" OR "minors") AND ("COVID-19" OR "coronavirus" OR "SARS-CoV-2" OR "cov-19" OR "2019-nCoV" OR "severe acute respiratory syndrome Coronavirus2"). Two authors (NK and MS) independently searched all six databases using the same syntax to ensure that no relevant studies were missed.

2.2. Inclusion and Exclusion Criteria

All survey studies investigating depression and anxiety in children and adolescents in the context of COVID-19 were included in this review. In accordance with the age definitions proposed by the WHO and the United Nations, we included people aged 0-19 years (children and adolescents) (31, 32). All types of survey data, including questionnaires and interviews, reporting the prevalence of depression or anxiety or reporting depression or anxiety scores in different countries were considered. There were no restrictions regarding the type of intervention,

the comparison, or data analysis methods. Articles that exclusively reported data on participants older than 19 years, studies on populations with clinical conditions (patients with COVID-19 or other diseases), those unrelated to our epidemiological and clinical aspects of interest (i.e., psychological consequences), and studies not written in English were excluded.

2.3. Study Selection and Data Extraction

Two authors (MA and FM) independently reviewed the titles and abstracts of all publications and read their full text for qualification. The information of each study, including the first author's name, country, publication date, population under study, study design, sample size, age, interventions, type of depression or anxiety, and outcomes, was extracted and recorded into a password-protected Microsoft Excel spreadsheet. If the two researchers disagreed, a third party (the lead researcher, MS) intervened in the discussion and resolved any disagreement. When articles presented results from the same study, we selected those that reported outcomes that most closely matched our eligibility criteria. We also carefully reviewed the references of the included studies to find potentially relevant research.

2.4. Quality Assessment

The quality of the studies was assessed using a 14-item quality assessment tool presented in the National Heart, Lung, and Blood Institutes (NHLBI) study for observational cohort and cross-sectional studies (33). These items included the statistical power of the study, the strength of the causal relationship between the exposure and outcomes, and the potential risk of selection bias, information bias, measurement bias, or confounding bias. The studies were then categorized as high, moderate, or low quality, and the overall rating of each study was determined (see Appendix 1 in the Supplementary File).

2.5. Data Synthesis

In order to ensure accurate reporting, we followed the PRISMA checklist in this review. Given the diversity of the data from the initial studies, we created descriptive summary tables using Microsoft Excel version 2016 (Microsoft Corp, Redmond, WA, USA) to provide a concise overview of the literature.

3. Results

We identified 4,516 potentially relevant articles. After removing duplicates and irrelevant records, 3,879 articles remained. After screening titles and abstracts, we excluded

3,871 articles due to the lack of fulfillment of inclusion criteria in at least one of the following steps: (1) title screening, (2) abstract screening, and (3) retrieving of results. When reading the full text of the remaining 242 articles, 75 papers were removed for not having useful data, 58 for being conducted on participants older than 19 years, 72 for reporting insufficient data, and 29 for insufficient information. Ultimately, 8 eligible studies were selected to be comprehensively reviewed here. The screening process is summarized in a PRISMA 2020 flowchart (Figure 1).

The results of all studies that met the inclusion criteria were carefully reviewed. A summary of the content of these articles is provided in Table 1. The objectives, measurements, and conclusions of these studies are summarized in Table 2.

The spread of the COVID-19 disease nurtured psychological consequences and anxiety in children and adolescents. The review of the studies conducted in this field showed that the most important consequences were as follows.

3.1. Behavioral and Emotional Problems

The closure of schools, quarantine at home, fear of contracting the COVID-19 infection, and lack of communication with classmates decreased physical activities and social interactions of children and adolescents. Almost all these events cause or intensify behavioral and emotional problems in children and adolescents (37, 44).

3.2. Anxiety and Stress

In epidemics like COVID-19, fear of disease, death, and disruption in daily functioning cause anxiety due to fear of contagious diseases. The spread of the COVID-19 disease and continuous quarantine rules overexposed families to sad pandemic news and children and teenagers to fear and anxiety. The fear of losing parents, the rapid spread of the disease, the fear and anxiety of contracting the disease, as well as the fear of a painful death were mentioned in studies (45). According to studies (39, 40, 44, 46-48), COVID-19 has a high prevalence and increases stress and anxiety in parents and families.

3.3. Depression and Suicidal Thoughts

Concerns over depression and suicidal thoughts increased during the pandemic, especially among female teenagers. The high and unexpected rise in the prevalence of anxiety and depression requires an urgent solution to reduce harm and ensure students' mental health in the next epidemic crises. Studies suggest continuous screening for depression and suicidal thoughts among




Figure 1. The selection procedure is shown in the PRISMA 2020 flowchart. It is recommended, if possible, to specify the number of records identified from each individual database or register searched and not the total number from all databases or registers. If automation tools were used, it is also important to indicate the number of records excluded by humans and the number of records excluded by automation tools. This information was taken from (34, 35). Further details can be found at: <http://www.prisma-statement.org>.

Table 1. Characteristics of Reviewed Trials^a

Authors	Year	Country	Design	Ethics Approval	Informed Consent	Sample Size	Population	Age, y	Results	
									Anxiety	Depression
Giannopoulou et al. (36)	2020	Greece	Cross-sectional	Yes	Yes	442	High school students	Senior high school students (SHSS)	10.59 ± 5.79	13.59 ± 7.25
Burdzovic Andreas and Brunborg (37)	2020	Norway	Longitudinal	Yes	Yes	915	Adolescents students	8th -10th grade		8.58 ± 7.41
Belanger et al. (38)	2020	Canada	Longitudinal	Yes	Yes	2099	Students in grades 9th -12th	14.1 ± 1.0	6.23 ± 5.44	
Black et al. (39)	2020	United States	Cross-sectional	No	Yes	362	Children under the age of 18 years	10.68 ± 2.28		12.02 ± 5.13
Westrupp et al. (40)	2020	Australia	Cross-sectional	Yes	Not report (NR)	2,365	Children aged 0 -18 years	8.66 ± 5.14	3.04 ± 2.65	5.04 ± 12.35
Mayne et al. (41)	2020	United States	Cross-sectional	No	No	47,684	Adolescents aged 12 -21 years	15.3 ± 2.1		10,929 (22.92)
Gladstone et al. (42)	2021	United States	Longitudinal	Yes	Not report (NR)	228	Adolescents in 7th, 9th and 11th grades	14.5 ± 1.60		4.37 ± 4.88
Jolliff et al. (43)	2021	United States	Cross-sectional	Yes	Yes	134	Teenagers aged 13 -17 years	15.2 ± 1.4	5.37 ± 5.18	5.27 ± 5.06

^a Values are expressed as mean ± SD or No. (%)

adolescents and high school students is a crucial task (36, 41).

3.4. Abuse and Emotional Abuse

The relationship between parents and children is crucial for creating safety and affection. The closure and bankruptcy of some businesses at the peak of the outbreak created many economic problems for some families, causing family disputes and increasing tensions and arguments in families, which would often lead to the mistreatment of children (49). Economic problems and family conflicts can affect the interaction between parents and children (50). Therefore, the more severe the psychological problems of parents, such as anxiety and stress, the higher the risk of behavioral disorders, such as aggression and anxiety, in children (21).

3.5. Depression and Post-traumatic Stress

Being quarantined at home during the COVID-19 outbreak was traumatic for many children and teenagers and might have led to loneliness and depression in this sensitive society group. Children and adolescents who lose a family member due to COVID-19 show the symptoms of post-traumatic stress or acute stress (42, 48).

Most studies have reported the adverse effects of COVID-19 on the mental health of children and adolescents. However, some studies also argued that neither being inflicted with COVID-19 disease nor experiencing changes in daily activities due to COVID-19 disease was associated with the exaggeration of depression or anxiety compared to before the pandemic. However, we found that young people with a lower socioeconomic status (SES) were more

anxious during the pandemic than their peers with a higher SES. According to these results, instrumental and psychological interventions are necessary for low-income teenagers (38, 43).

4. Discussion

The present systematic review aimed to determine the severity of depression and anxiety in children and adolescents under 19 years of age during the outbreak of COVID-19. The current study demonstrated that COVID-19 could be associated with increased depression, anxiety, and stress among children and adolescents. The results of this research align with the findings of a review study by Viner et al. (51), who reported that the closure of schools increased psychological problems in children and adolescents (51). In this context, mental health support and similar interventions seem necessary. The results of various studies showed that in critical and emergency conditions, such as the coronavirus epidemic, the effects of the disease not only cause mental health problems in children and adolescents but also lead to persistent negative emotions (22, 52-55). Also, during disease outbreaks, the resultant economic and social consequences and financial damages to families will cause many problems for society and families, leading to fear, tension, frustration, and anxiety (17, 24, 56).

Generally, various studies indicate that children and teenagers are less vulnerable than adults to COVID-19 (57), but psychologically and socially, this group is more vulnerable than adults (58-60). Fear of illness, excruciating death, rumors, interruption in daily

Table 2. Aims, Measurements, and Conclusions of Included Studies

Authors	Study's Aims	Measurement Tools	Conclusions
Giannopoulou et al. (36)	Investigating how the nationwide lockdown affected anxiety and depression in high school students preparing for the upcoming national college entrance exams in Greece.	Anxiety (GAD-7), Depression (PHQ-9)	The surprisingly high levels of anxiety and depression required immediate action to address and manage students' mental health issues in future pandemic outbreaks.
Burdzovic Andreas and Brunborg (37)	Investigating the self-reported mental and physical well-being of adolescents in Norway before and during the pandemic and the health impact of pandemic-related anxiety.	Depression (PHQ-9)	The adolescents in the group affected by COVID-19, who felt very anxious due to the pandemic, were much more likely to have poorer mental and physical health compared to the pre-pandemic period.
Belanger et al. (38)	Assessing the impact of the initial response to COVID-19 on adolescents' mental well-being and general health in a real-world study.	Anxiety (GAD-7)	The early phases of COVID-19 lockdown measures had no negative impact on the mental health of young people. The decline in mental well-being during the first COVID-19 wave was not as bad as the decline before the pandemic.
Black et al. (39)	Investigating the impact of COVID-19 on the relationship between depressive symptoms and irritability in children and their parents.	Depression (MFQ-A)	Stress caused by COVID-19 was associated with higher levels of symptoms in parents and children, and it was suggested that family dynamics could play a role in interactions between these symptoms.
Westrupp et al. (40)	To examine trends in parents' and children's (0-18 years) mental health, parental substance use, marital conflicts, parenting practice, and family dynamics during COVID-19 compared to pre-pandemic data and to identify families that were particularly vulnerable to negative outcomes caused by demographic and individual factors as well as stressors caused by COVID-19.	Anxiety (SCAS), Depression (MFQ)	When it comes to children's mental health, existing parental health problems and the psychological and environmental stress caused by COVID-19 were repeatedly linked to increased anxiety and depression symptoms in children. This was observed in cases where the child was diagnosed with ADHD or ASD.
Mayne et al. (41)	Explaining changes in screening, signs of depression, and suicide risk in teenagers during the COVID-19 pandemic.	Depression (PHQ-9)	Depression and suicidal thoughts increased among female teens during the pandemic, underscoring the need for regular screening for depression and suicidal thoughts, as well as for consistent mental health monitoring of this population.
Gladstone et al. (42)	To investigate how the COVID-19 pandemic affected the depressive feelings of a group of young people and to examine the relationship between the stress caused by COVID-19 and factors that make them more or less vulnerable to depression, and how these factors contributed to changes in depressive symptoms over time.	Depression (PHQ-8)	The study investigated how major negative life events could influence the risk of depression in adolescents in their future lives. Because depression can have lasting effects on adolescents and because negative life events during adolescence can predict depression risk in adulthood, it is important that mental health professionals and school personnel be alert to the potential rise in depression in adolescents during the pandemic.
Jolliff et al. (43)	Comparing the levels of depression and anxiety in two groups of teenagers in the United States before and during the COVID-19 pandemic and examining the correlation of demographic factors and daily activities with depression and anxiety.	Anxiety (GAD-7), Depression (PHQ-8)	Adolescents who were part of the pandemic group or for whom the pandemic led to changes in their daily activities did not show higher levels of depression or anxiety. However, it was found that adolescents from less privileged backgrounds had significantly higher levels of anxiety during the pandemic than their more advantaged peers. It is important to offer low-income young people both practical support and psychological help.

Abbreviations: GAD, general anxiety disorder-7; PHQ, Patient Health Questionnaire-9; MFQ, Mood and Feelings Questionnaire, SCAS, Spence Children's Anxiety Scale; ADHD, attention-deficit/hyperactivity disorder; ASD, autism spectrum disorder; SES, socioeconomic status.

activities, prohibitions or restrictions on travel and commute, financial problems, limitation of social interactions, psychological burden and pressure on families, lack of playing during home quarantine, staying away from the school, lack of communication with friends and classmates, having no entertainment, and domestic violence all threaten the mental health of children and teenagers (57-63). Among the most common psychological consequences were the stressful situation caused by the COVID-19 disease and the lack of information about effective and efficient treatments, exaggerating anxiety

(11, 64, 65).

Strengths and limitations: One of the strengths of this review was that the eight papers analyzed included more than 54,000 children and adolescents aged 0 - 19 years in five countries worldwide. They provided data on various mental health issues and evaluated changes in mental health before and after the pandemic. However, this study also had some limitations. Due to variabilities in study designs and mental health outcomes, no meta-analysis was performed. In addition, only peer-reviewed English-language papers were included,

excluding studies in other languages. Although eight studies were included, only one presented a representative population-based data set, limiting the assessment of anxiety and depression on a larger scale. In addition, there were few studies on children under the age of ten years, making the extrapolation of findings to this age group difficult. Finally, all reviewed studies focused on changes in the early phase of the pandemic, so future research should focus on the impact of later waves of the pandemic on the mental health of children and young people.

4.1. Conclusions

Several survey studies have shown a strong link between the COVID-19 pandemic (and its consequences, such as social distancing, quarantine, and school closure) and mental health problems in children and adolescents. Anxiety, depression, stress, loneliness, and tension have been frequently reported. It is crucial to urgently apply positive coping strategies and provide family and social support to these individuals, as recommended in clinical practice guidelines. However, existing evidence is limited, and there is a need for more comprehensive studies, including intervention-based research, on this topic.

Supplementary Material

Supplementary material(s) is available [here](#) [To read supplementary materials, please refer to the journal website and open PDF/HTML].

Acknowledgments

We thank the research deputy of the Abadan University of Medical Sciences for financially supporting this project.

Footnotes

Authors' Contribution: NK and MS obtained the funding and developed the idea for the review article; MA and FM performed the literature search and screening; NK, MS, MA, and FM drafted the manuscript; MA, FM, MS, and NK critically revised the manuscript.

Conflict of Interests: The authors declare that they have no conflict of interest.

Ethical Approval: The article does not contain clinical studies or patient data. The Research Ethics Committee of Abadan University of Medical Sciences approved this study with a specific code [IR.ABADANUMS.REC.1401.098](#).

Funding/Support: The Abadan University of Medical Sciences supported this work under grant number 1555.

References

- Zhou P, Yang XL, Wang XG, Hu B, Zhang L, Zhang W, et al. A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature*. 2020;579(7798):270-3. [PubMed ID: 32015507]. [PubMed Central ID: PMC7095418]. <https://doi.org/10.1038/s41586-020-2012-7>.
- Shen K, Yang Y, Wang T, Zhao D, Jiang Y, Jin R, et al. Diagnosis, treatment, and prevention of 2019 novel coronavirus infection in children: experts' consensus statement. *World J Pediatr*. 2020;16(3):223-31. [PubMed ID: 32034659]. [PubMed Central ID: PMC7090771]. <https://doi.org/10.1007/s12519-020-00343-7>.
- Protudjer JLP, Golding M, Salisbury MR, Abrams EM, Roos LE. High anxiety and health-related quality of life in families with children with food allergy during coronavirus disease 2019. *Ann Allergy Asthma Immunol*. 2021;126(1):83-88 e1. [PubMed ID: 32950685]. [PubMed Central ID: PMC7498414]. <https://doi.org/10.1016/j.anaai.2020.09.010>.
- Nikolaidis A, Paksarian D, Alexander L, Derosa J, Dunn J, Nielson DM, et al. The Coronavirus Health and Impact Survey (CRISIS) reveals reproducible correlates of pandemic-related mood states across the Atlantic. *Sci Rep*. 2021;11(1):8139. [PubMed ID: 33854103]. [PubMed Central ID: PMC8046981]. <https://doi.org/10.1038/s41598-021-87270-3>.
- Fitzpatrick O, Carson A, Weisz JR. Using Mixed Methods to Identify the Primary Mental Health Problems and Needs of Children, Adolescents, and Their Caregivers during the Coronavirus (COVID-19) Pandemic. *Child Psychiatry Hum Dev*. 2021;52(6):1082-93. [PubMed ID: 33108612]. [PubMed Central ID: PMC7590914]. <https://doi.org/10.1007/s10578-020-01089-z>.
- Shanafelt T, Ripp J, Trockel M. Understanding and Addressing Sources of Anxiety Among Health Care Professionals During the COVID-19 Pandemic. *JAMA*. 2020;323(21):2133-4. [PubMed ID: 32259193]. <https://doi.org/10.1001/jama.2020.5893>.
- Li W, Yang Y, Liu ZH, Zhao YJ, Zhang Q, Zhang L, et al. Progression of Mental Health Services during the COVID-19 Outbreak in China. *Int J Biol Sci*. 2020;16(10):1732-8. [PubMed ID: 32226291]. [PubMed Central ID: PMC7098037]. <https://doi.org/10.7150/ijbs.45120>.
- Nshimiyimana R, Guzzetta CE, Brown MM, Zhou Q, Johnson JM, Sato T, et al. Pilot study of anxiety, depression, and quality of life in patients with the diagnosis of metastatic uveal melanoma. *Annals of Oncology*. 2018;29:viii560. <https://doi.org/10.1093/annonc/mdy296.011>.
- Davila J, Starr LR, Stroud CB, Li Y. Mood and anxiety disorders. In: Fiese BH, Celano M, Deater-Deckard K, Jouriles EN, Whisman MA, editors. *APA handbook of contemporary family psychology: Applications and broad impact of family psychology*. Washington DC, USA: American Psychological Association; 2019. p. 21-36.
- Muris P, Simon E, Lijphart H, Bos A, Hale W3, Schmeitz K, et al. The Youth Anxiety Measure for DSM-5 (YAM-5): Development and First Psychometric Evidence of a New Scale for Assessing Anxiety Disorders Symptoms of Children and Adolescents. *Child Psychiatry Hum Dev*. 2017;48(1):1-17. [PubMed ID: 27179521]. [PubMed Central ID: PMC5243875]. <https://doi.org/10.1007/s10578-016-0648-1>.
- Luo D, Liu Q, Chen Q, Huang R, Chen P, Yang BX, et al. Mental Health Status of the General Public, Frontline, and Non-frontline Healthcare Providers in the Early Stage of COVID-19. *Front Psychiatry*. 2021;12:553021. [PubMed ID: 33815157]. [PubMed Central ID: PMC8012479]. <https://doi.org/10.3389/fpsy.2021.553021>.
- Chaabane S, Chaabna K, Bhagat S, Abraham A, Doraiswamy S, Mamtani R, et al. Perceived stress, stressors, and coping strategies among nursing students in the Middle East and North Africa: an overview of systematic reviews. *Syst Rev*. 2021;10(1):136. [PubMed ID: 33952346]. [PubMed Central ID: PMC8101235]. <https://doi.org/10.1186/s13643-021-01691-9>.

13. Kocsis RN. Book Review: Diagnostic and Statistical Manual of Mental Disorders: Fifth Edition (DSM-5. *Int J Offender Ther Comp Criminol*. 2013;**57**(12):1546–8. <https://doi.org/10.1177/0306624X13511040>.
14. Ueda R, Okada T, Kita Y, Ozawa Y, Inoue H, Shioda M, et al. The quality of life of children with neurodevelopmental disorders and their parents during the Coronavirus disease 19 emergency in Japan. *Sci Rep*. 2021;**11**(1):3042. [PubMed ID: 33589663]. [PubMed Central ID: PMC7884401]. <https://doi.org/10.1038/s41598-021-82743-x>.
15. Teubert D, Pinquart M. A meta-analytic review on the prevention of symptoms of anxiety in children and adolescents. *J Anxiety Disord*. 2011. <https://doi.org/10.1016/j.janxdis.2011.07.001>.
16. McCormick BF, Connolly EJ, Nelson DV. Mild Traumatic Brain Injury as a Predictor of Classes of Youth Internalizing and Externalizing Psychopathology. *Child Psychiatry Hum Dev*. 2021;**52**(1):166–78. [PubMed ID: 32372378]. <https://doi.org/10.1007/s10578-020-00992-9>.
17. Anticich SA, Barrett PM, Silverman W, Lacherez P, Gillies R. The prevention of childhood anxiety and promotion of resilience among preschool-aged children: a universal school based trial. *Adv Sch Ment Health Promot*. 2013;**6**(2):93–121. <https://doi.org/10.1080/1754730x.2013.784616>.
18. Cheng FF, Zhan SH, Xie AW, Cai SZ, Hui L, Kong XX, et al. Anxiety in Chinese pediatric medical staff during the outbreak of Coronavirus Disease 2019: a cross-sectional study. *Transl Pediatr*. 2020;**9**(3):231–6. [PubMed ID: 32775241]. [PubMed Central ID: PMC7347772]. <https://doi.org/10.21037/tp.2020.04.02>.
19. Qu P, Zhao D, Jia P, Dang S, Shi W, Wang M, et al. Changes in Mental Health of Women Undergoing Assisted Reproductive Technology Treatment During the COVID-19 Pandemic Outbreak in Xi'an, China. *Front Public Health*. 2021;**9**:645421. [PubMed ID: 34113596]. [PubMed Central ID: PMC8185191]. <https://doi.org/10.3389/fpubh.2021.645421>.
20. Koksal E, Dost B, Terzi O, Ustun YB, Ozdin S, Bilgin S. Evaluation of Depression and Anxiety Levels and Related Factors Among Operating Theater Workers During the Novel Coronavirus (COVID-19) Pandemic. *J Perianesth Nurs*. 2020;**35**(5):472–7. [PubMed ID: 32855053]. [PubMed Central ID: PMC7303603]. <https://doi.org/10.1016/j.jopan.2020.06.017>.
21. Ghosh R, Dubey MJ, Chatterjee S, Dubey S. Impact of COVID -19 on children: special focus on the psychosocial aspect. *Minerva Pediatr*. 2020;**72**(3):226–35. [PubMed ID: 32613821]. <https://doi.org/10.23736/S0026-4946.20.05887-9>.
22. Zhao X, Lan M, Li H, Yang J. Perceived stress and sleep quality among the non-diseased general public in China during the 2019 coronavirus disease: a moderated mediation model. *Sleep Med*. 2021;**77**:339–45. [PubMed ID: 32482485]. [PubMed Central ID: PMC7240276]. <https://doi.org/10.1016/j.sleep.2020.05.021>.
23. Fegert JM, Vitiello B, Plener PL, Clemens V. Challenges and burden of the Coronavirus 2019 (COVID-19) pandemic for child and adolescent mental health: a narrative review to highlight clinical and research needs in the acute phase and the long return to normality. *Child Adolesc Psychiatry Ment Health*. 2020;**14**:20. [PubMed ID: 32419840]. [PubMed Central ID: PMC7216870]. <https://doi.org/10.1186/s13034-020-00329-3>.
24. Bate J, Malberg N. Containing the Anxieties of Children, Parents and Families from a Distance During the Coronavirus Pandemic. *J Contemp Psychother*. 2020;**50**(4):285–94. [PubMed ID: 32836378]. [PubMed Central ID: PMC7358677]. <https://doi.org/10.1007/s10879-020-09466-4>.
25. Park E, Logan H, Zhang L, Kamigaichi N, Kulapichitr U. Responses to Coronavirus Pandemic in Early Childhood Services Across Five Countries in the Asia-Pacific Region: OMEP Policy Forum. *Int J Early Child*. 2020;**52**(3):249–66. [PubMed ID: 33518791]. [PubMed Central ID: PMC7829089]. <https://doi.org/10.1007/s13158-020-00278-0>.
26. Holling H, Schlack R, Petermann F, Ravens-Sieberer U, Mauz E, Ki GG. [Psychopathological problems and psychosocial impairment in children and adolescents aged 3-17 years in the German population: prevalence and time trends at two measurement points (2003-2006 and 2009-2012): results of the KiGGS study: first follow-up (KiGGS Wave 1)]. *Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz*. 2014;**57**(7):807–19. German. [PubMed ID: 24950830]. <https://doi.org/10.1007/s00103-014-1979-3>.
27. Han L, Wong FKY, She DLM, Li SY, Yang YF, Jiang MY, et al. Anxiety and Depression of Nurses in a North West Province in China During the Period of Novel Coronavirus Pneumonia Outbreak. *J Nurs Scholarsh*. 2020;**52**(5):564–73. [PubMed ID: 32652884]. [PubMed Central ID: PMC7405411]. <https://doi.org/10.1111/jnu.12590>.
28. Adibelli D, Sumen A. The effect of the coronavirus (COVID-19) pandemic on health-related quality of life in children. *Child Youth Serv Rev*. 2020;**119**:105595. [PubMed ID: 33071408]. [PubMed Central ID: PMC7550976]. <https://doi.org/10.1016/j.childyouth.2020.105595>.
29. Moher D, Liberati A, Tetzlaff J, Altman DG, Prisma Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Ann Intern Med*. 2009;**151**(4):264–9. W64. [PubMed ID: 19622511]. <https://doi.org/10.7326/0003-4819-151-4-200908180-00135>.
30. Wohlin C. Guidelines for snowballing in systematic literature studies and a replication in software engineering. In: Shepperd M, editor. *EASE '14: Proceedings of the 18th International Conference on Evaluation and Assessment in Software Engineering*. New York, SA. Association for Computing Machinery; 2014. p. 1–10.
31. *Recognizing adolescence*. World Health Organization; 2020, [cited 2020 Jul 1]. Available from: <https://apps.who.int/adoles-cent/second-decade/section2/page1/recognizing-adolescence.html>.
32. *Convention on the Rights of the Child*. UN General Assembly; 2003, [cited 2021 Apr 29]. Available from: https://web.archive.org/web/20101031104336/http://www.hakani.org/en/convention/Convention_Rights_Child.pdf.
33. *Study Quality Assessment Tools: quality assessment tool for observational cohort and cross-sectional studies*. National Heart Lung and Blood Institute; 2021. Available from: <https://www.nhlbi.nih.gov/health-topics/study-quality-assessment-tools>.
34. Subirana M, Sola I, Garcia JM, Gich I, Urrutia G. A nursing qualitative systematic review required MEDLINE and CINAHL for study identification. *J Clin Epidemiol*. 2005;**58**(1):20–5. [PubMed ID: 15649667]. <https://doi.org/10.1016/j.jclinepi.2004.06.001>.
35. Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ*. 2021;**372**:n71. [PubMed ID: 33782057]. [PubMed Central ID: PMC8005924]. <https://doi.org/10.1136/bmj.n71>.
36. Giannopoulou I, Efstathiou V, Triantafyllou G, Korkoliakou P, Douzenis A. Adding stress to the stressed: Senior high school students' mental health amidst the COVID-19 nationwide lockdown in Greece. *Psychiatry Res*. 2021;**295**:113560. [PubMed ID: 33187723]. [PubMed Central ID: PMC7647459]. <https://doi.org/10.1016/j.psychres.2020.113560>.
37. Burdzovic Andreas J, Brunborg GS. Self-reported Mental and Physical Health Among Norwegian Adolescents Before and During the COVID-19 Pandemic. *JAMA Netw Open*. 2021;**4**(8):e2121934. [PubMed ID: 34427678]. [PubMed Central ID: PMC8385591]. <https://doi.org/10.1001/jamanetworkopen.2021.21934>.
38. Belanger RE, Patte KA, Leatherdale ST, Ganssone RJ, Haddad S. An Impact Analysis of the Early Months of the COVID-19 Pandemic on Mental Health in a Prospective Cohort of Canadian Adolescents. *J Adolesc Health*. 2021;**69**(6):917–24. [PubMed ID: 34565667]. [PubMed Central ID: PMC8457891]. <https://doi.org/10.1016/j.jadohealth.2021.07.039>.
39. Black SR, Evans ML, Aaron L, Brabham DR, Kaplan RM. Covariance Between Parent and Child Symptoms Before and During the COVID-19 Pandemic. *J Pediatr Psychol*. 2021;**46**(10):1182–94. [PubMed ID: 34405876]. [PubMed Central ID: PMC8385985]. <https://doi.org/10.1093/jpepsy/jsab086>.
40. Westrupp EM, Bennett C, Berkowitz T, Youssef GJ, Toumbourou

- JW, Tucker R, et al. Child, parent, and family mental health and functioning in Australia during COVID-19: comparison to pre-pandemic data. *Eur Child Adolesc Psychiatry*. 2023;**32**(2):317-30. [PubMed ID: 34417875]. [PubMed Central ID: PMC8379590]. <https://doi.org/10.1007/s00787-021-01861-z>.
41. Mayne SL, Hannan C, Davis M, Young JF, Kelly MK, Powell M, et al. COVID-19 and Adolescent Depression and Suicide Risk Screening Outcomes. *Pediatrics*. 2021;**148**(3). [PubMed ID: 34140393]. <https://doi.org/10.1542/peds.2021-051507>.
 42. Gladstone TRG, Schwartz JA, Posselt P, Richer AM, Buchholz KR, Rintell LS. Depressive Symptoms Among Adolescents: Testing Vulnerability-Stress and Protective Models in the Context of COVID-19. *Child Psychiatry Hum Dev*. 2022;**53**(6):1372-82. [PubMed ID: 34235630]. [PubMed Central ID: PMC8262586]. <https://doi.org/10.1007/s10578-021-01216-4>.
 43. Jolliffe A, Zhao Q, Eickhoff J, Moreno M. Depression, Anxiety, and Daily Activity Among Adolescents Before and During the COVID-19 Pandemic: Cross-sectional Survey Study. *JMIR Form Res*. 2021;**5**(12). e30702. [PubMed ID: 34609316]. [PubMed Central ID: PMC8641699]. <https://doi.org/10.2196/30702>.
 44. Zhang J, Shuai L, Yu H, Wang Z, Qiu M, Lu L, et al. Acute stress, behavioural symptoms and mood states among school-age children with attention-deficit/hyperactive disorder during the COVID-19 outbreak. *Asian J Psychiatr*. 2020;**51**:102077. [PubMed ID: 32315967]. [PubMed Central ID: PMC7195413]. <https://doi.org/10.1016/j.ajp.2020.102077>.
 45. Liu JJ, Bao Y, Huang X, Shi J, Lu L. Mental health considerations for children quarantined because of COVID-19. *Lancet Child Adolesc Health*. 2020;**4**(5):347-9. [PubMed ID: 32224303]. [PubMed Central ID: PMC7118598]. [https://doi.org/10.1016/S2352-4642\(20\)30096-1](https://doi.org/10.1016/S2352-4642(20)30096-1).
 46. Dalton L, Rapa E, Stein A. Protecting the psychological health of children through effective communication about COVID-19. *Lancet Child Adolesc Health*. 2020;**4**(5):346-7. [PubMed ID: 32243784]. [PubMed Central ID: PMC7270522]. [https://doi.org/10.1016/S2352-4642\(20\)30097-3](https://doi.org/10.1016/S2352-4642(20)30097-3).
 47. Zolfaghari A, Elahi T. [Children's level of anxiety in relation to their level of awareness and attitude towards corona virus based on the health belief model and the level of stress, anxiety and depression of mothers]. *J Res Psychol Health*. 2020;**14**(1):40-55. Persian.
 48. Roccella M. Children and Coronavirus Infection (Covid-19): What to Tell Children to Avoid Post-traumatic Stress Disorder (PTSD). *Open Pediatr Med J*. 2020;**10**(1):1-2. <https://doi.org/10.2174/1874309902010010001>.
 49. Altafim ERP, Linhares MBM. Universal violence and child maltreatment prevention programs for parents: A systematic review. *Psychosoc Interv*. 2016;**25**(1):27-38. <https://doi.org/10.1016/j.psi.2015.10.003>.
 50. Wang C, Pan R, Wan X, Tan Y, Xu L, Ho CS, et al. Immediate Psychological Responses and Associated Factors during the Initial Stage of the 2019 Coronavirus Disease (COVID-19) Epidemic among the General Population in China. *Int J Environ Res Public Health*. 2020;**17**(5). [PubMed ID: 32155789]. [PubMed Central ID: PMC7084952]. <https://doi.org/10.3390/ijerph17051729>.
 51. Viner RM, Russell SJ, Croker H, Packer J, Ward J, Stansfield C, et al. School closure and management practices during coronavirus outbreaks including COVID-19: a rapid systematic review. *Lancet Child Adolesc Health*. 2020;**4**(5):397-404. [PubMed ID: 32272089]. [PubMed Central ID: PMC7270629]. [https://doi.org/10.1016/S2352-4642\(20\)30095-X](https://doi.org/10.1016/S2352-4642(20)30095-X).
 52. Yue J, Zang X, Le Y, An Y. Anxiety, depression and PTSD among children and their parent during 2019 novel coronavirus disease (COVID-19) outbreak in China. *Curr Psychol*. 2022;**41**(8):5723-30. [PubMed ID: 33223783]. [PubMed Central ID: PMC7666617]. <https://doi.org/10.1007/s12144-020-01191-4>.
 53. Yildirim M, Ozaslan A, Arslan G. Perceived risk and parental coronavirus anxiety in healthcare workers: a moderated mediation role of coronavirus fear and mental well-being. *Psychol Health Med*. 2022;**27**(5):1095-106. [PubMed ID: 33410335]. <https://doi.org/10.1080/13548506.2021.1871771>.
 54. Sritharan J, Sritharan A. Emerging Mental Health Issues from the Novel Coronavirus (COVID-19) Pandemic. *J Health Med Sci*. 2020;**3**(2):157-62. <https://doi.org/10.31014/aior.1994.03.02.109>.
 55. Rodriguez-Rey R, Garrido-Hernansaiz H, Collado S. Psychological Impact and Associated Factors During the Initial Stage of the Coronavirus (COVID-19) Pandemic Among the General Population in Spain. *Front Psychol*. 2020;**11**:1540. [PubMed ID: 32655463]. [PubMed Central ID: PMC7325630]. <https://doi.org/10.3389/fpsyg.2020.01540>.
 56. Bahn GH. Coronavirus Disease 2019, School Closures, and Children's Mental Health. *Soa Chongsonyon Chongsin Uihak*. 2020;**31**(2):74-9. [PubMed ID: 32595345]. [PubMed Central ID: PMC7289477]. <https://doi.org/10.5765/jkacap.200010>.
 57. Mangolian Shahrabaki P, Dehghan M, Maazallahi M, Asadi N. Fear and anxiety in girls aged 7 to 11 years old and related factors during the coronavirus pandemic. *Clin Child Psychol Psychiatry*. 2022;**27**(1):259-68. [PubMed ID: 33951970]. [PubMed Central ID: PMC8829152]. <https://doi.org/10.1177/13591045211013873>.
 58. Duan L, Shao X, Wang Y, Huang Y, Miao J, Yang X, et al. An investigation of mental health status of children and adolescents in china during the outbreak of COVID-19. *J Affect Disord*. 2020;**275**:112-8. [PubMed ID: 32658812]. [PubMed Central ID: PMC7329661]. <https://doi.org/10.1016/j.jad.2020.06.029>.
 59. Dreher A, Pietrowsky R, Loerbroks A. Pandemic-related attitudes, stressors and work outcomes among medical assistants during the SARS-CoV-2 ("Coronavirus") pandemic in Germany: A cross-sectional Study. *PLoS One*. 2021;**16**(1). e0245473. [PubMed ID: 33444419]. [PubMed Central ID: PMC7808691]. <https://doi.org/10.1371/journal.pone.0245473>.
 60. Cheng C, Wang HY, Chan L. Multiple forms of mass anxiety in coronavirus disease-2019 pandemic. *J Affect Disord*. 2021;**291**:338-43. [PubMed ID: 34087629]. [PubMed Central ID: PMC8460399]. <https://doi.org/10.1016/j.jad.2021.05.034>.
 61. Wu Z, McGoogan JM. Characteristics of and Important Lessons From the Coronavirus Disease 2019 (COVID-19) Outbreak in China: Summary of a Report of 72314 Cases From the Chinese Center for Disease Control and Prevention. *JAMA*. 2020;**323**(13):1239-42. [PubMed ID: 32091533]. <https://doi.org/10.1001/jama.2020.2648>.
 62. Valero-Moreno S, Lacomba-Trejo L, Casana-Granell S, Prado-Gasco VJ, Montoya-Castilla I, Perez-Marin M. Psychometric properties of the questionnaire on threat perception of chronic illnesses in pediatric patients. *Rev Lat Am Enfermagem*. 2020;**28**. e3242. [PubMed ID: 32022154]. [PubMed Central ID: PMC7000189]. <https://doi.org/10.1590/1518-8345.3144.3242>.
 63. Bajema KL, Oster AM, McGovern OL, Lindstrom S, Stenger MR, Anderson TC, et al. Persons Evaluated for 2019 Novel Coronavirus - United States, January 2020. *MMWR Morb Mortal Wkly Rep*. 2020;**69**(6):166-70. [PubMed ID: 32053579]. [PubMed Central ID: PMC7017962]. <https://doi.org/10.15585/mmwr.mm6906e1>.
 64. Sarner M. Maintaining mental health in the time of coronavirus. *New Sci*. 2020;**246**(3279):40-6. [PubMed ID: 32372828]. [PubMed Central ID: PMC7194563]. [https://doi.org/10.1016/S0262-4079\(20\)30819-8](https://doi.org/10.1016/S0262-4079(20)30819-8).
 65. Luca DL, Margiotta C, Staatz C, Garlow E, Christensen A, Zivin K. Financial Toll of Untreated Perinatal Mood and Anxiety Disorders Among 2017 Births in the United States. *Am J Public Health*. 2020;**110**(6):888-96. [PubMed ID: 32298167]. [PubMed Central ID: PMC7204436]. <https://doi.org/10.2105/AJPH.2020.305619>.