



# The Effect of Slow Stroke Back Massage on Sleep Quality in Hemodialysis Patients: A Clinical Trial Study

Mohammad Zayer<sup>1</sup>, Masumeh Akbarbegloo <sup>1,\*</sup>, Fatemeh Mehraji<sup>1</sup>

<sup>1</sup> Student Research Committee, Khoy University of Medical Sciences, Khoy, Iran

\*Corresponding Author: Khoy University of Medical Sciences, Khoy, Iran. Email: m.akbarbegloo@yahoo.com

Received: 7 July, 2024; Revised: 22 November, 2025; Accepted: 24 January, 2025

## Abstract

**Background:** Hemodialysis patients frequently experience sleep disturbances, substantially influencing their quality of life. Safe and non-invasive therapies such as massage are essential in mitigating the adverse effects of sleep deprivation.

**Objectives:** This study aimed to assess the impact of slow stroke back massage (SSBM) on the sleep quality of hemodialysis patients.

**Methods:** This clinical trial included 60 patients selected through convenient sampling in Khoy Shahid Madani Hospital, Iran (2023), and randomly assigned to intervention and control groups. The intervention group received back massages thrice a week for 10 minutes over one month. The control group continued with routine care. Sleep quality was assessed using the Pittsburgh Sleep Quality Index (PSQI) before, and 15 and 30 days after the intervention. Data were analyzed using SPSS version 19.

**Results:** Repeated measures analysis showed a significant improvement in sleep quality in the intervention group compared to baseline and on the 15th and 30th days ( $P < 0.001$ ). On the 30th day, the mean sleep quality score was  $10.19 \pm 5.39$  in the intervention group compared to  $12.55 \pm 6.2$  in the control group. The mean changes in sleep quality scores in the intervention group were  $0.73 \pm 4.2$  on day 15 and  $1.63 \pm 3.9$  on day 30. The independent  $t$ -test confirmed that sleep quality significantly improved in the intervention group compared to the control group ( $P < 0.001$ ).

**Conclusions:** The findings suggest that back massage using the SSBM is a safe and cost-effective intervention to enhance relaxation and improve sleep quality in hemodialysis patients.

**Keywords:** Slow Stroke Back Massage, Sleep Quality, Hemodialysis

## 1. Background

Chronic kidney disease (CKD) is a progressive, irreversible condition in which the body's ability to maintain fluid and electrolyte balance is compromised, leading to uremia and azotemia (1). According to data from the centers for disease control (CDC), approximately 52% of advanced CKD patients undergo hemodialysis (2). In Iran, the prevalence of diagnosed CKD patients is about 253 per million population (3). Hemodialysis patients suffer from various issues, including sleep disturbances, with prevalence rates ranging between 20% and 80% (4). Insomnia significantly impacts individuals' quality of life, leading to substantial declines in daily functioning across emotional, social, and physical domains (5). Using

pharmaceutical drugs to improve sleep quality may pose additional risks for patients. Therefore, adopting alternative strategies is essential (6). Studies emphasize the effectiveness of massage and therapeutic touch in promoting relaxation and reducing emotional distress (7, 8). One such technique is slow stroke back massage (SSBM) (3). The SSBM promotes lymphatic drainage, increases blood circulation, stimulates veins, improves muscle metabolism, and reduces sympathetic activity, inducing patient relaxation (9).

A study by Lubbna et al. demonstrated that SSBM positively affects sleep quality and anxiety levels in elderly stroke patients (10). In contrast, Unal and Balci Akpinar concluded that the effectiveness of SSBM in patients remains controversial, requiring further studies (11). The time of back massage in various studies

is mostly mentioned as occurring during the night and before the patient sleeps. It is thought that there is a need for studies that evaluate the effect of massage earlier (12). On the other hand, some studies determined that there was no statistically significant relationship between back massage and sleep (11). Also, in the study by Harris and Richards, in which light pressure was applied and the session duration was the shortest at 3 minutes, it was found that back massage did not improve sleep quality (13). There is no common view on the duration of back massage, but it seems that the longer the duration of massage, the more effective it is.

These uncertainties within medical communities highlight the need for robust research backing to integrate these complementary therapies into nursing practices. Therefore, this study evaluates the effect of back massage on sleep quality.

## 2. Objectives

The SSBM is a simple, cost-effective, non-invasive, and safe technique without side effects (11). Therefore, this study aims to determine the impact of SSBM on sleep quality in hemodialysis patients, as well as sleep quality over time after the intervention.

## 3. Methods

### 3.1. Study Design and Setting

This study was a randomized controlled clinical trial (IRCT20231122060139N1) conducted at Shahid Madani Hospital in Khoy, Iran. The sample size, considering ( $\beta = 85\%$ ), ( $\alpha = 0.05$ ), (SD of the sleep quality = 4.24), the mean difference of sleep quality ( $d = 2.4$ ) (14), and with a consideration of 10% attrition (3), was calculated to be 60.

Patients were selected through convenience sampling and then assigned to two groups using simple random sampling. Using a simple randomization method, each of the selected patients was randomly assigned a number, and their name was written on envelopes given to someone unaware of the study objectives. All the sealed envelopes were placed in a container, and based on chance, the first 30 numbers taken out were assigned to the intervention group, and the second 30 numbers were assigned to the control group. In this way, neither the researcher nor the participant knew who was in which group.

Inclusion criteria were a Pittsburgh Sleep Quality Index (PSQI) score of 5 or higher, at least 3 months of dialysis history, and healthy skin in the massage area. Exclusion criteria were the use of complementary

therapies in the past three months, use of sleeping pills, severe neuropathy, hospitalization, and inability to attend massage sessions.

### 3.2. Intervention


After obtaining ethical approval and patient consent, an initial session was held to explain the study's objectives to participants. In the intervention group, a research assistant provided the SSBM thrice weekly for 10 minutes per session. Sleep quality was assessed in both groups using the PSQI before the intervention and again at two and four weeks post-intervention (Figure 1). Massages were conducted between 3 and 7 PM in a designated dialysis unit room free of other patients (15). Meanwhile, the control group received standard care and remained unaware of the intervention group's massage therapy.

To learn massage, the researcher was first taught SSBM by a physiotherapist, and after several practice sessions and repetitions, the researcher, under the supervision and approval of the physiotherapist, began performing the intervention. The stages of SSBM are as follows: The patient lies on a hard bed, and 20 small circular movements are made with the thumbs for 1 minute on the neck. Then the lumbar spine is massaged from the base of the skull to the sacrum with the palm of one hand, and this process is repeated on the other side of the spine with the palm of the other hand, while the first hand is moved towards the base of the skull. The number of movements was 60 times within 4 minutes. After that, 20 strokes are made with the thumb along the shoulder blades for 30 seconds. Strokes with the thumbs are made on both sides of the spine from the shoulder to the lower back for 1 minute. Then, sweeping strokes are made with both hands from the neck to near the sacrum with 40 movements within 3.5 minutes (15).

### 3.3. Data Collection and Analysis

The data collection tool was the PSQI, developed by Buysse et al. It includes 18 items with a 4-point Likert scale ranging from none = 0 to severe = 3, and seven subscales. The total score of the tool ranges from 0 to 21, with higher scores indicating poorer sleep quality. A score of 5 is the cut-off for distinguishing between good and poor sleep (16).

Data analysis was performed using SPSS software version 16, with P values less than 0.05 considered significant. The analysis utilized descriptive statistics, repeated measures tests, paired *t*-tests, and independent *t*-tests.



**Figure 1.** Consort diagram of research participants

#### 4. Results

Comparison of demographic characteristics before the intervention shows that there was no significant difference between the two groups (Table 1).

The sleep quality in both groups showed no significant difference before the intervention ( $P = 0.156$ ). The mean scores for all PSQI domains, except for daytime dysfunction, significantly decreased after the intervention compared to baseline and on the 15th and 30th days ( $P < 0.05$ ) (Table 2).

The results showed that the mean PSQI before the intervention in the intervention and control groups was  $11.82 \pm 6.36$  and  $11.89 \pm 6.32$ , respectively. On the 15th day after the intervention, the mean PSQI in the intervention and control groups was  $11.09 \pm 5.95$  and  $13.1 \pm 6.15$ , respectively. On the 30th day, the mean PSQI in the intervention and control groups was  $10.19 \pm 5.39$  and  $12.55 \pm 6.2$ , respectively. The repeated measures test showed that SSBM significantly improved the sleep quality of patients ( $P < 0.001$ ) compared to before the intervention, and on the 15th and 30th days in the

**Table 1.** Comparison of Demographic Characteristics Between the Two Groups Before the Intervention <sup>a</sup>

| Demographic Variables        | Control Group | Intervention Group | P-Value |
|------------------------------|---------------|--------------------|---------|
| <b>Sex</b>                   |               |                    | 0.59    |
| Female                       | 15 (50)       | 14 (46.6)          |         |
| Male                         | 15 (50)       | 16 (53.3)          |         |
| <b>Education</b>             |               |                    | 0.33    |
| Illiterate                   | 6 (20)        | 5 (16.6)           |         |
| Elementary                   | 5 (16.6)      | 7 (23.3)           |         |
| Diploma                      | 17 (56.6)     | 15 (50)            |         |
| Bachelor and above           | 2 (6.6)       | 3 (10)             |         |
| <b>Marital</b>               |               |                    | 0.38    |
| Single                       | 4 (13.3)      | 5 (16.6)           |         |
| Married                      | 19 (63.3)     | 20 (66.6)          |         |
| Widow                        | 7 (23.3)      | 5 (16.6)           |         |
| <b>Income</b>                |               |                    | 0.46    |
| Sufficiency                  | 11 (36.6)     | 13 (43.3)          |         |
| Insufficiency                | 19 (63.6)     | 17 (56.6)          |         |
| <b>Employment</b>            |               |                    | 0.79    |
| Employed                     | 14 (13.3)     | 13 (43.3)          |         |
| Unemployed                   | 16 (53.3)     | 17 (56.6)          |         |
| <b>Hemodialysis sessions</b> |               |                    | 0.55    |
| Twice a week                 | 18 (60)       | 16 (53.3)          |         |
| Three times a week           | 12 (40)       | 14 (46.6)          |         |
| <b>Hours of hemodialysis</b> |               |                    | 0.32    |
| Three hours                  | 21 (70)       | 19 (63.3)          |         |
| More than three hours        | 9 (30)        | 11 (36.6)          |         |
| <b>Age</b>                   | 48.14 ± 12.33 | 47.88 ± 12.33      | 0.89    |

<sup>a</sup> Values are expressed as No. (%) or mean ± SD.

intervention group. Although the sleep quality of the control group worsened on the 15th day, this change was not statistically significant ( $P > 0.05$ ) and may have been due to other underlying factors such as stress that were beyond the researcher's control (Table 3).

The independent *t*-test revealed that the sleep quality of patients in the intervention group was significantly better than that in the control group at the end of the 15th and 30th days ( $P < 0.05$ ) (Figure 2). Sleep quality in the SSBM group significantly improved ( $P < 0.001$ ) (Table 4).

## 5. Discussion

The findings indicated that SSBM improved sleep quality in all domains, except for daytime dysfunction. A study by Mirawati revealed a significant difference in four components of sleep quality: Subjective sleep quality, sleep latency, sleep efficiency, and daytime dysfunction, as well as the overall sleep quality score in elderly individuals after massage. However, no

differences were observed in three domains: Use of sleep medications, sleep disturbances, and sleep duration (17). The discrepancy between results could be attributed to the fact that Mirawati focused on an elderly population that generally experiences more sleep problems, and perhaps a more extended intervention would be needed.

A study by Hojat Ansari et al. showed improvement in sleep quality in SSBM or hot stone massage groups. However, no difference was found between the two intervention groups with increasing massage durations (18). Unlike this study, the current research did not use two intervention groups. However, it only assessed the effect of massage on sleep quality in one group at two time points, with significant results. The results confirmed that SSBM improved sleep quality in the intervention group compared to before the intervention and on the 15th and 30th days.

Studies by Sihaloho et al. and Oshvandi et al. indicated that the time to fall asleep or return to sleep after waking up significantly decreased following

**Table 2.** Comparison of the Mean Scores of Components of Pittsburgh Sleep Quality Index Before and After the Intervention <sup>a</sup>

| Sleep Quality Domain                | Control Group | Day 15th    | Day 30th    |
|-------------------------------------|---------------|-------------|-------------|
| <b>Subjective sleep quality</b>     |               |             |             |
| Before intervention                 | 1.51 ± 0.85   | 1.48 ± 1.06 | 1.45 ± 0.85 |
| After intervention                  | 1.38 ± 0.84   | 1.36 ± 0.74 | 1.35 ± 0.98 |
| Paired <i>t</i> -test               | 0.1           | 0.004       | 0.001       |
| Cohen's d                           | 0.23          | 0.16        | 0.11        |
| <b>Sleep latency</b>                |               |             |             |
| Before intervention                 | 2.29 ± 0.86   | 2.03 ± 0.91 | 2.25 ± 0.93 |
| After intervention                  | 2.23 ± 0.88   | 1.68 ± 0.9  | 1.78 ± 0.81 |
| Paired <i>t</i> -test               | 0.3           | 0.03        | 0.001       |
| Cohen's d                           | 0.06          | 0.38        | 0.58        |
| <b>Sleep duration</b>               |               |             |             |
| Before intervention                 | 1.71 ± 0.99   | 2.16 ± 0.86 | 1.93 ± 0.96 |
| After intervention                  | 1.67 ± 1.04   | 1.75 ± 0.88 | 1.51 ± 0.84 |
| Paired <i>t</i> -test               | 0.3           | 0.002       | 0.001       |
| Cohen's d                           | 0.04          | 0.47        | 0.5         |
| <b>Sleep efficiency</b>             |               |             |             |
| Before intervention                 | 2 ± 1.06      | 2.35 ± 0.98 | 2 ± 0.1     |
| After intervention                  | 2.06 ± 1.06   | 1.92 ± 1.05 | 1.74 ± 1.05 |
| Paired <i>t</i> -test               | 0.3           | 0.02        | 0.03        |
| Cohen's d                           | 0.05          | 0.43        | 0.26        |
| <b>Sleep disturbances</b>           |               |             |             |
| Before intervention                 | 1.53 ± 0.61   | 1.54 ± 0.67 | 1.67 ± 0.54 |
| After intervention                  | 1.58 ± 0.67   | 1.42 ± 0.67 | 1.51 ± 0.58 |
| Paired <i>t</i> -test               | 0.9           | 0.04        | 0.04        |
| Cohen's d                           | 0.08          | 0.17        | 0.29        |
| <b>Sedative medication use</b>      |               |             |             |
| Before intervention                 | 1.48 ± 1.18   | 2 ± 0.1     | 1.67 ± 1.16 |
| After intervention                  | 1.45 ± 1.15   | 1.57 ± 1.03 | 1.48 ± 0.97 |
| Paired <i>t</i> -test               | 0.3           | 0.001       | 0.003       |
| Cohen's d                           | 0.02          | 0.43        | 0.19        |
| <b>Daily functioning impairment</b> |               |             |             |
| Before intervention                 | 1.32 ± 0.7    | 1.54 ± 0.67 | 1.58 ± 0.76 |
| After intervention                  | 1.45 ± 0.72   | 1.39 ± 0.68 | 1.33 ± 0.62 |
| Paired <i>t</i> -test               | 0.1           | 0.2         | 0.04        |
| Cohen's d                           | 0.18          | 0.22        | 0.40        |


<sup>a</sup> Values are expressed as mean ± SD.**Table 3.** Mean and Difference in Sleep Quality Scores of Hemodialysis Patients in Intervention and Control Groups

| Variables           | Intervention Group         |                   |                        | Control Group              |                   |                        |
|---------------------|----------------------------|-------------------|------------------------|----------------------------|-------------------|------------------------|
|                     | Mean ± SD                  | Mean ± SD Changes | Repeated Measures Test | Mean ± SD                  | Mean ± SD Changes | Repeated Measures Test |
| Time point          | F = 7.1, P > 0.001, Df = 2 |                   |                        | F = 4.2, P > 0.131, Df = 2 |                   |                        |
| Before intervention | 11.82 ± 6.36               | -                 |                        | 11.89 ± 6.32               | -                 |                        |
| Day 15th            | 11.09 ± 5.95               | 0.73 ± 4.2        |                        | 13.1 ± 6.15                | 1.21 ± 5.9        |                        |
| Day 30th            | 10.19 ± 5.39               | 1.63 ± 3.9        |                        | 12.55 ± 6.2                | 0.66 ± 5.4        |                        |

massage (19, 20), which aligns with the results of the current study.

Studies by Naderi et al., Cecen and Lafci, and Doner and Tasci showed improved sleep quality, depression,

stress, anxiety, pain, fatigue, and quality of life between two groups during all follow-up weeks after the intervention (21-23). Contrary to the results of the present study, Williamson et al. showed there was no



**Figure 2.** Comparison of the results in the control group and the intervention group (compared with the control group,  $P < 0.05$ ).

**Table 4.** Comparison of Mean Sleep Quality Scores of Hemodialysis Patients Between the Intervention and Control Groups

| Variables           | Mean Differences | SE    | Degrees of Freedom | t     | P-Value |
|---------------------|------------------|-------|--------------------|-------|---------|
| Before intervention | 0.03             | 0.421 | 58                 | 0.523 | 0.156   |
| Day 15th            | 4.99             | 0.365 | 58                 | 0.385 | 0.023   |
| Day 30th            | 3.64             | 0.657 | 58                 | 0.374 | 0.026   |

difference between sleep adequacy scores in the massage therapy and control groups (24). The lack of consistency may be attributed to the study population, as the research by Williamson focused on postmenopausal women, and the physiological and psychological changes during this period can affect sleep in women. Additionally, massage in cancer patients did not show a significant decrease in fatigue, pain, and discomfort (25). One of the main reasons for this could be that fatigue, pain, and discomfort in cancer patients are often chronic, so massage interventions may need to be applied over a more extended period.

Considering the effect of massage on patients' sleep quality, it is suggested that future studies compare the effect of massage with other complementary medicine methods on patients' sleep quality. Also, the effect of the amount of pressure applied during massage, the duration of massage, and different times of massage on patients' sleep should be investigated.

In terms of research limitations, given that sleep quality is a subjective phenomenon, many factors such as individual factors, family problems, and cultural and social differences affect it, which could not be controlled by the researchers. Additionally, the need for patients to go to the dialysis department in the afternoon can also be one of the possible factors affecting the results, as



most of the patients had dialysis in the morning shift. One of the strengths of this study is that it effectively improved sleep quality in patients, such as increasing sleep duration, falling asleep faster, and sleeping better without any side effects.

### 5.1. Conclusions

Based on the results, implementing massage therapy programs can improve patients' psychological symptoms and sleep quality. This is because massage therapy can be applied in various settings without specific equipment or the side effects associated with medications. The findings of this study can be utilized by health policymakers, nurses, and healthcare providers in hemodialysis units and other medical groups to incorporate non-pharmacological methods into the treatment and care plans for patients.

### Acknowledgements

This article is the result of a project by the Student Research Committee of Khoy University of Medical Sciences. The authors sincerely thank the administrators, staff, and patients of the Shahid Modani Hospital, Khoy, hemodialysis unit for their collaboration.

### Footnotes

**Authors' Contribution:** M. A.: Conceptualization, formal analysis, data curation, and supervision; F. M.: Methodology and data collection; M. Z.: Project administration and resources.

**Clinical Trial Registration Code:** [IRCT20231122060139N1](https://www.clinicaltrials.gov/ct2/show/study?term=IRCT20231122060139N1).

**Conflict of Interests Statement:** The authors declare no conflict of interest.

**Data Availability:** The dataset presented in the study is available on request from the corresponding author during submission or after publication.

**Ethical Approval:** This study was approved by Khoy University of Medical Sciences, Iran and all data collection and analysis procedures were conducted in accordance with the tenets of the Declaration of Helsinki ([IR.KHOY.REC.1402.039](https://www.khoymedicalscience.ir/KHOY.REC.1402.039)).

**Funding/Support:** This research received no funding/support.

**Informed Consent:** All the subjects voluntarily agree to participate in this research. Written informed consents were provided by all the subjects participated in the study.

### References

- Murdeswar HN, Anjum F. *Hemodialysis*. Treasure Island (FL): StatPearls; 2025. [PubMed ID: [33085443](#)].
- Canaud B, Chazot C, Koomans J, Collins A. Fluid and hemodynamic management in hemodialysis patients: challenges and opportunities. *J Bras Nefrol*. 2019;**41**(4):550-9. [PubMed ID: [31661543](#)]. [PubMed Central ID: [PMC6979572](#)]. <https://doi.org/10.1590/2175-8239-JBN-2019-0135>.
- Raiesifar Z, Molavynejad S, Raiesifar A, Maraghi E, Vaismoradi M, Helalat A. [The effect of slow-stroke back massage on fatigue and sleep quality in hemodialysis patients: A clinical trial study]. *J Hayat*. 2021;**27**(1):59-71. FA.
- Malekshahi F, Aryamanesh F, Fallahi S. The Effects of Massage Therapy on Sleep Quality of Patients with End-Stage Renal Disease Undergoing Hemodialysis. *Sleep Hypnosis - Int J*. 2017;**20**(2):91-5. <https://doi.org/10.5350/Sleep.Hypn.2017.19.0138>.
- Lyons OD. Sleep disorders in chronic kidney disease. *Nat Rev Nephrol*. 2024;**20**(10):690-700. [PubMed ID: [38789686](#)]. <https://doi.org/10.1038/s41581-024-00848-8>.
- Baykal D, Comlekci N. Non-Pharmacologic Approaches to Sleep Problems for Palliative Care Cancer Patients: A Systematic Review. *Florence Nightingale J Nurs*. 2023;**31**(2):131-7. [PubMed ID: [37404216](#)]. [PubMed Central ID: [PMC10440962](#)]. <https://doi.org/10.5152/FNjN.2023.23051>.
- Ghasemi M, Rejeh N, Bahrami T, Heravi-Karimooi M, Tadrissi SD, Vaismoradi M. Aromatherapy Massage vs. Foot Reflexology on the Severity of Restless Legs Syndrome in Female Patients Undergoing Hemodialysis. *Geriatrics (Basel)*. 2021;**6**(4). [PubMed ID: [34698176](#)]. [PubMed Central ID: [PMC8544351](#)]. <https://doi.org/10.3390/geriatrics6040099>.
- Suarilah I, Muhammad H, Retno I. The Effect of Cutaneous Stimulation Slow Stroke Back Massage on Sleep Quality Improvement in Elderly. *Int J Psychosocial Rehab*. 2020;**11**(3):9428-32.
- Patonengan GS, Mendrofa FAM, Hani U. Effectiveness of Slow Stroke Back Massage (SBBM) on blood pressure, anxiety, and depression among older people with hypertension: A quasi-experimental study. *Public Health Indonesia*. 2023;**9**(2):66-73. <https://doi.org/10.36685/phi.v9i2.678>.
- Lubbnas S, Dani AH, Asih SW. The effects of slow-stroke back massage on anxiety and sleep problems in elderly stroke patients. *Malahayati Int J Nurs Health Sci*. 2023;**6**(1):53-9. <https://doi.org/10.33024/minh.v6i1.8874>.
- Unal KS, Balci Akpınar R. The effect of foot reflexology and back massage on hemodialysis patients' fatigue and sleep quality. *Complement Ther Clin Pract*. 2016;**24**:139-44. [PubMed ID: [27502815](#)]. <https://doi.org/10.1016/j.ctcp.2016.06.004>.
- Akpınar RB, Akın G, Ay E. The Effect of Back Massage on Sleep Quality: A Systematic Review. *Kafkas J Med Sci*. 2022;**12**(2):179-84. <https://doi.org/10.5505/kjms.2022.25932>.
- Harris M, Richards KC. The physiological and psychological effects of slow-stroke back massage and hand massage on relaxation in older people. *J Clin Nurs*. 2010;**19**(7-8):917-26. [PubMed ID: [20492036](#)]. <https://doi.org/10.1111/j.1365-2702.2009.03165.x>.
- Mohammadpourhodki R, Sargolzaei MS, Basirinezhad MH. Evaluating the effect of massage based on slow stroke back massage

- on the anxiety of candidates for cataract surgery. *Rom J Ophthalmol*. 2019;**63**(2):146-52. [PubMed ID: 31334393]. [PubMed Central ID: PMC6626929].
15. Nahamin M, Akbarbegloo M, Habibipur Z. The Impact of Slow-Stroke Back Massage on Anxiety Among Patients Receiving Hemodialysis: A Randomized Clinical Trial. *Mod Care J*. 2016;**In Press**(In Press). <https://doi.org/10.17795/modernc.10273>.
  16. Buysse DJ, Reynolds CF, Monk TH, Berman SR, Kupfer DJ. The Pittsburgh Sleep Quality Index: a new instrument for psychiatric practice and research. *Psychiatry Res*. 1989;**28**(2):193-213. [PubMed ID: 2748771]. [https://doi.org/10.1016/0165-1781\(89\)90047-4](https://doi.org/10.1016/0165-1781(89)90047-4).
  17. Mirawati D. The Effect of Back Massage on Decreasing Insomnia in the Elderly at the Posyandu Lansia Marsudi Waras Jebres Surakarta. *Gaster*. 2021;**19**(1). <https://doi.org/10.30787/gaster.v19i1.581>.
  18. Hojat Ansari M, Aemmi SZ, Farhadi Faruji A, Esmaily H. The therapeutic effect of two different massages on the sleep quality of orthopedic surgery patients: A randomized controlled study. *Explore (NY)*. 2025;**21**(1):103084. [PubMed ID: 39586193]. <https://doi.org/10.1016/j.explore.2024.103084>.
  19. Sihaloho E, Dalimunthe SY, Simamora M, Ristiani R. Effectiveness of Slow Stroke Back Massage Therapy on the Sleep Quality of Postpartum Mother. *Proceedings Int Conference Nurs Health Sci*. 2024;**3**(1):299-306.
  20. Oshvandi K, Mirzajani Letomi F, Soltanian AR, Shamsizadeh M. The effects of foot massage on hemodialysis patients' sleep quality and restless leg syndrome: a comparison of lavender and sweet orange essential oil topical application. *J Complement Integr Med*. 2021;**18**(4):843-50. [PubMed ID: 33838094]. <https://doi.org/10.1515/jcim-2020-0121>.
  21. Naderi A, Rezvani MH, Aminian-Far A, Hamood-Ahvazi S. Can a six-week Swedish massage reduce mood disorders and enhance the quality of life in individuals with Multiple Sclerosis? A randomized control clinical trial. *Explore (NY)*. 2024;**20**(5):103032. [PubMed ID: 39018656]. <https://doi.org/10.1016/j.explore.2024.103032>.
  22. Cecen S, Lafci D. The effect of hand and foot massage on fatigue in hemodialysis patients: A randomized controlled trial. *Complement Ther Clin Pract*. 2021;**43**:101344. [PubMed ID: 33714863]. <https://doi.org/10.1016/j.ctcp.2021.101344>.
  23. Doner A, Tasci S. Effect of massage therapy with lavender oil on severity of restless legs syndrome and quality of life in hemodialysis patients. *J Nurs Scholarsh*. 2022;**54**(3):304-14. [PubMed ID: 34779137]. <https://doi.org/10.1111/jnu.12738>.
  24. Williamson J, White A, Hart A, Ernst E. Randomised controlled trial of reflexology for menopausal symptoms. *BJOG*. 2002;**109**(9):1050-5. [PubMed ID: 12269681]. <https://doi.org/10.1111/j.1471-0528.2002.01504.x>.
  25. Cassileth BR, Vickers AJ. Massage therapy for symptom control: outcome study at a major cancer center. *J Pain Symptom Manage*. 2004;**28**(3):244-9. [PubMed ID: 15336336]. <https://doi.org/10.1016/j.jpainsymman.2003.12.016>.