



A survey on the Frequency of *Sarcocystis* in Bandar Abbas, Iran in 2019 - 2020

Masoud Sheikhi¹, Abdoreza Salahi-Moghaddam¹, Majid Najafi Asl², Abbas Farahani¹ and Jebreil Shamseddin^{1,*}

¹Infectious and Tropical Diseases Research Center, Hormozgan Health Institute, Hormozgan University of Medical Sciences, Bandar Abbas, Iran

²Department of Laboratory Sciences, Faculty of Para-Medicine, Hormozgan University of Medical Sciences, Bandar Abbas, Iran

*Corresponding author: Infectious and Tropical Diseases Research Center, Hormozgan Health Institute, Hormozgan University of Medical Sciences, Bandar Abbas, Iran. Email: shams.jebreil@gmail.com

Received 2020 October 03; Revised 2020 October 12; Accepted 2020 October 19.

Abstract

Background: The zoonotic *Sarcocystis* parasite has an obligatory two-host life cycle that mainly involves herbivorous animals as intermediate hosts and carnivorous animals as definitive hosts.

Objectives: Lack of reliable study and published data about frequency of *Sarcocystis* in livestock of Hormozgan Province and consumed meat led us to investigate in abattoirs and slaughterhouses of Bandar Abbas, Iran, in 2019 - 2020.

Methods: In this descriptive cross-sectional study, 400 meat samples of three types of animals (cow, sheep, goat) belonging to Hormozgan, Fars, and Kerman provinces were studied from September 2019 to January 2020 using naked eye examination for detection of macroscopic *Sarcocystis* cysts and pepsin digestion method accompanied by squeezing methods performed to examine striated muscles for microscopic cyst types. Isolated tissues of the esophagus, heart, and diaphragm of 400 slaughtered animals were examined for *Sarcocystis*.

Results: The carcasses of all the animals were investigated to detect *Sarcocystis* macroscopic cysts, all of which were negative. However, microscopic examination of isolated tissues by pepsin digestion showed a total frequency of 92.25% in these animals. Analysis revealed that cows bred in Hormozgan and goats imported from Kerman are significantly infected and play an important role in the distribution of disease ($P < 0.05$).

Conclusions: These results and obtained data indicated that a large volume of imported meats to Hormozgan is contaminated with this parasite, and more control should be applied over slaughtered livestock. Also, the people of the community who consume these meats should be given proper and complete training.

Keywords: Slaughtered Livestock, *Sarcocystis*, Pepsin-Digestion Method

1. Background

Sarcocystis parasites are coccidian protozoan organisms with worldwide distribution and one of the most common parasites in domestic livestock that can cause severe infection in some hosts such as cattle and sheep (1, 2). Also, some species of the parasite transmit to humans and cause clinical presentations. *Sarcocystis* parasites have significant health importance, especially in certain parts of the world that animal breeding is a main occupation, thus impose economic losses. Reduced milk production, abortion, and sometimes death occurs in intermediate hosts such as cows, sheep, goats, and pigs (2). *Sarcocystis* parasites need two hosts to complete their life cycle. In its life cycle, there is a definitive host and an intermediate host. The sexual life cycle occurs in the definitive host that often

belongs to carnivorous animals that accidentally ingest oocysts. The asexual part of the life cycle takes place in the intermediate host (mostly herbivores) (3). The dogs and cats play an important role in the spreading of the parasite in the environment. They can excrete up to 200 million infective oocysts in the course of disease. Animals remain asymptomatic and rarely have mild diarrhea. Humans are infected by eating schizonts (asexual form) of *Sarcocystis hominis* and *S. suihominis* in undercooked beef, pork, mutton, and raw or poorly cooked meat products (hamburgers, sausages, and hot dogs) (1, 3). People who voluntarily ate *Sarcocystis*-infected meat excreted *Sarcocystis* oocysts in feces after eight days or more. The severity of clinical signs depends on the type of meat consumed (beef or pork) and *Sarcocystis* species causative agent (4).

The *Sarcocystis* can act as an opportunistic organism in

immunocompromised patients, which can cause intestinal or muscular disturbances, but epidemiology and clinical impact of human infections by *Sarcocystis* are largely unknown, and reports continue to be rare (5). The prevalence of sarcocystosis in slaughtered animals has been studied in many regions around the world. The digestion technique of infected tissues is the most common and sensitive method for detecting infections with microscopic forms of *Sarcocystis*, which comprises taking pieces of the contaminated tissue and crushing it, then solubilizing it in a digestive solution containing chloric acid and pepsin, and subsequently staining with Giemsa. Contamination rates from 3% to 100% have been reported by different methods in different parts of the world (6, 7). The method of identification of organisms affects the rate of positivity, e.g., microscopic or molecular methods. Its frequency in livestock such as cattle, sheep, and goats has been reported up to 100% in some parts of Iran (8). Some carnivores are the definitive host for many species of *Sarcocystis*, but species transmitted by dogs are usually not transmitted by cats (9). Systematic analysis revealed that the total prevalence (95% confidence intervals) of *Sarcocystis* spp. in Iranian ruminants, according to published studies, was an average of 74.40% (10).

2. Objectives

To the best of our knowledge, there is no reliable study and published data about *Sarcocystis* in Hormozgan abattoirs (Industrial and traditional), which investigate consumed meat contamination. Thus, we designed a descriptive cross-sectional study to observe the situation. A large amount of consumed meats in Hormozgan is imported from nearby provinces, and determination of *Sarcocystis* frequency in Hormozgan can help researchers to estimate Sarcocystosis in those regions and get more ideas for further investigations.

3. Methods

The study was done from September 2019 to January 2020 in Hormozgan. The population of the study was 400 samples of slaughtered animals belonging to Hormozgan, Fars, and Kerman provinces.

After taking and processing tissue specimens, 20 g of the tissues, i.e., heart, diaphragm, and esophagus, were crushed with an electric shredder and placed in 2 mL of digestion solution (Figure 1). To make a digestion buffer, it is necessary to solve 1.3 g of pepsin, 3.5 mL HCl and 2.5 g NaCl in 500 mL of distilled water at pH = 7.4. Meat samples were placed in the incubator at 40°C for 3 hours to release

parasites and passed through a non-sterile gauze. The filtrated liquids were centrifuged at 2500 rpm for 5 min, and a direct smear was prepared from the precipitate. It is important to allow those smears to air-dry at least 1-hour before staining. Smears were fixed with methanol 96% (Germany, Merck) then stained with diluted Giemsa according to instructions for 15 minutes; after staining, slides were washed with tap water and allowed to air-dry completely. The stained slides were examined microscopically at magnifications of 400× and 1000×.



Figure 1. Taking of Specimens from 3 targeted tissues, heart, esophagus and liver in Bandar Abbas abattoirs

A minimum of 100 high-power fields must be examined. It should be noted that in cases that smears are strongly positive and especially when there was high contamination, plenty of parasites were readily detectable in magnification of 400×.

3.1. Statistical Analysis

The obtained data were analyzed via SPSS software ver. 22 (Chicago, IL, USA). Moreover, the chi-square test was used for the analysis of data. To control the confounder variables, the logistic regression analysis was implemented. The results would be considered statistically significant if the P value was less than 0.05 ($P < 0.05$).

3.2. Ethical Clearance

This descriptive cross-sectional study was approved by the Ethics Committee of Hormozgan University of Medical Sciences (HUMS) numbered IR.HUMS.REC.1397.143.

4. Results

In this study, a total of 400 samples were collected from slaughtered animals in two industrial and traditional slaughterhouses in Bandar Abbas, Iran, and transferred to the parasitology department laboratory. Slaughtered animals and carcasses were examined for macroscopic cysts, and all were negative. All samples were evaluated by the digestion method and direct smear microscopy. The slaughtered animals were 212 cows (53.0%), 152 goats (38.0%) and 36 sheep (9.0%). According to the information obtained, it was found that a large number of slaughtered cattle were imported to Bandar Abbas from other provinces.

The total frequency was 92.5%. Most of the imported cattle for meat consumption were from Kerman and Fars provinces. Therefore, at the time of sampling, important information, and animal characteristics such as sex, type of livestock, and breeding location were considered and recorded for each animal. Most of the samples belonged to animals bred in Hormozgan Province with 188 (47.0%), Kerman province with 38.3% (153), Fars 13.3% (49), and other provinces 2.5% (10). Hormozgan Province had the highest number of cows with 157 (74.1%), and Kerman province had the highest number of goats and sheep with 84 (55.26%) and 18 (50%), respectively (Table 1). The highest frequency of *Sarcocystis* in all three kinds of animals was detected in esophageal tissue, with 87.5% (350). Heart tissue with 72.8% (291) had the lowest frequency rate (Table 2).

Statistics showed that total frequency had no significant relation with animal gender ($P > 0.05$). The breeding location was determined and considered during sampling. According to the gender variable, infection in male cows of Hormozgan Province with 163 (47.7%) is significantly higher than animals imported from other provinces. The total frequency in each province based on all three infected tissues includes Kerman 94.8%, Hormozgan 93.6%, Fars 83.7%, and 70% in other provinces, respectively. According to the results of the above-mentioned infection rates, Kerman Province had the highest level of contamination rate (94.8%). The results show that among all three provinces, the cows of Hormozgan with 151 (96.2%) had a significant frequency ($P < 0.005$). Using the chi-square test and comparing the total infections and by animal type of each province, goats in Kerman Province with 60.9% were significantly infected compared to goats of other provinces.

5. Discussion

In this study, we investigated the frequency of *Sarcocystis* parasites in domestic animals slaughtered in Bandar

Abbas slaughterhouses. A total of 400 animals and 3 different tissues were studied. These animals consisted of 212 cows (53%), 152 goats (38%), and 36 sheep (9%), respectively. The carcasses of all the animals were examined for the presence of *Sarcocystis* macroscopic cysts, all of which were negative. However, microscopic examination by pepsin digestion showed a total frequency of 92.25% in these animals. Rate of infection indicates the high frequency of parasites in slaughtered animals that supply edible meat in Bandar Abbas. In other parts of the country, various studies have been conducted that show the high frequency of *Sarcocystis* in livestock in Iran, and our results are similar to many of these studies. Investigations in other provinces in Iran showed the reported frequency between 90% - 100% (6). In other parts of the world, *Sarcocystis* has been detected in 70 - 100 of examined meat samples (11).

In two studies carried out in Shahriar (Alborz Province) with a time interval, infection with macroscopic cysts was not reported (12). Also, studies in Isfahan and Kerman did not report any contamination with macroscopic cysts, or macroscopic cysts were restricted to a specific organ like the esophagus that is in agreement with our study (6, 13, 14). However, in some studies, macroscopic cysts have been reported at a low frequency (15, 16).

The results of this study showed that the total frequency of *Sarcocystis* has a significant relation with infected organ and livestock (animal) type ($P < 0.005$). The highest esophageal infection was reported in cows with 92.2%, the highest diaphragm infection in sheep with 91.7%, and the highest heart infection occurred in cows with 84.9%. In 2013, Alibeigi et al. (12), by direct microscopic method, detected the same results (12), but Mirzaei and Rezaei (17) identified the heart as the most infected organ in 2008. Some other studies indicated that all three examined organs were 100% infected.

Due to the geographical location and climatic conditions of Hormozgan Province, animal husbandry is not very prosperous in this province. Therefore, in order to supply edible meat, it is necessary to import livestock from other provinces of the country, especially neighboring provinces. The obtained information indicates that more than half of the animals slaughtered in Bandar Abbas slaughterhouses come from other provinces, e.g., Fars and Kerman. By reviewing the statistical results, it was found that most of the animals slaughtered in this study were related to Hormozgan Province with 47%, Kerman Province 38.3%, Fars Province 13.3%, and other provinces 2.5%, respectively.

Total *Sarcocystis* frequency had no significant statistical relation with animal gender. However, the frequency of *Sarcocystis* infection in male animals in Hormozgan Province was significantly higher than the livestock of

Table 1. The Number of Livestock Imported from Other Places (Provinces) to Hormozgan Abattoirs^a

Province	Total	Others	Fars	Kerman	Hormozgan
Kind of animals					
Cow	212 (53.0)	4 (40.0)	0 (0.0)	51 (33.3)	157 (83.5)
Goat, No	152 (38.0)	4 (40.0)	42 (85.7)	84 (54.9)	22 (11.7)
Sheep	36 (9.0)	2 (20.0)	7 (14.3)	18 (11.8)	9 (4.8)
The number of cattle according to province	400	10 (2.5)	49 (13.3)	153 (38.3)	188 (47.0)

^aValues are expressed as No. (%).

Table 2. Total Contamination Rate in Three Types of Cattle Slaughtered According to Infected Organs

Infected Tissue	Kinds of Cattle ^a			Total ^a	P Value
	Sheep	Goat	Cow		
Heart					< 0.001 ^b
Negative	13 (36.1)	64 (42.1)	32 (15.1)	109 (27.3)	
Positive	23 (63.9)	88 (57.9)	180 (84.9)	291 (72.8)	
Diaphragm					0.009 ^b
Negative	3 (8.3)	22 (14.5)	53 (25.0)	78 (19.5)	
Positive	33 (91.7)	130 (85.5)	159 (75.0)	322 (80.5)	
Esophagus					0.004 ^b
Negative	5 (13.9)	29 (19.1)	16 (7.5)	50 (12.5)	
Positive	31 (86.1)	123 (80.9)	196 (92.5)	350 (87.5)	
Total					0.021 ^b
Negative	2 (5.6)	19 (12.5)	10 (4.7)	31 (7.75)	
Positive	34 (94.4)	133 (87.5)	202 (95.3)	369 (92.25)	

^aValues are expressed as No. (%).

^bSignificant.

other provinces based on the breeding location ($P < 0.5$). In the studies conducted in Kerman, Jahrom, and Tabriz, the frequency of infection was not significantly related to gender (2). In this study, the highest total infection based on breeding location is related to Kerman Province livestock with 94.8%, which is significantly higher than the frequency in the livestock of other provinces. These studies are consistent with our results. The key point of this study is that the current data of this study can be an estimate for the neighboring provinces because of a large number of slaughtered animals imported from neighbors. On the other hand, the lack of information about this parasite in Hormozgan was filled because there was no reliable and published data before.

5.1. Conclusions

The foodborne and zoonotic *Sarcocystis* parasite gives an idea to evaluate and examine fecal samples of people in future studies. Diagnostic limitations have impeded our

ability to increase our understanding of the epidemiology and public health significance of *Sarcocystis* infection in humans. This study seems to be the first of its own kind carried out in meat samples of Hormozgan. This parasite also causes health, economic and pathological problems in humans and animals. Therefore, it is important to note that macroscopic methods implemented in abattoirs and slaughterhouses are not a good way to detect this parasite in livestock, and microscopic digestion methods should be used to obtain the correct results. The most important conclusion of this study is that more control should be applied over slaughtered livestock and that the people of the community who consume these meats should be given proper and complete training, such as full cooking or freezing of meat before consuming it.

Footnotes

Authors' Contribution: MS and JS conceived the project and designed the experiments. MS, MN, JS, and AF designed and collected samples. MS, JS, MN, AS, and AF analyzed the data. JS and MS supervised the collection of the samples. MS, JS, MN, AS, and AF wrote the manuscript. All authors reviewed and approved the manuscript.

Conflict of Interests: There are no conflicts of interest for authors.

Ethical Approval: This study was approved by the Ethics Committee of Hormozgan University of Medical Sciences (HUMS) numbered IR.HUMS.REC.1397.143.

Funding/Support: Hormozgan University of Medical Sciences funded this research.

References

- Hajimohammadi B, Zohourtabar A, Fattahi BA, Omidpanah A, Rayati T, Ghasemi A, et al. Occurrence and distribution of Sarcocystis parasite isolated from sheep in Yazd Province, Iran. *J Community Health Res.* 2014;**3**(3):204-9.
- Mirzaei M, Rezaei H. A survey on Sarcocystis spp. infection in cattle of Tabriz city, Iran. *J Parasit Dis.* 2016;**40**(3):648-51. doi: [10.1007/s12639-014-0551-2](https://doi.org/10.1007/s12639-014-0551-2). [PubMed: [27605760](https://pubmed.ncbi.nlm.nih.gov/27605760/)]. [PubMed Central: [PMC4996164](https://pubmed.ncbi.nlm.nih.gov/PMC4996164/)].
- Fayer R, Esposito DH, Dubey JP. Human infections with Sarcocystis species. *Clin Microbiol Rev.* 2015;**28**(2):295-311. doi: [10.1128/CMR.00113-14](https://doi.org/10.1128/CMR.00113-14). [PubMed: [25715644](https://pubmed.ncbi.nlm.nih.gov/25715644/)]. [PubMed Central: [PMC4402950](https://pubmed.ncbi.nlm.nih.gov/PMC4402950/)].
- Nesbakken T. *Biological pathogens in animals*. Woodhead Publishing; 2005. doi: [10.1533/9781845691028.1.3](https://doi.org/10.1533/9781845691028.1.3).
- Reiner G, Eckert J, Peischl T, Bochert S, Jäkel T, Mackenstedt U, et al. Variation in clinical and parasitological traits in Pietrain and Meishan pigs infected with Sarcocystis miescheriana. *Veterinary Parasitology.* 2002;**106**(2):99-113. doi: [10.1016/s0304-4017\(02\)00041-9](https://doi.org/10.1016/s0304-4017(02)00041-9).
- Nourollahi Fard SR, Asghari M, Nouri F. Survey of Sarcocystis infection in slaughtered cattle in Kerman, Iran. *Trop Anim Health Prod.* 2009;**41**(8):1633-6. doi: [10.1007/s11250-009-9358-z](https://doi.org/10.1007/s11250-009-9358-z). [PubMed: [19390981](https://pubmed.ncbi.nlm.nih.gov/19390981/)].
- Imre K, Darabus G, Tirziu E, Morariu S, Imre M, Plutzer J, et al. Sarcocystis spp. In romanian slaughtered cattle: Molecular characterization and epidemiological significance of the findings. *Biomed Res Int.* 2019;**2019**:4123154. doi: [10.1155/2019/4123154](https://doi.org/10.1155/2019/4123154). [PubMed: [31737662](https://pubmed.ncbi.nlm.nih.gov/31737662/)]. [PubMed Central: [PMC6815619](https://pubmed.ncbi.nlm.nih.gov/PMC6815619/) publication of this paper].
- Mirzaei Dehaghi M, Fallahi M, Sami M, Radfar MH. Survey of sarcocystis infection in slaughtered sheep in Kerman Abattoir, Kerman, Iran. *Comp Clin Path.* 2012;**22**(3):343-6. doi: [10.1007/s00580-012-1414-9](https://doi.org/10.1007/s00580-012-1414-9).
- Lindsay DS, Blagburn BL, Braund KG. Sarcocystis spp. and sarcocystosis. *Br Med J.* 1995;**5**(3):249-54.
- Anvari D, Narouei E, Hosseini M, Narouei MR, Daryani A, Shariatzadeh SA, et al. Sarcocystosis in ruminants of Iran, as neglected food-borne disease: a systematic review and meta-analysis. *Acta Parasitol.* 2020;**65**(3):555-68. doi: [10.2478/s11686-020-00210-5](https://doi.org/10.2478/s11686-020-00210-5). [PubMed: [32297107](https://pubmed.ncbi.nlm.nih.gov/32297107/)].
- de J. Pena HF, Ogassawara S, Sinhorini IL. Occurrence of cattle sarcocystis species in raw kibbe from Arabian food establishments in the city of Sao paulo, Brazil, and experimental transmission to humans. *J Parasitol.* 2001;**87**(6):1459. doi: [10.2307/3285319](https://doi.org/10.2307/3285319).
- Alibeigi Z, Rahbari S, Hoghooghird N, Naisi S. Macroscopic and microscopic survey of sarcocystosis in ruminants Shahriar slaughterhouse, during 2012-2013. *J Vet Res.* 2015;**70**(4).
- Dong H, Su R, Wang Y, Tong Z, Zhang L, Yang Y, et al. Sarcocystis species in wild and domestic sheep (Ovis ammon and Ovis aries) from China. *BMC Vet Res.* 2018;**14**(1):377. doi: [10.1186/s12917-018-1712-9](https://doi.org/10.1186/s12917-018-1712-9). [PubMed: [30509255](https://pubmed.ncbi.nlm.nih.gov/30509255/)]. [PubMed Central: [PMC6276146](https://pubmed.ncbi.nlm.nih.gov/PMC6276146/)].
- Hosseini SR, Shakerian A, Tahamtan N. Survey of sarcocystis infection in slaughtered sheep in Isfahan, Qom and Shahre-Kord, Iran. *J Anim Vet Adv.* 2012;**11**(15):2683-6. doi: [10.3923/jvavaa.2012.2683.2686](https://doi.org/10.3923/jvavaa.2012.2683.2686).
- Bahari P, Salehi M, Seyedabadi M, Mohammadi A. Molecular identification of macroscopic and microscopic cysts of Sarcocystis in sheep in North Khorasan province, Iran. *Int J Mol Cell Med.* 2014;**3**(1):51.
- Farhang-Pajuh F, Yakhchali M, Mardani K. Molecular determination of abundance of infection with Sarcocystis species in slaughtered sheep of Urmia, Iran. *Veterinary Research Forum: an International Quarterly Journal.* Faculty of Veterinary Medicine, Urmia University, Urmia, Iran; 2014. 181 p.
- Mirzaei M, Rezaei H. The role of sheep in the epidemiology of Sarcocystis spp. in Tabriz area northwest of Iran. *J Parasit Dis.* 2016;**40**(2):285-8. doi: [10.1007/s12639-014-0495-6](https://doi.org/10.1007/s12639-014-0495-6). [PubMed: [27413294](https://pubmed.ncbi.nlm.nih.gov/27413294/)]. [PubMed Central: [PMC4927477](https://pubmed.ncbi.nlm.nih.gov/PMC4927477/)].