



Effect of Interval Training and Curcumin on BAX, Bcl-2, and Caspase-3 Enzyme Activity in Rats

Narges Kashani Vahid¹, Farah Nameni^{1,*} and Bahareh Yazdanparast Chaharmahali¹

¹Department of Sport Physiology, Varamin Pishva Branch, Islamic Azad University, Varamin, Iran

*Corresponding author: Department of Sport Physiology, Varamin Pishva Branch, Islamic Azad University, Varamin, Iran. Email: dastavard96@gmail.com

Received 2021 January 09; Revised 2022 February 07; Accepted 2022 February 13.

Abstract

Background: Apoptosis is a type of programmed cell death in extracellular organisms. High-intensity interval training and curcumin can make some changes in this process.

Objectives: This study aimed to investigate the role of intense interval training with curcumin supplementation on BAX and Bcl-2 proteins and caspase-3 enzyme activity in rats.

Methods: In this study, 48 elderly rats were randomly divided into four groups: (1) control, (2) training, (3) curcumin, and (4) training + curcumin. Then, high-intensity interval training group rats ran on the treadmill for eight weeks, five sessions per week, for 30 - 50 min, and curcumin was fed to the supplement group at 25 mg/kg of body weight three times per week for eight weeks. Gene expression levels of BAX and Bcl-2 and myocardial caspase enzyme were measured in the heart tissue. The Shapiro-Wilk test, one-way ANOVA, and Tukey test were used for data analysis.

Results: Curcumin consumption and intense interval training increased the expression of BAX ($P = 0.001$), Bcl-2 ($P = 0.002$), and caspase ($P = 0.001$). Besides, BAX, Bcl-2, and caspase genes expression significantly changed in the groups compared to the control group. The ratio of BAX to Bcl-2 in the curcumin group and interval training was significantly lower than the other groups. The Tukey post hoc test confirmed a significant difference between the groups and the control group.

Conclusions: High-intensity interval training did not reduce BAX protein, but the training and curcumin supplementation increased Bcl-2 protein expression and neutralized the BAX effect. Curcumin supplementation combined with intense interval training resulted in synergy and reduced cell programming mortality.

Keywords: Caspase, Apoptosis, High-intensity Interval Training

1. Background

Apoptosis is one of the main ways of cell homeostasis in the body that occurs under the control of different genes in multicellular and unicellular organisms (1). Bcl-associated X (BAX) protein and B-cell lymphoma 2 (Bcl-2) protein are significant in forming mitochondrial apoptotic channels. Caspase protease enzymes, including caspase-3 and 7, are also involved in various apoptotic pathways (2).

Curcumin is a polyphenol used as a painkiller, anti-inflammatory agent, and a source of powerful antioxidants. Numerous studies have confirmed the biological effects of curcumin and introduced it into various pathological conditions (3). It contains many anti-diabetic, anti-tumor, anti-inflammatory, and antioxidant agents (4). Curcumin activates the caspase-3 enzyme and releases cytochrome c from the mitochondria to the cytosol (5). Also, curcumin can improve athletic performance and delay ex-

ercise fatigue (6). Studies have shown that curcumin synergizes with 5-fluorouracil and enhances apoptosis in cancer cells in xenograft mice (7). If curcumin enters cancer cells, it activates programmed cell death pathways.

Exercise is another critical factor in promoting health and fighting planned cell death. The simultaneous physiological and immunological effects of curcumin and physical activity have been less studied. Regular exercise can reduce the apoptosis rate in healthy people (8). The impact of interval training on the expression of myocardial infarction and leukemia inhibitory factor (LIF) in skeletal muscle atrophy rats showed that Bcl-2/BAX increased in the interval training group and decreased in the sedentary group (9). Aerobic activity may reduce the signal of mitochondrial apoptosis in the heart muscle by increasing Bcl-2 protein and BAX/Bcl-2 ratio. The BAX protein reduction in mitochondria and the cytochrome c accumulation in the cytosol have also been observed (10). Some studies have

shown that cardiac receptors associated with the obesity-associated apoptosis pathway are activated by exercise. Mitochondrial pathway-dependent apoptosis also decreases with an increase in Bcl-2 and a decrease in BAX and caspase-9 (11). The effects of endurance training on caspase-3 and 9 proteins have been investigated in the tumor tissue of rats with breast cancer (12).

The curcumin consumption at doses of 150 - 1500 mg per day before and during exercise and up to 72 h after exercise could reduce the inflammation caused by physical activity and improve performance. In addition, it appears that humans can tolerate high doses of curcumin without side effects (13). Also, curcumin reduced muscle pain compared to the placebo and decreased post-exercise inflammatory responses without adverse effects. However, it has been reported not to affect TAC, TNF- α , and MDA. Research on curcumin and its mechanical effects on muscle recovery have not yet been performed (14). Nevertheless, the effects of curcumin on inflammatory factors, muscle injury, and exercise training have been investigated in sedentary individuals (15).

2. Objectives

The role of curcumin and other supplements has been investigated in apoptosis and its indicators. Most research on curcumin and exercise has been conducted in the recovery period, focusing on inflammatory, oxidative, cancer, and human factors. However, this study addresses the simultaneous effects of curcumin supplementation and intense interval training, as the physiological changes caused by physical activity can affect the immune system. The antioxidant effects of curcumin are another intervention. Therefore, this study aimed to investigate the effects of intense interval training and curcumin consumption on some indicators of apoptosis and caspase-3 in female rats.

3. Methods

3.1. Groups

In this experimental study, the statistical population included 45 elderly female Wistar rats. Rats were obtained from the Pasteur Institute and housed in the University Laboratory Animal House. Then, 28 mice were randomly selected as a statistical sample. Rats were randomly divided into four groups (control, high-intensity interval training, curcumin, and high-intensity interval training + curcumin). Water was freely available in a 500 ml bottle for the laboratory animals (16). Rats were kept in separate polycarbonate cages with 20 × 27 × 47 cm dimensions. The ambient temperature was set at 22 ± 1.4°C, the light cycle at 12:12

h, and the humidity at 55 ± 0.6%. Rats were fed with foods produced by the Pars-Iran Company Animal Feed Production Center. Also, the water required for each animal was provided in 500 mL bottles for laboratory animals. All research stages were carried out as per the ethical principles of working with animals.

3.2. Exercise Protocol

The training groups were adapted to a rodent treadmill for one week before the start of the protocol (17). The training groups were trained five minutes per day for five days before starting the research protocol to get acquainted with the treadmill. The training program consisted of three sessions per week for six weeks. Each session comprised 10 sets of one-minute activity with 75% intensity for the HIIT group (High-intensity interval training) and two minutes of rest between sets, which started at a speed of 14 meters per minute in the first week. In the following weeks, 2 meters per minute was added to the speed. Until it reached a maximum of 28 meters per minute in the eighth week. Also, five minutes before and after the training, the animals were provided with warming and cooling.

3.3. Supplementation

Curcumin (25 mg) was given to the curcumin and curcumin-training groups by gavage three days a week (18). Eight weeks after the study, all animals with the same conditions followed 13 h fasting, and 48 h after the last training session, they were anesthetized by intraperitoneal injection of ketamine and xylosine to take a sample of heart tissue. The tissue levels of variables were measured using the Western blot method (19).

3.4. Tissue Sampling and Laboratory Measurements

Six weeks after the study, all rats with the same conditions, after 13 hours of fasting and 48 hours of the last training session, were anesthetized with ketamine (60 mg per kg body weight) and xylosine (5 mg per kg body weight) intra-peritoneal with a ratio of 5 to 2. Samples of heart tissue were taken and frozen at -70°C. The tissue levels of variables were measured using the Western blot method (19). We first cut the heart tissue into small pieces in Western blot analysis and homogenized them. This step was performed in the Radio Immunoprecipitation Assay and a protease inhibitor buffer (phenyl-methylsulfonyl fluoride) (complete cocktail tablet, Roche Applied Science, Mannheim, Germany). The supernatant was collected using centrifugation at 12,000 rpm for 20 min. Protein concentrations were determined using BCA (bicinchoninic acid), a protein assay kit. The isolated protein

was loaded and isolated by SDS polyacrylamide gel electrophoresis (sodium dodecyl sulfate), and dissolved proteins were transferred to PVDF (polyvinyl difluoride) membranes. The membranes were blocked with 5% skim milk in tris buffer salt and incubated at 4°C overnight with different primary antibodies. After washing three times with Tween-20 saline buffer, the membranes were proportional to the secondary antibodies (1:12,000) and incubated for 1.5 h.

The next step was to measure changes in BAX, Bcl-2, caspase-3 enzyme activity, and BAX/Bcl-2 ratio. Devices were used to process the data and indicate whether the cells underwent apoptosis or not.

3.5. Statistical Analysis

The normal distribution of data was assessed using the Shapiro-Wilk test, and the homogeneity of variances using the Levin test. Significant changes in variables were determined using ANOVA and the Tukey post hoc test ($P \leq 0.05$).

4. Results

Mean changes in caspase-3, BAX, and Bcl-2 protein expression were determined as the mean and standard deviation. The Shapiro-Wilk test was normal. The Leven test confirmed the homogeneity of variances.

The results showed that caspase-3 had the highest increase in the intense interval training and intense interval training + curcumin groups. One-way ANOVA evaluated the difference between the mean caspase-3 values. According to the test results, the expression of caspase-3 protein was significantly different between the groups. The Tukey post hoc test confirmed a significant increase in the three experimental groups compared to the control group (Figure 1).

The results showed that BAX had the highest increase in the intense interval training + curcumin and intense interval training groups. One-way ANOVA ($P = 0.001$) evaluated changes in BAX protein expression. According to the test results, the BAX protein expression was significantly different between the groups. Tukey post hoc test confirmed a significant decrease in exercise + curcumin and curcumin groups compared with the control and exercise groups. (Figure 2).

The results showed that Bcl-2 had the highest increase in the intense interval training and intense interval training + curcumin groups. One-way ANOVA ($P = 0.002$) evaluated the differences between the means of Bcl-2. According to the test results, the Bcl-2 protein expression was significantly different between the groups. The Tukey post hoc test confirmed a significant increase between the experimental groups compared to the control group. (Figure 3).

The results showed that the BAX to Bcl-2 ratio had the highest decrease in the intense interval training + curcumin and intense interval training groups. The ratio of BAX to Bcl-2 was significantly different between the groups. The Tukey post hoc test confirmed the significant difference between all groups and the control group (Figure 4).

5. Discussion

The results showed that the expression of BAX proapoptotic protein decreased in the interval training-curcumin supplement and curcumin supplement groups, but the anti-apoptotic protein Bcl-2 increased. This study showed that HIIT was an essential factor in the induction of apoptosis, and the expression of Bax, Bcl-2, and caspase-3 increased during HIIT. The ratio of BAX to Bcl-2 was lower in the supplement + exercise group than in the other groups, indicating a direct effect of the two independent variables. The ratio of BAX to Bcl-2 decreased. Apoptosis limits tissue damage and induces direct signals for adaptation induced by mediators such as reactive oxygen species, cytokines, and hormones.

Exercise creates physiological stress and induces adaptive responses. This confirms the decrease in apoptosis in the exercise + curcumin supplement group. Physical activity stimulates the apoptotic pathways and affects the balance between the release of free radicals and the antioxidant defense mechanism. Reactive oxygen species also affect apoptotic pathways. However, training and curcumin reduced this interaction. One of the possibilities of these changes is the direct prevention of apoptosis due to exercise protocol and curcumin supplementation. Interval training and supplementation can increase heart rate and blood lactate threshold level, improve body composition, and decrease fat percentage. Also, increased stimulation of the sarcoplasmic reticulum and contractile pumps in the mitochondrial membrane makes molecular changes, consistent with the results of Siddighi's research. Maintaining homeostasis and oxidative compatibility can reduce apoptosis. Of course, Sedighi et al. used endurance training to assess cardiac apoptosis. The tissue studied and the length of the training period are the reasons for the similarity of the results of the two studies (20). Exercise duration and curcumin use can significantly increase Bcl-2 and BAX levels and the BAX/ Bcl-2 ratio depending on the exercise intensity.

In addition, the combination of exercise and curcumin had an antagonistic effect. Chiu et al. studied breast cancer and apoptotic agents and obtained different results (21). The study of Safarzadeh Gargari et al. also showed that the expression of proteins and their ratio were not significantly different from the control group. In their study,




Figure 1. Caspase-3 expression levels in four research groups. *Significant increases compared to the control group at $P \leq 0.05$.




Figure 2. BAX expression levels in four research groups. #Significant increases compared to the control group at $P \leq 0.05$. *Significant decreases curcumin group to another group at $P \leq 0.05$.




Figure 3. Bcl-2 expression levels in four research groups. #Significant increases compared to the control group at $P \leq 0.05$.




Figure 4. Bax/Bcl-2 expression levels in four research groups. #Significant decreases compared to the control group at $P \leq 0.05$.

hydrogen peroxide injection into the tissue was much stronger than endurance training, and apoptosis was not stopped (22). In the study of Karimi et al., the expression of BAX and Bcl-2 genes did not change significantly, but caspase-3 activity showed a significant increase in the liver

of diabetic rats before and after treatment with aqueous garlic extract. The discrepancy between the results seems to be due to the use of garlic supplements and sampled tissue (23, 24). The main reason for the alignment of the results is the direct and potential impact of exercise and

matched texture.

Severe oxidative stress destabilizes mitochondrial membrane homeostasis, stimulates the formation of permeable pores of mitochondrial membranes, and releases pre-apoptotic agents such as cytochrome c. Exercise can have the opposite effect and reduce pro-apoptotic genes by increasing antioxidant capacity and modulating stress (25). Intense exercise triggers an inflammatory response, and taking supplements can be effective too. Intense activity, unlike moderate aerobic activity, causes DNA to fragment and caspase cascades to begin (26). Flavonoid and cerebroside compounds in curcumin supplementation can reduce these effects. In this study, HIIT reduced the expression of pro-apoptotic genes by increasing antioxidant capacity and modulating oxidative stress (14). Therefore, training and curcumin supplementation prevented the increase of caspase-3 expression, while apoptosis was reduced due to the compatibility of training and supplementation.

5.1. Conclusions

High-intensity interval training did not affect reducing BAX protein. However, training and curcumin supplementation increased the expression of Bcl-2 protein, and this ratio has been able to reduce the effect of BAX. Therefore, curcumin supplementation and intermittent exercise can reduce cell apoptosis and programmed cell death.

Footnotes

Authors' Contribution: Study concept and design: NK, FN, and BY; acquisition of data: FN; analysis and interpretation of data: NK, FN, and BY; drafting of the manuscript: NK, FN, and BY; critical revision of the manuscript for important intellectual content: FN; statistical analysis: FN and NK; administrative, technical, and material support: NK; study supervision: FN.

Clinical Trial Registration Code: Researchers received introduction letters from the Varamin Pishva Branch of Islamic Azad University with code IR.IAU.VARAMIN.REC.1398.009.

Conflict of Interests: There is no conflict of interest.

Data Reproducibility: It was not declared by the authors.

Ethical Approval: IR.IAU.VARAMIN.REC.1398.009. Link: ethics.research.ac.ir/EthicsProposalView.php?id=71209

Funding/Support: There is no funding/support.

References

1. Russo V, Candeloro P, Malara N, Perozziello G, Iannone M, Scicchitano M, et al. Key Role of Cytochrome C for Apoptosis Detec-

- tion Using Raman Microimaging in an Animal Model of Brain Ischemia with Insulin Treatment. *Appl Spectrosc.* 2019;**73**(10):1208-17. doi: [10.1177/0003702819858671](https://doi.org/10.1177/0003702819858671). [PubMed: [31219322](https://pubmed.ncbi.nlm.nih.gov/31219322/)].
2. Jafari A, Pourrazi H, Nikookheslat S, Baradaran B. Effect of Exercise Training on Bcl-2 and Bax Gene Expression in the Rat Heart. *Gene Cell Tissue.* 2015;**2**(4). doi: [10.17795/gct-32833](https://doi.org/10.17795/gct-32833).
3. Aktas C, Kanter M, Kocak Z. Antiapoptotic and proliferative activity of curcumin on ovarian follicles in mice exposed to whole body ionizing radiation. *Toxicol Ind Health.* 2012;**28**(9):852-63. doi: [10.1177/0748233711425080](https://doi.org/10.1177/0748233711425080). [PubMed: [22075690](https://pubmed.ncbi.nlm.nih.gov/22075690/)].
4. Zhao L, Gu Q, Xiang L, Dong X, Li H, Ni J, et al. Curcumin inhibits apoptosis by modulating Bax/Bcl-2 expression and alleviates oxidative stress in testes of streptozotocin-induced diabetic rats. *Ther Clin Risk Manag.* 2017;**13**:1099-105. doi: [10.2147/TCRM.S141738](https://doi.org/10.2147/TCRM.S141738). [PubMed: [28894373](https://pubmed.ncbi.nlm.nih.gov/28894373/)]. [PubMed Central: [PMC5584885](https://pubmed.ncbi.nlm.nih.gov/PMC5584885/)].
5. Zhu L, Han MB, Gao Y, Wang H, Dai L, Wen Y, et al. Curcumin triggers apoptosis via upregulation of Bax/Bcl-2 ratio and caspase activation in SW872 human adipocytes. *Mol Med Rep.* 2015;**12**(1):151-6. doi: [10.3892/mmr.2015.3450](https://doi.org/10.3892/mmr.2015.3450). [PubMed: [25760477](https://pubmed.ncbi.nlm.nih.gov/25760477/)].
6. Hadi A, Pourmasoumi M, Ghaedi E, Sahebkar A. The effect of Curcumin/Turmeric on blood pressure modulation: A systematic review and meta-analysis. *Pharmacol Res.* 2019;**150**:104505. doi: [10.1016/j.phrs.2019.104505](https://doi.org/10.1016/j.phrs.2019.104505). [PubMed: [31647981](https://pubmed.ncbi.nlm.nih.gov/31647981/)].
7. Zhang P, Lai ZL, Chen HF, Zhang M, Wang A, Jia T, et al. Curcumin synergizes with 5-fluorouracil by impairing AMPK/ULK1-dependent autophagy, AKT activity and enhancing apoptosis in colon cancer cells with tumor growth inhibition in xenograft mice. *J Exp Clin Cancer Res.* 2017;**36**(1):190. doi: [10.1186/s13046-017-0661-7](https://doi.org/10.1186/s13046-017-0661-7). [PubMed: [29273065](https://pubmed.ncbi.nlm.nih.gov/29273065/)]. [PubMed Central: [PMC5741949](https://pubmed.ncbi.nlm.nih.gov/PMC5741949/)].
8. Santana ET, Serra AJ, Silva Junior JA, Bocalini DS, Barauna VG, Krieger JE, et al. Aerobic exercise training induces an anti-apoptotic milieu in myocardial tissue. *Mot Rev Educ.* 2014;**20**(2):233-8. doi: [10.1590/s1980-65742014000200015](https://doi.org/10.1590/s1980-65742014000200015).
9. Jia D, Cai M, Xi Y, Du S; ZhenjunTian. Interval exercise training increases LIF expression and prevents myocardial infarction-induced skeletal muscle atrophy in rats. *Life Sci.* 2018;**193**:77-86. doi: [10.1016/j.lfs.2017.12.009](https://doi.org/10.1016/j.lfs.2017.12.009). [PubMed: [29223542](https://pubmed.ncbi.nlm.nih.gov/29223542/)].
10. Peterson JM, Bryner RW, Sindler A, Frisbee JC, Alway SE. Mitochondrial apoptotic signaling is elevated in cardiac but not skeletal muscle in the obese Zucker rat and is reduced with aerobic exercise. *J Appl Physiol (1985).* 2008;**105**(6):1934-43. doi: [10.1152/jappphysiol.00037.2008](https://doi.org/10.1152/jappphysiol.00037.2008). [PubMed: [18832755](https://pubmed.ncbi.nlm.nih.gov/18832755/)]. [PubMed Central: [PMC2612474](https://pubmed.ncbi.nlm.nih.gov/PMC2612474/)].
11. Lee SD, Shyu WC, Cheng IS, Kuo CH, Chan YS, Lin YM, et al. Effects of exercise training on cardiac apoptosis in obese rats. *Nutr Metab Cardiovasc Dis.* 2013;**23**(6):566-73. doi: [10.1016/j.numecd.2011.11.002](https://doi.org/10.1016/j.numecd.2011.11.002). [PubMed: [22402061](https://pubmed.ncbi.nlm.nih.gov/22402061/)].
12. Kazemi A, Mirza-zade E. The Effect of Endurance Training on Tumor Tissue Levels of Caspase-3 and Caspase-9 in Mice with Breast Cancer. *Iran J Breast Dis.* 2018:32-43. doi: [10.30699/acadpub.ijbd..11.3.32](https://doi.org/10.30699/acadpub.ijbd..11.3.32).
13. Fernandez-Lazaro D, Mielgo-Ayuso J, Seco Calvo J, Cordova Martinez A, Caballero Garcia A, Fernandez-Lazaro CI. Modulation of Exercise-Induced Muscle Damage, Inflammation, and Oxidative Markers by Curcumin Supplementation in a Physically Active Population: A Systematic Review. *Nutrients.* 2020;**12**(2). doi: [10.3390/nu12020501](https://doi.org/10.3390/nu12020501). [PubMed: [32075287](https://pubmed.ncbi.nlm.nih.gov/32075287/)]. [PubMed Central: [PMC7071279](https://pubmed.ncbi.nlm.nih.gov/PMC7071279/)].
14. Basham S, Waldman H, Krings B, Lamberth J, Smith J, McAllister M. Effect of Curcumin Supplementation on Exercise-Induced Oxidative Stress, Inflammation, Muscle Damage, and Muscle Soreness. *J Diet Suppl.* 2020;**17**(4):401-14. doi: [10.1080/19390211.2019.1604604](https://doi.org/10.1080/19390211.2019.1604604). [PubMed: [31025894](https://pubmed.ncbi.nlm.nih.gov/31025894/)].
15. Dias KA, da Conceicao AR, Oliveira LA, Pereira SMS, Paes SDS, Monte LF, et al. Effects of Curcumin Supplementation on Inflammatory Markers, Muscle Damage, and Sports Performance during Acute Physical Exercise in Sedentary Individuals. *Oxid Med Cell Longev.* 2021;**2021**:9264639. doi: [10.1155/2021/9264639](https://doi.org/10.1155/2021/9264639). [PubMed: [34659641](https://pubmed.ncbi.nlm.nih.gov/34659641/)]. [PubMed Central: [PMC8516555](https://pubmed.ncbi.nlm.nih.gov/PMC8516555/)].

16. Brown MB, Neves E, Long G, Graber J, Gladish B, Wiseman A, et al. High-intensity interval training, but not continuous training, reverses right ventricular hypertrophy and dysfunction in a rat model of pulmonary hypertension. *Am J Physiol Regul Integr Comp Physiol.* 2017;**312**(2):R197-210. doi: [10.1152/ajpregu.00358.2016](https://doi.org/10.1152/ajpregu.00358.2016). [PubMed: [27784688](https://pubmed.ncbi.nlm.nih.gov/27784688/)]. [PubMed Central: [PMC5504398](https://pubmed.ncbi.nlm.nih.gov/PMC5504398/)].
17. Abdelbasset WK, Tantawy SA, Kamel DM, Alqahtani BA, Soliman GS. A randomized controlled trial on the effectiveness of 8-week high-intensity interval exercise on intrahepatic triglycerides, visceral lipids, and health-related quality of life in diabetic obese patients with nonalcoholic fatty liver disease. *Medicine (Baltimore).* 2019;**98**(12). e14918. doi: [10.1097/MD.00000000000014918](https://doi.org/10.1097/MD.00000000000014918). [PubMed: [30896648](https://pubmed.ncbi.nlm.nih.gov/30896648/)]. [PubMed Central: [PMC6708753](https://pubmed.ncbi.nlm.nih.gov/PMC6708753/)].
18. Li W, Suwanwela NC, Patumraj S. Curcumin prevents reperfusion injury following ischemic stroke in rats via inhibition of NFkappaB, ICAM-1, MMP-9 and caspase-3 expression. *Mol Med Rep.* 2017;**16**(4):4710-20. doi: [10.3892/mmr.2017.7205](https://doi.org/10.3892/mmr.2017.7205). [PubMed: [28849007](https://pubmed.ncbi.nlm.nih.gov/28849007/)]. [PubMed Central: [PMC5647023](https://pubmed.ncbi.nlm.nih.gov/PMC5647023/)].
19. Hunter AL, Choy JC, Granville DJ. Detection of Apoptosis in Cardiovascular Diseases. *Molecular Cardiology.* Humana Press Inc; 2005. p. 277-89.
20. Sadighi A, Abdi A, Azarbayjani MA, Barari A. [Effect of aerobic exercise on some factors of cardiac apoptosis in male rats]. *Feyz J Kashan Univ Med Sci.* 2019;**23**(5):495-502. Persian.
21. Chiu TL, Su CC. Curcumin inhibits proliferation and migration by increasing the Bax to Bcl-2 ratio and decreasing NF-kappaBp65 expression in breast cancer MDA-MB-231 cells. *Int J Mol Med.* 2009;**23**(4):469-75. doi: [10.3892/ijmm_00000153](https://doi.org/10.3892/ijmm_00000153). [PubMed: [19288022](https://pubmed.ncbi.nlm.nih.gov/19288022/)].
22. Safrazadeh Gargari S, Matin Homaei H, Azarbayjani MA. [Effects of continuous exercise on BAX and BCL-2 heart proteins following by different dos of H2O2 consumption in rat male]. *J Shahid Sadoughi Univ Med Sci.* 2018;**26**(4):363-79. Persian.
23. Karimi MN, Abbasalipourkabir R, Arab Sadeghabadi Z, Ziamajidi N. [The level of gene expression of Bax and Bcl-2 and the activity of caspase 3 in the liver tissues of normal, type 1 and type 2 diabetic rats before and after treatment with aqueous extract of garlic]. *J Shahid Sadoughi Univ Med Sci.* 2017;**25**(7):547-55. Persian.
24. Yang J, Ning J, Peng L, He D. Effect of curcumin on Bcl-2 and Bax expression in nude mice prostate cancer. *Int J Clin Exp Pathol.* 2015;**8**(8):9272-8. [PubMed: [26464676](https://pubmed.ncbi.nlm.nih.gov/26464676/)]. [PubMed Central: [PMC4583908](https://pubmed.ncbi.nlm.nih.gov/PMC4583908/)].
25. Sharafi H, Rahimi R. The effect of resistance exercise on p53, caspase-9, and caspase-3 in trained and untrained men. *J Strength Cond Res.* 2012;**26**(4):1142-8. doi: [10.1519/JSC.0b013e31822e58e5](https://doi.org/10.1519/JSC.0b013e31822e58e5). [PubMed: [22446679](https://pubmed.ncbi.nlm.nih.gov/22446679/)].
26. Siahkohian M, Asgharpour Arshad M, Bolboli L, Jafari A, Sheikhzadeh Hesari F. [Effect of 12-Weeks Aerobic Training on Some Indices of Skeletal Muscle Apoptosis In Male Rats]. *Med J Tabriz Univ Med Sci Health Serv.* 2018;**39**(6):35-43. Persian.