Published online 2023 January 22.

Synergistic Effect of Taxol and Vincristine Against MCF-7 Cell Line

Mohammad Zandi 回 1,*

¹Department of Agriculture, Iranian Research Organization for Science and Technology (IROST), 33535111, Tehran, Iran

^{*} Corresponding author: Department of Agriculture, Iranian Research Organization for Science and Technology (IROST), 33535111, Tehran, Iran. Email: mz1075@yahoo.com

Received 2022 November 05; Revised 2022 December 04; Accepted 2022 December 10.

Abstract

Background: Women worldwide are concerned about breast cancer, resulting in high mortality and morbidity.

Objectives: In this research, to investigate the function of Taxol, Vinblastine, and Vincristine on the MCF-7 cell line, different concentration levels (10, 20, 40, 80, 160, and 320 μ mol/mL) of the drugs were examined. Taxolwas also used in combination with Vinblastine and Vincristine.

Methods: The examined compounds were applied to the cells for 48 h. The MTT test was employed for assessing cytotoxicity, and the nonlinear regression method was run to estimate the values of IC50 (half maximal inhibitory concentration) values.

Results: It was found that taxol + vincristine, taxol, and vinblastine treatments with concentrations of 41.45, 64.46, and 67.12 μ mol/mL resulted in the best 50% inhibitory effect. The lowest effective concentration that had significant effects on cancer cells was vinblastine (160 μ mol/mL), taxol + vincristine (80 μ mol/mL), taxol (80 μ mol/mL), taxol + vinblastine (80 μ mol/mL), and vincristine (40 μ mol/mL), respectively (P < 0.05).

Conclusions: From the results of the current study, a synergistic reaction was found between taxol and vincristine based on calculated CI values.

Keywords: MCF-7, Taxol, Vinblastine, Vincristine

1. Background

Cancer is considered as the leading cause of death in the sense that there have been almost 13 million newly diagnosed cases and 8 million deaths within a year (1). One of the most widely used breast cancer cell line models is the breast adenocarcinoma cell line (MCF-7), which is used for looking at anti-cancer drugs (2). Women in the United States suffer from MCF-7 as one of the most prevalent women's cancers, with 266,120 new cases and 40,920 breast cancer deaths in 2017 (3). According to Jemal et al.'s final report, the yearly rate of breast cancer is enhancingby 0.4% globally (4). Chemotherapy is often used as adjuvant anti-cancer therapy. Chemotherapy is applied to most patients prior to or after surgery (1).

Attempts to explore new cancer chemotherapeutics have led to the introduction of natural products. The substances obtained from plants have been determined as models for developing new drugs. The adaptation of plants results in synthesizing second metabolites whose compounds additionally exist in pharmacological activities, amongst which the antibacterial, antihypertensive, anti-inflammatory, and anti-cancer tasks are the most regularly reported ones (5).

Taxol (generic name paclitaxel) is a clinically effective antitumor drug being approved for treating breast carcinomas (6). Also, taxol is a poly-oxygenated cyclic diterpenoid with a characteristic taxane ring system. This drug has been the most efficient and pervasively utilized chemotherapeutic for treating cancers and virus-related sarcomas. Besides, Kumar et al.'s therapy of cells with taxol hinders the regular reconstruction of the microtubule network and also prevents the regular spindle development at the metaphase required for mitosis as well as cell expansion (7). This yields the cells' arrestment at the G2/M phase of the cell cycle and apoptotic cell death as a consequence (8). Taxol and other hydrophobic antitumor drugs are substrates for P-glycoprotein, which is the MDR1 gene product. Along with the overexpression of P-glycoprotein, other actively involved mechanisms consist of c-erbB2/neuoverexpression in breast cancer cells, changes in L-tubulin isotypes, anomalies in tubulin and also particular changes in many properties of signal transduction paths. The activation of Raf-1 and mitogenactivated protein kinase (MAP kinase), which are molecular changes in molecules engaged with the cascades of signal transduction, has resulted after treating cells that are sensitive to drugs with taxol (9).

Copyright © 2023, Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (http://creativecommons.org/licenses/by-nc/4.0/) which permits copy and redistribute the material just in noncommercial usages, provided the original work is properly cited.

Vinblastine and Vincristine effectively applied in chemotherapy regimens, are recognized as the first two vinca alkaloid compounds. These agents resolve to apprehend the separating cells at metaphase by binding to the β -subunit of tubulin heterodimers to avoid polymerization and incorporation into microtubules (10). The different vinca alkaloids have their unique properties. For example, vinblastine has an angiogenesis inhibitory effect. It is also related to the anti-diuretic hormonal agent secretion and angina, which are used to deal with Hodgkin's illness, non-Hodgkin's lymphoma, and breast cancer (11). Vincristine contains a high fondness for tubulin dimers, and its response to dimers is rapidly reversible. This implies that a vincristine particle will affix to a dimer at one site, break off, and afterward reattach to an additional site, resulting in two sites per dimer "poisoned" and incapable of reassembling into the protein. Hence, vincristine is potentially able to destabilize tubulin (12). Although these alkaloids meet only minor structural differences, namely the methyl group on the vindoline N atom in vinblastine replaced by a formyl group in vincristine, they significantly differ in terms of the toxicity and clinical activity spectrum. Neuropathy has often been observed following vincristine administration, while the dose-limiting toxicity for vinblastine has been carried out by myelosuppression (13).

Additionally, mixed treatment is an appealing technique for minimizing the side effects of vinca alkaloids, which are incorporated into various chemotherapy medicines to boost their results. Medications are commonly administered simultaneously as a cocktail or sequentially to optimize their therapeutic impact (14).

2. Objectives

The present study has emphasized breast cancer treatment using potential anti-cancer drugs. Here, a possible synergism has been hypothesized among taxol, vinblastine, and vincristine against the MCF-7 breast cancer cell line.

3. Methods

3.1. Chemicals

Dulbecco's modified eagle medium (DMEM) and penicillin-streptomycin from Inoclon (Iran), and fetal bovine serum (FBS) from Gibco (USA), were responsible for providing the culture media, and other chemicals such as phosphate-buffer saline (PBS), MTT (code 1001904784) and DMSO from Sigma-Aldrich (Germany). Besides purchasing taxol, vinblastine, and vincristine from Sobhan oncology, Iran, the plastic was done from Sorfa (China).

3.2. Human Cancer Cell Line

Iranian Biological Research Center (IBRC) contributed to purchasing the human cancer cell line. The cell line, which has been employed in the present study, includes human breast adenocarcinoma (MCF-7; IBRC No. C10682). The IBRC instructions were employed for culturing the cells. DMEM was specified as the culture medium. Penicillin, streptomycin, and 10% fetal bovine serum were used to supplement the medium. Moreover, a humidified incubator at 37°C and 5% CO2 was used to maintain the cells.

3.3. MTT Assay

Trypsinizing the MCF-7 cells were conducted at 80 -90% confluence. They were also plated into a 96-well culture dish whose density was found to be 1 \times 10⁴ cells/well overnight. The exposure of cells to different concentrations of taxol, vinblastine, and vincristine was carried out. Besides, they were combined with taxol for 48 h. The modified version of Shabani et al.'s (15) viability test of MTT, including 3-(4,5-dimethylthiazol-2-yl)-2,5diphenyl tetrazolium bromide, was done. Changing the cell medium was conducted with a fresh one (100 μ L per well) after incubation. Besides, 50 μ L of MTT solution (0.5 mg/mL in PBS) was added. Incubation of the plates was done for four h at 37°C. Dissolution of the formed formazan crystals was also carried out in 100 μ L DMSO per well upon mixing. In order to measure the absorbance at 570 nm, a 96-well plate reader was used. The control's cell viability was presented as a percentage of cytotoxicity aligned with cell viability. The untreated cell viability was found to be 100% when all cytotoxicity was investigated. In addition, nonlinear regression via AAT Bioquest program (www.aatbio.com) was employed for estimating the IC50 values.

3.4. Experimental Design

As to the evaluation of taxol, vinblastine, and vincristine performance on the MCF-7 cell line, different concentration levels (10, 20, 40, 80, 160, and 320 μ mol/mL) of each of the drugs were examined using the MTTcellproliferationassay. Based on the cytotoxicity results of the studied treatments, the synergistic effects of taxol, along with each of the vinblastine and vincristine treatments, were studied.

3.5. Synergy Testing

The Chou-Talalay method was used to assess interactions taking place among taxol, vinblastine, and vincristine on MCF-7 cells. Besides, CalcuSyn software (Biosoft, Cambridge, UK) was used for generating CI values.

3.6. Statistical Analysis

The repetition of each experiment occurred at least three times. A statistical software program (SPSS 16) was run for data analysis. Statistics through one-way ANOVA and Duncan measure were done to examine whether there were significant differences among the mean values. Results have been stated as mean \pm SEM, and a P-value of less than 0.05 (P < 0.05) stands for the statistically significant difference.

4. Results

4.1. Effectsof Taxol, Vinblastine, and Vincristine on MCF-7 Cell Line

The cytotoxicity of taxol, vinblastine, and vincristine was investigated separately, and the calculation of IC50 for all drugs was conducted. To this end, MCF-7 cells were incubated with different concentrations of taxol, vinblastine, and vincristine (Figure 1). The cytotoxicity of the examined drugs was evaluated within a concentration range of 10 -320 μ mol/mL. The present results indicated that with increasing concentrations of taxol (from 10 to 20 μ mol/mL), the cell viability significantly decreases after 48 h of incubation (P < 0.05). At a taxol concentration level of 80 μ mol/mL, the estimation of cancer cells survival was presented as 40.17 \pm 2.59. Investigating the influence of vinblastine on MCF-7 cells, it can be concluded that the survival of cancer cells (25.22 ± 0.09) significantly decreases with an increment in the concentration level up to 160 μ mol/mL (P < 0.05) (Figure 1).

The results corresponding to the effect of vincristine on MCF-7 cells indicated that the survival of cancer cells at 40 μ mol/mL has significantly decreased (60.67 ± 3.61) (P < 0.05).

The results obtained for the effect of the taxol + vinblastine combination on the survival of MCF-7 cells indicated that there is a significant decrease in the survival of cancer cells up to 80 μ mol/mL (58.9 ± 3.38) (P < 0.05) (Figure 1).

The effect of combining various concentrations of taxol and vincristine on cancer cells indicated that the 80 μ mol/mL level of concentration significantly reduces the survival of the examined cells (39.01 ± 5.8) (P < 0.05) (Figure 1).

4.2. Cytotoxicity in MCF-7 Cell Line

The examined compounds were applied to the cells for 48 h at different concentrations ranging from 10 - 320 μ mol/mL. The MTT test assessed the cytotoxicity, and the values of IC50 were also estimated. Table 1 presents the findings of 50% inhibitory features associated with the investigated experiments. The best inhibitory effects are associated with the taxol + vincristine, taxol, and vinblastine treatments with concentrations of 41.45, 64.46, and 67.12 μ mol/mL, respectively.

4.3. Synergistic Effect of Taxol

When loading of the surviving fractions into CalcuSyn software was carried out, CI (combination index), which is derived from the Chou and Talalay method for assessing synergy, antagonism, or addition, was calculated. Results revealed that a synergistic reaction was recorded between the taxol and vincristine combination of drugs (CI < 1).

5. Discussion

Today, one way to increase the anti-cancer effect is by combining chemotherapy with natural compounds, which can lead to improving the therapeutic window and reversing resistance to the drug (16). The present results indicated that taxol at a concentration of 64.46 μ mol/mL meets 50% inhibitory properties on MCF-7 cells. The calculated IC50 values were higher than those of vinblastine (67.12 μ mol/mL) and vincristine (239.51 μ mol/mL). Cell cycle arrest and apoptosis might result from exposing cells to taxol, which might lead to effects that align with the different concentrations of drugs and employed tissues and/or cell lines (8). A number of researchers have acknowledged that taxol can activate various signal transduction pathways. For instance, (1) the activation process of c-jun Nterminal kinase (JNK) can be carried out by taxol using the Ras and apoptosis signal-regulating kinase (ASK1) pathways. JNK and p38 can also be activated by ASK1, which is a MAP kinase. When there is TNF-K treatment, apoptotic cell death can be induced by ASK1; (2) when extracellular signalregulated kinases (ERK) are to be activated, the process is induced by taxol. Taxol-treated RAW 264.7 cells were also significant in forming anShc/Grb2 complex and Tyrosine phosphorylation of Shc; (3) p38 tyrosine phosphorylation results from taxol (9). Additionally, several investigations have indicated that the taxol resistance in breast cancer cells might be due to the increased levels of β III isotypes (17). The major cause of the reduction of clinical effects can be attributed to the achieved resistance to taxol, unfortunately (16). In this regard, the present study examined the effect of taxol along with vinca alkaloids. According to the results obtained, the best inhibitory effect is related to taxol + vincristine treatment with concentrations of 41.45 μ mol/mL. The use of combination therapy has been shown to play a key role in the therapy of cancer. It also results in a more therapeutic improvement than the monotherapeutic modality (2). Possessing the ability to induce specific



Table 1. The Investigated Compounds' Cytotoxicity Against the MCF-7 Cell Line					
Treatments (µmol/mL)					
Cell Line	Taxol	Vinblastine	Vincristine	Taxol + Vinblastine	Taxol + Vincristine
MCF-7	64.46	67.12	239.51	236.71	41.45

changes in DNA, such as micronuclei and oxidative, but not others (i.e., SCEs) appears to be the potential of vinca alkaloids. This ability can be attributed to the vincaalkaloid's performance characteristics, which is in alignment with apoptosis activation. The activation process proceeds when there is interference with spindle fiber formation and mitochondrial function. Furthermore, changes in models, doses, and duration regarding the treatment of the drug can be caused by improper variations occurring in genotoxicity, which are aligned with vincristine in vitro cultured cell methods and mammalian models (11). The side effects of vinca alkaloids can be reduced by a promising approach known as combination therapy. In other words, a combination occurs when other chemotherapy drugs are involved, which can increase the potential for antitumor treatment. Administering the drugs occurs concurrently as a cocktail, or it might take place sequentially to enhance their therapeutic effectiveness (14).

Being resistant to chemotherapy can be a significant hindrance to therapeutic success. There appears to be significant demand to seek more effective techniques for therapy (16). Cell viability measurement was conducted to specify the most synergistic effect for the different combinations. Results revealed that, according to the measured CI, synergistic performance occurred between taxol and vincristine.

Acknowledgments

The Agricultural Institute of IROST is greatly appreciated for preparing the required facilities for the laboratory.

Footnotes

Authors' Contribution: MZ, conceptualization, formal analysis, writing – review & editing.

Conflict of Interests: The author declares no conflict of interest. The author is grateful to the Agricultural Institute of IROST for providing this project with laboratory facilities and other technical support.

Ethical Approval: This is an in vitro study, and ethical approval is optional. The protocol used in the present research was approved by the Iranian Research Organization for Science and Technology(IROST)(approval number: 321713).

Funding/Support: This study was supported in part by grant 1011097005 from the Iranian Research Organization for Science and Technology (IROST) and by a research scholarship from IROST (Dr. Mohammad Zandi).

References

- Hassanzadeganroudsari M, Heydarinasab A, Soltani M, Chen P, Akbarzadeh Khiyavi A. Enhancing anti-cancer efficacy of carboplatin by PEGylated poly(butyl cyanoacrylate) nano-particles. J Drug Deliv Sci Technol. 2019;54:101218. https://doi.org/10.1016/j.jddst.2019.101218.
- Odeh F, Naffa R, Azzam H, Mahmoud IS, Alshaer W, Al Bawab A, et al. Co-encapsulation of thymoquinone with docetaxel enhances the encapsulation efficiency into PEGylated liposomes and the chemosensitivity of MCF7 breast cancer cells to docetaxel. *Heliyon*. 2019;5(11). e02919. [PubMed ID: 31844767]. [PubMed Central ID: PM-CPmc6895652]. https://doi.org/10.1016/ji.heliyon.2019.e02919.
- Siegel RL, Miller KD, Jemal A. Cancer statistics, 2018. CA Cancer J Clin. 2018;68(1):7-30. [PubMed ID: 29313949]. https://doi.org/10.3322/caac.21442.
- Jemal A, Ward EM, Johnson CJ, Cronin KA, Ma J, Ryerson B, et al. Annual Report to the Nation on the Status of Cancer, 1975-2014, Featuring Survival. J Natl Cancer Inst. 2017;109(9). [PubMed ID: 28376154]. [PubMed Central ID: PMCPmc5409140]. https://doi.org/10.1093/jnci/djx030.
- Fagundes Rosales P, Gower A, Lucia Ruiz Benitez M, Silveira Pacheco B, Vieira Segatto N, Roesch-Ely M, et al. Extraction, isolation and in vitro evaluation of affinisine from Tabernaemontana catharinensis in human melanoma cells. *Bioorg Chem.* 2019;90:103079. [PubMed ID: 31255990]. https://doi.org/10.1016/j.bioorg.2019.103079.
- Yang CP, Horwitz SB. Distinct mechanisms of taxol-induced serine phosphorylation of the 66-kDa Shc isoform in A549 and RAW 264.7 cells. *Biochim Biophys Acta*. 2002;**1590**(1-3):76-83. [PubMed ID: 12063170]. https://doi.org/10.1016/s0167-4889(02)00200-8.
- Kumar P, Singh B, Thakur V, Thakur A, Thakur N, Pandey D, et al. Hyper-production of taxol from Aspergillus fumigatus, an endophytic fungus isolated from Taxus sp. of the Northern Himalayan region. *Biotechnol Rep (Amst)*. 2019;24. e00395.

[PubMed ID: 31799144]. [PubMed Central ID: PMCPmc6881681]. https://doi.org/10.1016/j.btre.2019.e00395.

- Panno ML, Giordano F, Mastroianni F, Morelli C, Brunelli E, Palma MG, et al. Evidence that low doses of Taxol enhance the functional transactivatory properties of p53 on p21 waf promoter in MCF-7 breast cancer cells. *FEBS Lett.* 2006;**580**(9):2371-80. [PubMed ID: 16616141]. https://doi.org/10.1016/j.febslet.2006.03.055.
- Yang CP, Galbiati F, Volonte D, Horwitz SB, Lisanti MP. Upregulation of caveolin-1 and caveolae organelles in Taxol-resistant A549 cells. *FEBS Lett.* 1998;**439**(3):368-72. [PubMed ID: 9845355]. https://doi.org/10.1016/s0014-5793(98)01354-4.
- Mora E, Smith EM, Donohoe C, Hertz DL. Vincristine-induced peripheral neuropathy in pediatric cancer patients. *Am J Cancer Res.* 2016;6(11):2416–30. [PubMed ID: 27904761]. [PubMed Central ID: PM-CPmc5126263].
- Mhaidat NM, Alzoubi KH, Khabour OF, Alawneh KZ, Raffee LA, Alsatari ES, et al. Assessment of genotoxicity of vincristine, vinblastine and vinorelbine in human cultured lymphocytes: a comparative study. *Balkan J Med Genet.* 2016;19(1):13–20. [PubMed ID: 27785403]. [PubMed Central ID: PMCPmc5026275]. https://doi.org/10.1515/bjmg-2016-0002.
- Haque MU, Ferdiousi N, Sajon SR. Anticancer agents derived from plant and dietary sources: a review. Int J Pharmacogn. 2016;32:55–66.
- Sethi VS, Jackson DJ, White DR, Richards F2, Stuart JJ, Muss HB, et al. Pharmacokinetics of vincristine sulfate in adult cancer patients. *Cancer Res.* 1981;41(9 Pt 1):3551–5. [PubMed ID: 7260915].
- Lee CT, Huang YW, Yang CH, Huang KS. Drug delivery systems and combination therapy by using vinca alkaloids. *Curr Top Med Chem.* 2015;**15**(15):1491–500. [PubMed ID: 25877096]. [PubMed Central ID: PM-CPmc4997956]. https://doi.org/10.2174/1568026615666150414120547.
- Shabani H, Zandi M, Ofoghi H, Sanjabi MR, Hoseini Pajooh K. The effect of combining vitamin E and C on the viability improvement of transfected ovine spermatogonial stem cells after cryopreservation and thawing. *Turkish J Vet Anim Sci.* 2017;41(5):648–55. https://doi.org/10.3906/vet-1701-38.
- Li DD, Qin XC, Yang Y, Chu HX, Li RL, Ma LX, et al. Daurinoline suppressed the migration and invasion of chemo-resistant human non-small cell lung cancer cells by reversing EMT and Notch-1 and sensitized the cells to Taxol. *Environ Toxicol Pharmacol.* 2019;66:109–15. [PubMed ID: 30641414]. https://doi.org/10.1016/j.etap.2018.12.005.
- Feizabadi MS, Hernandez MAV, Breslin JB, Akintola ,I. The regulatory effect of Tau protein on polymerization of MCF7 microtubules in vitro. *Biochem Biophys Rep.* 2019;17:151-6. [PubMed ID: 30671547]. [PubMed Central ID: PMCPmc6327910]. https://doi.org/10.1016/j.bbrep.2018.12.010.