



The Effect of Aerobic Exercise and Pistachio Soft Hull Extract on the Expression of the IL-6 and IL-1 β Genes in the Soleus Muscle of Female Rats Fed a High-Fat Diet

Hadi Barshan¹, Mohammad Ali Azarbayjani ^{1,*}, Saleh Rahmati ², Maghsoud Peeri ¹

¹ Central Tehran Branch, Islamic Azad University, Tehran, Iran

² Pardis Branch, Islamic Azad University, Tehran, Iran

* Corresponding author: Central Tehran Branch, Islamic Azad University, Tehran, Iran. Email: m_azarbayjani@iauctb.ac.ir

Received 2023 December 22; Revised 2024 February 18; Accepted 2024 February 20.

Abstract

Background: Obesity is considered the most important public health problem. Previous studies have confirmed that a high-fat diet (HFD) can cause the development of systemic and tissue inflammation due to the increase in pro-inflammatory markers.

Objectives: This study aims to evaluate the effect of aerobic exercise and pistachio soft hull extract (PSHE) on inflammatory gene expression in the soleus muscle of rats fed a HFD.

Methods: In an experimental trial, 30 female Wistar rats weighing between 180 and 200 grams were selected as subjects and randomly divided into five groups: normal diet control, HFD control, HFD aerobic exercise, HFD-PSHE, and HFD aerobic exercise-PSHE. The aerobic exercise program consisted of four weeks of five weekly sessions of running on a treadmill at an intensity of 50 - 60% VO_{2max} . Pistachio soft hull extract at a dose of 60 mg/kg was administered to the rats by gavage for four weeks, five times a week. Forty-eight hours after the last interventions, the rats were anesthetized, and the soleus muscle was removed to determine the expression of the IL-6 and IL-1 β genes.

Results: Induction of HFD significantly decreased IL-6 ($P = 0.001$) and increased IL-1 β ($P = 0.001$) gene expression. IL-1 β gene expression in the aerobic exercise ($P = 0.001$), PSHE ($P = 0.003$), and aerobic exercise-PSHE ($P = 0.001$) groups was significantly lower than in the HFD control group. Aerobic exercise ($P = 0.002$), PSHE ($P = 0.002$), and the combination of aerobic exercise and PSHE ($P = 0.003$) caused a significant increase in IL-6 gene expression compared to the HFD control group.

Conclusions: High-fat diet promotes inflammation in skeletal muscle by increasing the expression of the IL-1 β and IL-6 genes. Aerobic exercise, PSHE, and the combination of aerobic exercise and PSHE can exert their anti-inflammatory effects by reducing the expression of the IL-1 β and IL-6 genes in skeletal muscle tissue.

Keywords: Obesity, Public Health, Inflammation, Anti-inflammatory Effects, Exercise Training

1. Background

Obesity is one of the greatest health problems (1). This disorder is an important risk factor for metabolic diseases associated with premature death (2). One of the main causes of overweight and obesity is the consumption of a high-calorie diet and lack of physical activity (3). It is well known that obesity is associated with many diseases related to low-grade chronic inflammation (4-6). Obesity-induced inflammation causes significant disturbances in skeletal muscle structure and function, including impaired muscle metabolism, insulin signaling, atrophy, and reduced

hypertrophy (7). Feeding with a high-fat diet (HFD) increases the gene expression of IL-1 β and decreases IL-6 mRNA in skeletal muscle (8, 9). A significant increase in the expression of pro-inflammatory cytokines and a decrease in the expression of anti-inflammatory genes in the muscle cause changes in the signaling pathways that disrupt the metabolic functions of the muscle, especially lipid and glucose metabolism (10). For example, a disruption in the skeletal muscle insulin signaling pathway has been reported with a significant increase in the gene expression of pro-inflammatory cytokines in myotubes exposed to palmitic acid (11). Disturbance in the muscle signaling pathway eventually

leads to type 2 diabetes (12). It is well known that aerobic exercise is one of the most effective interventions for the prevention and treatment of insulin resistance. Several studies showed that aerobic exercise decreased the gene expression of skeletal muscle pro-inflammatory cytokines, especially IL-1 β and TNF- α , under obesity conditions. It increased the gene expression of anti-inflammatory cytokines (13, 14). Since the over-expression of inflammatory mediator genes and the reduction of anti-inflammatory mediators inhibit insulin resistance, aerobic exercises are increasingly considered a protective factor for muscle against insulin resistance (15).

On the other hand, medicinal herbs and supplements can reduce inflammation in obesity due to their bioactive compounds (16-18). One of the plants that has various anti-inflammatory compounds is the pistachio soft hull. It contains anthocyanins, flavan-3-ols, proanthocyanidins, flavonols, isoflavones, flavanones, stilbenes, and phenolic acids (19). The presence of these bioactive compounds gives pistachio soft hull anti-inflammatory properties (20). It has been reported that the compounds in pistachio soft hull inhibit IL-1 β gene expression (21-23).

2. Objectives

Aerobic exercise and pistachio soft hull can alter the gene expression of pro-inflammatory and anti-inflammatory mediators in skeletal muscle under conditions of obesity. However, no study has examined the simultaneous effect of these interventions on the expression of IL-6 and IL-1 β . The purpose of the current study is to assess the impact of aerobic exercise and pistachio soft hull extract (PSHE) on IL-6 and IL-1 β in the soleus muscle of female rats fed a HFD.

3. Methods

3.1. Animals

In an experimental trial, 30 female Wistar rats, weighing between 180 and 200 grams, were purchased from the Pasteur Institute and transferred to the Laboratory Center of Azad Islamic University, Tehran Branch. To allow the rats to adapt to the environment and to control confounding factors, the rats were acclimated for two weeks before the start of the protocols (24, 25). The laboratory environment was prepared according to the following standards: Regulation of relative humidity at 50% \pm 10%, regulation of light-dark cycles to 12/12 hours, temperature maintained at 25 \pm 2 degrees Celsius, a ventilation

system with a silent fan, housing 6 rats in a standard cage, and providing free access to tap water and rat chow. The rats were randomly divided into five groups: 1, normal diet control; 2, HFD control; 3, HFD-aerobic exercise; 4, HFD-PSHE; and 5, HFD-aerobic exercise-PSHE.

3.2. High-Fat Diet Induction

The normal diet of the control group consisted of commercial pellet feed (Behparvar Company, Karaj, Iran) for laboratory mice. To induce a HFD, the normal diet was supplemented with 20% palm oil, 1.5% cholesterol, and 0.25% cholic acid (26).

3.3. Aerobic Exercise Program

The aerobic exercise program consisted of four weeks of five weekly sessions of running on a treadmill. One week before the formal training, the rats were acclimated to the treadmill by running at a speed of 9 m/min for 20 minutes. The intensity and duration of aerobic exercise from the first to the last day were as follows: Moderate-intensity (MET) training sessions at 50 – 60% Vo_{2max} , consisting of five sessions per week with a 5-minute warm-up, a 20-minute exercise period, and a 5-minute cool-down. On the first day of training, the speed started at 16 m/min and, according to the program, increased by 2.5 m/min each week. By the last day, after four weeks, the speed reached 26 m/min (27).

3.4. Pistachio Soft Hull Extract Preparation and Induction

The soft pistachio hull was obtained from Rafsanjan and verified by a botanist. After drying in the shade, it was powdered using a mill. A 100 mg sample was immersed in 5 mL of four solvents (70% acetone, 50% ethanol, 50% methanol, and water) for two hours at room temperature on a shaker, completing the extraction process. The samples were then centrifuged at 3000 g for 10 minutes. The prepared extract in pure liquid form was dissolved in distilled water and administered to the rats via gavage at a dose of 60 mg/kg for four weeks, five times a week (28, 29).

Forty-eight hours after the last interventions (aerobic exercise and PSHE) and after twelve hours of fasting, the rats were anesthetized with a combination of ketamine and xylazine, and the soleus muscle was removed to determine the expression of the IL-6 and IL-1 β genes.

3.5. Sacrifice and Collection of Soleus Muscle Tissue

After fasting for 8 – 10 hours, all Wistar rats were sacrificed, and their body weights were measured before

tissue harvesting. Once the animals were fully anesthetized and unconscious, in accordance with the Declaration of Helsinki, the rat soleus muscle tissue was quickly removed and washed with PBS (phosphate-buffered saline). Mucus, blood, and excess material were cleaned from the tissues, which were then placed in 2 mL microtubes. The microtubes were placed in a nitrogen tank and stored in an 80-degree freezer until final testing and cell analysis (30).

3.6. Real-time PCR Analysis

The qPCR method was used to study gene expression in soleus muscle tissue. The β -actin reference gene was used as a control, and the expression of other genes was compared to it. First, primers were designed, and then total RNA was extracted from the tissues and converted into cDNA. The cDNA was then amplified using PCR, and the expression of these genes was analyzed. RNA extraction was carried out manually using Trizol reagent prepared by the Kiazist Company, following the currently accepted standard procedure for the trizol method. cDNA synthesis was performed using Oligo dT primers and according to the instructions of the German Fermentase kit. The primers were designed using Gene Runner version 6.5 (Table 1) (31).

3.7. Statistical Method

The Shapiro-Wilk test was used to determine the normality of the data distribution. After confirming the data were normally distributed, an independent *t*-test was used to test the differences between the healthy control group and the HFD group. A two-way analysis of variance was used to examine the differences between the other groups. Bonferroni's post hoc test was also used to determine the main interaction effect of aerobic exercise and PSHE on IL-1 β and IL-6. A significance level of $P < 0.05$ was considered for all calculations. All analyses were performed with SPSS version 25.

4. Results

Induction of HFD significantly increased IL-1 β gene expression ($P = 0.001$) in the control group compared to the normal diet control group (Figure 1).

IL-1 β gene expression in the aerobic exercise ($P = 0.001$), PSHE ($P = 0.003$), and aerobic exercise-PSHE ($P = 0.001$) groups was significantly lower than in the HFD control group. The interaction of aerobic exercise and PSHE on the expression of the IL-1 β gene in the soleus muscle of rats fed an HFD was significant ($P = 0.001$), indicating that these two interventions enhance each

other's effect in reducing the expression of this gene (Figure 2).

The expression of the IL-6 gene of soleus muscle in the group fed with HFD was significantly lower than the group fed with a normal diet ($P = 0.001$) (Figure 3).

Aerobic exercise ($P = 0.002$), PSHE ($P = 0.002$), and the combination of aerobic exercise- PSHE ($P = 0.003$) caused a significant increase in IL-6 gene expression compared to the HFD control group. The interaction of aerobic exercise and PSHE was significant in the increase of IL-6 gene expression ($P = 0.003$) (Figure 4).

5. Discussion

The first finding of this study showed that feeding with a HFD increased IL-1 β and decreased IL-6 gene expression in the soleus muscle. Eating a HFD causes low-grade chronic inflammation and insulin resistance (32). From a molecular perspective, obesity increases tissue and systemic inflammation due to increased gene expression of pro-inflammatory mediators at the cellular level (33-35). Skeletal muscle, like other tissues, is affected by a HFD. Increased inflammation, oxidative stress, and disturbance in skeletal muscle metabolism with HFD have been reported in rodents (36, 37). Like other tissues such as the liver, brain, and adipose tissue, feeding with a HFD is generally associated with increased gene and protein expression of inflammatory mediators such as IL-1 β and TNF- α in skeletal muscle.

The exact molecular mechanism of increased IL-1 β expression in skeletal muscle following HFD is not well known, but possible mechanisms have been proposed. One suggested mechanism involves the activation of the cytoplasmic nucleotide-binding oligomerization domain-like receptor family (NOD-like) pyrin domain containing 3 (NLRP3) inflammasome (9). Pyrin domain containing 3 is associated with increased inflammation and insulin resistance. Fatty acids in a HFD, especially palmitic acid, activate the TLR4/NF κ B signaling pathway by acting on the TLR4 receptor (38). The activation of NF κ B leads to the activation of NLRP3, which then triggers the expression of the IL-1 β gene (9). Additionally, feeding with a HFD activates caspase-1 and increases IL-1 β gene and protein expression (39).

In the present study, aerobic exercise decreased IL-1 β gene expression. This exercise increases the uptake and oxidation of plasma fatty acids by active muscles, thus reducing the activation of the TLR4/NF κ B pathway and subsequently decreasing IL-1 β mRNA expression. Increased oxidative stress is one of the factors that activate inflammatory mediators in skeletal muscle tissue (8). It seems that in the present study, aerobic

Table 1. The Specific Primers Used in Real-time PCR

Variables	Reverse (5' -> 3')	Forward (5' -> 3')
IL-1 β	CACGGTTTTCTTATGGCTCTG	CTATTCCTAATGCCTTCCCAG
IL-6	TGTCATCAGCCTCCAATTC	ACCAAGACCATCCAACATC
β -actin	GCTTAGGCATAGCACACTAGG C	ACCGTGAAAAGATGACCCAG

Abbreviations: IL-1 β , interleukin 1 beta; IL-6, interleukin 6; β -actin, Beta-actin.




Figure 1. Comparison of IL-1 β gene expression between the normal diet control group and high-fat diet (HFD) control group.* Denote significant differences from the normal diet control group. Data are reported based on scale and standard deviation.

exercise inhibited IL-1 β gene expression by reducing oxidative stress.

On the other hand, PSHE also decreased IL-1 β gene expression. Pistachio soft hull extract bioactive compounds have hypolipidemic properties. It is reported that anthocyanins (40), proanthocyanidins (41), and flavonols (42) exert their hypolipidemic effects by inhibiting the absorption of fats in the small intestine and reducing circulating lipids. As a result, these compounds can decrease the expression of IL-1 β mRNA by reducing circulating fatty acids and subsequently reducing the activation of the TLR4/NF κ B pathway. Therefore, it seems that PSHE reduces IL-1 β gene expression by decreasing circulating lipids and affecting signaling pathways. Additionally, PSHE bioactive compounds, due to their high antioxidant

properties, reduce the oxidative stress caused by HFD and subsequently lower the expression of IL-1 β .

The significant finding of this study was the notable interaction of aerobic exercise and PSHE on the reduction of IL-1 β gene expression. It appears that aerobic exercise and PSHE each reduced fatty acids and oxidative stress through different mechanisms, thereby decreasing IL-1 β gene expression in the soleus muscle.

Another finding of this study showed that HFD decreased IL-6 gene expression in the soleus muscle. IL-6 is an inflammatory cytokine expressed as an adipokine in adipose tissue and released into the circulation, which is associated with tissue damage and the development of many metabolic diseases (43). Since IL-6 is also expressed in skeletal muscle tissue and responds to muscle contractions, it has attracted the attention of researchers as a myokine with anti-inflammatory




Figure 2. Effect of aerobic exercise and PSHE on IL-1 β gene expression of soleus muscle in female Wistar rats. PSHE, pistachio soft hull extract; HFD, high-fat diet. Data represent mean \pm SD (n = 6). Two-way ANOVA, followed by the post hoc test of Bonferroni. * Denotes significant differences to the HFD control group, and † denotes significant interaction of aerobic exercise and PSHE.




Figure 3. Comparison of IL-6 gene expression between the normal diet control group and high-fat diet (HFD) control group. * Denote significant differences from the normal diet control group. Data are reported based on scale and standard deviation.

properties. During muscle contractions, IL-6 expression increases, enters the blood circulation, and enhances lipolysis, uptake, and oxidation of fatty acids by skeletal

muscles (44). It has been reported that skeletal muscle IL-6 expression decreases due to a HFD. Performing aerobic exercises can counteract the inhibitory effect of




Figure 4. Effect of aerobic exercise and PSHE on IL-6 gene expression of soleus muscle in female Wistar rats. PSHE, pistachio soft hull extract; HFD, High-fat diet. Data represent mean \pm SD (n=6). Two-way ANOVA, followed by the post hoc test of Bonferroni. * Denotes significant differences to the HFD control group, and † denotes significant interaction of aerobic exercise and PSHE.

a HFD on IL-6 gene expression and increase IL-6 expression in skeletal muscle (7). In the present study, aerobic exercise increased IL-6 gene expression. It seems that aerobic exercise improves the myokine profile by increasing the uptake and oxidation of fatty acids and stimulating IL-6 gene expression. Muscle contractions caused by aerobic exercises increase the level of cytosolic calcium ions and enhance the expression of IL-6 through the activation of MAPK and calcineurin (45).

In the present study, PSHE also increased the expression of the IL-6 gene in the soleus muscle. The reduction of plasma fatty acids caused by the compounds in PSHE led to a change in the activation pattern of myokine gene expression, especially IL-6. Both aerobic exercise and PSHE were able to amplify each other's effects on skeletal muscle IL-6 gene expression through different molecular mechanisms, indicating the synergistic effect of these two interventions on the expression of this gene.

5.1. Conclusions

A HFD has been implicated in promoting inflammation within skeletal muscle, primarily by elevating the expression of pro-inflammatory genes such as IL-1 β and IL-6. Aerobic exercise, characterized by sustained and rhythmic physical activity, has demonstrated efficacy in counteracting the deleterious effects of HFD-induced inflammation. It effectively downregulates the expression of IL-6 and IL-1 β genes in skeletal muscle tissue. Similarly, PSHE has emerged as a

noteworthy agent in attenuating skeletal muscle inflammation. Furthermore, combining aerobic exercise with PSHE represents an innovative approach, synergistically amplifying their individual benefits.

Acknowledgements

This article is a part of a doctoral thesis in the field of exercise physiology, which was approved by the Exercise Physiology Department of Islamic Azad University, Central Tehran Branch.

Footnotes

Authors' Contribution: Study concept and design, Maghsoud Peeri; analysis and interpretation of data, Saleh R. Ahmadabad; acquisition of data, Hadi Barshan; study supervision, Mohammad Ali Azarbayjani.

Conflict of Interests Statement: The authors declare no conflict of interest.

Data Availability: The dataset presented in the study is available on request from the corresponding author during submission or after publication.

Ethical Approval: This article is part of the results of a doctoral mega-project in 1400, which has been approved by the university's research unit.

Funding/Support: No funding is reported by the authors.

References

- McGown C, Biredinc A, Younossi ZM. Adipose Tissue Endocrine Organ. *Clin Liver Dis*. 2014;**18**(1):41-58. [PubMed ID: [24274864](#)]. <https://doi.org/10.1016/j.cld.2013.09.012>.
- Chong B, Kong G, Shankar K, Chew HSJ, Lin C, Goh R, et al. The global syndemic of metabolic diseases in the young adult population: A consortium of trends and projections from the Global Burden of Disease 2000-2019. *Metab*. 2023;**141**:155402. [PubMed ID: [36717058](#)]. <https://doi.org/10.1016/j.metabol.2023.155402>.
- Moschonis G, Trakman GL. Overweight and Obesity: The Interplay of Eating Habits and Physical Activity. *Nutr*. 2023;**15**(13). [PubMed ID: [37447222](#)]. [PubMed Central ID: [PMC10343820](#)]. <https://doi.org/10.3390/nu15132896>.
- Khanna D, Khanna S, Khanna P, Kahar P, Patel BM. Obesity: A Chronic Low-Grade Inflammation and Its Markers. *Cureus*. 2022. <https://doi.org/10.7759/cureus.22711>.
- Milano W, Carizzone F, Foia M, Marchese M, Milano M, Saetta B, et al. Obesity and Its Multiple Clinical Implications between Inflammatory States and Gut Microbiotic Alterations. *Dis*. 2022;**11**(1). [PubMed ID: [36648872](#)]. [PubMed Central ID: [PMC9844347](#)]. <https://doi.org/10.3390/diseases11010007>.
- Azeez S, Jafar S, Aziziaram Z, Fang L, Mawlood A, Ercisli M. Insulin-producing cells from bone marrow stem cells versus injectable insulin for the treatment of rats with type I diabetes. *Cell, Mol Biomedical Rep*. 2021;**1**(1):42-51. <https://doi.org/10.55705/cmbr.2021.138888.1006>.
- Sinha I, Sakthivel D, Varon DE. Systemic Regulators of Skeletal Regeneration in Obesity. *Front Endocrinol (Lausanne)*. 2017;**8**:29. [PubMed ID: [28261159](#)]. [PubMed Central ID: [PMC5311070](#)]. <https://doi.org/10.3389/fendo.2017.00029>.
- Andrich DE, Melbouci L, Ou Y, Auclair N, Mercier J, Grenier JC, et al. A Short-Term High-Fat Diet Alters Glutathione Levels and IL-6 Gene Expression in Oxidative Skeletal Muscles of Young Rats. *Front Physiol*. 2019;**10**:372. [PubMed ID: [31024337](#)]. [PubMed Central ID: [PMC6468044](#)]. <https://doi.org/10.3389/fphys.2019.00372>.
- Jorquera G, Meneses-Valdes R, Rosales-Soto G, Valladares-Ide D, Campos C, Silva-Monasterio M, et al. High extracellular ATP levels released through pannexin-1 channels mediate inflammation and insulin resistance in skeletal muscle fibres of diet-induced obese mice. *Diabetologia*. 2021;**64**(6):1389-401. [PubMed ID: [33710396](#)]. <https://doi.org/10.1007/s00125-021-05418-2>.
- Wu H, Ballantyne CM. Skeletal muscle inflammation and insulin resistance in obesity. *J Clin Invest*. 2017;**127**(1):43-54. [PubMed ID: [28045398](#)]. [PubMed Central ID: [PMC5199705](#)]. <https://doi.org/10.1172/JCI88880>.
- Varma V, Yao-Borengasser A, Rasouli N, Nolen GT, Phanavanh B, Starks T, et al. Muscle inflammatory response and insulin resistance: synergistic interaction between macrophages and fatty acids leads to impaired insulin action. *Am J Physiol Endocrinol Metab*. 2009;**296**(6):E1300-10. [PubMed ID: [19336660](#)]. [PubMed Central ID: [PMC2692398](#)]. <https://doi.org/10.1152/ajpendo.90885.2008>.
- Wang Y, Wang PY, Qin LQ, Davaasambuu G, Kaneko T, Xu J, et al. The development of diabetes mellitus in Wistar rats kept on a high-fat/low-carbohydrate diet for long periods. *Endocrine*. 2003;**22**(2):85-92. [PubMed ID: [14665711](#)]. <https://doi.org/10.1385/endo.22.2:85>.
- Gielen S, Adams V, Mobius-Winkler S, Linke A, Erbs S, Yu J, et al. Anti-inflammatory effects of exercise training in the skeletal muscle of patients with chronic heart failure. *J Am Coll Cardiol*. 2003;**42**(5):861-8. [PubMed ID: [12957433](#)]. [https://doi.org/10.1016/s0735-1097\(03\)00848-9](https://doi.org/10.1016/s0735-1097(03)00848-9).
- Li N, Shi H, Guo Q, Gan Y, Zhang Y, Jia J, et al. Aerobic Exercise Prevents Chronic Inflammation and Insulin Resistance in Skeletal Muscle of High-Fat Diet Mice. *Nutr*. 2022;**14**(18). [PubMed ID: [36145106](#)]. [PubMed Central ID: [PMC9503887](#)]. <https://doi.org/10.3390/nu14183730>.
- Hoffmann C, Weigert C. Skeletal Muscle as an Endocrine Organ: The Role of Myokines in Exercise Adaptations. *Cold Spring Harb Perspect Med*. 2017;**7**(11). [PubMed ID: [28389517](#)]. [PubMed Central ID: [PMC5666622](#)]. <https://doi.org/10.1101/cshperspect.a029793>.
- Karati D, Varghese R, Mahadik KR, Sharma R, Kumar D. Plant Bioactives in the Treatment of Inflammation of Skeletal Muscles: A Molecular Perspective. *Evid Based Complement Alternat Med*. 2022;**2022**:4295802. [PubMed ID: [35911155](#)]. [PubMed Central ID: [PMC9328972](#)]. <https://doi.org/10.1155/2022/4295802>.
- Aziziaram Z, Bilal I, Zhong Y, Mahmud A, Roshandel MR. Protective effects of curcumin against naproxen-induced mitochondrial dysfunction in rat kidney tissue. *Cell, Mol Biomedical Rep*. 2021;**1**(1):23-32. <https://doi.org/10.55705/cmbr.2021.138879.1001>.
- Park Y, Aminizadeh S, Lee J, Zarezadehmehrzi A, Najafipour H, Amiri-Deh Ahmadi M, et al. MitoQ supplementation improves oxygen uptake kinetic by reduced reactive oxygen species levels and altered expression of miR-155 and miR-181b. *FASEB J*. 2022;**36**(51). <https://doi.org/10.1096/fasebj.2022.36.51.R6226>.
- Arjeh E, Akhavan H, Barzegar M, Carbonell-Barrachina AA. Bio-active compounds and functional properties of pistachio hull: A review. *Trends Food Sci Tech*. 2020;**97**:55-64. <https://doi.org/10.1016/j.tifs.2019.12.031>.
- Grace MH, Esposito D, Timmers MA, Xiong J, Yousef G, Komarnytsky S, et al. Chemical composition, antioxidant and anti-inflammatory properties of pistachio hull extracts. *Food Chem*. 2016;**210**:85-95. [PubMed ID: [27211624](#)]. <https://doi.org/10.1016/j.foodchem.2016.04.088>.
- Blonska M, Czuba ZP, Krol W. Effect of flavone derivatives on interleukin-1beta (IL-1beta) mRNA expression and IL-1beta protein synthesis in stimulated RAW 264.7 macrophages. *Scand J Immunol*. 2003;**57**(2):162-6. [PubMed ID: [12588662](#)]. <https://doi.org/10.1046/j.1365-3083.2003.01213.x>.
- Kowalski J, Samojedny A, Paul M, Pietsz G, Wilczok T. Effect of apigenin, kaempferol and resveratrol on the expression of interleukin-1beta and tumor necrosis factor-alpha genes in J774.2 macrophages. *Pharmacol Rep*. 2005;**57**(3):390-4. [PubMed ID: [15985724](#)].
- Wongwichai T, Teeyakasem P, Pruksakorn D, Kongtawelert P, Pothacharoen P. Anthocyanins and metabolites from purple rice inhibit IL-1beta-induced matrix metalloproteinases expression in human articular chondrocytes through the NF-kappaB and ERK/MAPK pathway. *Biomed Pharmacother*. 2019;**112**:108610. [PubMed ID: [30797145](#)]. <https://doi.org/10.1016/j.biopha.2019.108610>.
- Al-Saedi HF, Panahi Y, Ghanimi HA, Abdolmaleki A, Asadi A. Enhancement of nerve regeneration with nimodipine treatment after sciatic nerve injury. *Fundam Clin Pharmacol*. 2023;**37**(1):107-15. [PubMed ID: [35989463](#)]. <https://doi.org/10.1111/fcp.12827>.
- Tamjid M, Abdolmaleki A, Mahmoudi F, Mirzaee S. Neuroprotective Effects of Fe3O4 Nanoparticles Coated with Omega-3 as a Novel Drug for Recovery of Sciatic Nerve Injury in Rats. *Gene, Cell Tissue*. 2022;**10**(2). <https://doi.org/10.5812/gct-124110>.
- Od-Ek P, Deenin W, Malakul W, Phoungpetchara I, Tunsophon S. Anti-obesity effect of Carica papaya in high-fat diet fed rats. *Biomed Rep*. 2020;**13**(4):30. [PubMed ID: [32802327](#)]. [PubMed Central ID: [PMC7412661](#)]. <https://doi.org/10.3892/br.2020.1337>.
- Nikbin S, Tajik A, Allahyari P, Matin G, Hoseini Roote SS, Barati E, et al. Aerobic exercise and eugenol supplementation ameliorated liver injury induced by chlorpyrifos via modulation acetylcholinesterase

- activation and antioxidant defense. *Environ Toxicol.* 2020;**35**(7):783-93. [PubMed ID: 32096903]. <https://doi.org/10.1002/tox.22913>.
28. Azadadel S, Hanachi P, Saboori A. [Investigation on antioxidant activity of pistachio (*Pistacia vera* L.) skin extraction]. *J Plant Res.* 2018;**30**(4):722-31. Persian.
 29. Kazemipour M, Mateen Homaie H, Farzanegi P. [The effect of aerobic exercise with pistachio skin extract on the expression of Bax, caspase-3 and Bcl-2 in heart tissue of fat-fed rats]. *Feyz Med Sci J.* 2022;**26**(5):521-9. Persian. <https://doi.org/10.48307/fmsj.2022.26.5.521>.
 30. Sobhani V, Rahmati-Ahmadabad S, Ghanbari-Niaki A, Shirvani H. Effects of endurance training and herb supplementation on tissue nesfatin-1/nucleobindin-2 and ghrelin mRNA expression. *Int J Appl Exercise Physiol.* 2017;**6**(1):71-84. <https://doi.org/10.22631/ijaep.v6i1.118>.
 31. Ghanbari-Niaki A, Rahmati S, Ansari-Pirsaraei Z. Effect of aerobic exercise on tissue nesfatin-1/nucleobindin-2 mRNA, plasma nesfatin-1 and high-density lipoprotein concentration in female rats. *Iran JHealth Physical Activity.* 2013;**4**:1-7.
 32. Kahn SE, Hull RL, Utzschneider KM. Mechanisms linking obesity to insulin resistance and type 2 diabetes. *Natur.* 2006;**444**(7121):840-6. [PubMed ID: 17167471]. <https://doi.org/10.1038/nature05482>.
 33. De Souza CT, Araujo EP, Bordin S, Ashimine R, Zollner RL, Boschero AC, et al. Consumption of a fat-rich diet activates a proinflammatory response and induces insulin resistance in the hypothalamus. *Endocrinol.* 2005;**146**(10):4192-9. [PubMed ID: 16002529]. <https://doi.org/10.1210/en.2004-1520>.
 34. Lagathu C, Yvan-Charvet L, Bastard JP, Maachi M, Quignard-Boulangé A, Capeau J, et al. Long-term treatment with interleukin-1 β induces insulin resistance in murine and human adipocytes. *Diabetologia.* 2006;**49**(9):2162-73. [PubMed ID: 16865359]. <https://doi.org/10.1007/s00125-006-0335-z>.
 35. Wieser V, Moschen AR, Tilg H. Inflammation, cytokines and insulin resistance: a clinical perspective. *Arch Immunol Ther Exp (Warsz).* 2013;**61**(2):119-25. [PubMed ID: 23307037]. <https://doi.org/10.1007/s00005-012-0210-1>.
 36. Asghar A, Akhtar T, Batool T, Khawar MB, Nadeem S, Mehmood R, et al. High-fat diet-induced splenic, hepatic, and skeletal muscle architecture damage: cellular and molecular players. *Mol Cell Biochem.* 2021;**476**(10):3671-9. [PubMed ID: 34050900]. <https://doi.org/10.1007/s11010-021-04190-6>.
 37. Pinho RA, Sepa-Kishi DM, Bikopoulos G, Wu MV, Uthayakumar A, Mohasses A, et al. High-fat diet induces skeletal muscle oxidative stress in a fiber type-dependent manner in rats. *Free Radic Biol Med.* 2017;**110**:381-9. [PubMed ID: 28690197]. <https://doi.org/10.1016/j.freeradbiomed.2017.07.005>.
 38. Korbecki J, Bajdak-Rusinek K. The effect of palmitic acid on inflammatory response in macrophages: an overview of molecular mechanisms. *Inflamm Res.* 2019;**68**(11):915-32. [PubMed ID: 31363792]. [PubMed Central ID: PMC6813288]. <https://doi.org/10.1007/s00011-019-01273-5>.
 39. Boucher D, Chan A, Ross C, Schroder K. Quantifying Caspase-1 Activity in Murine Macrophages. *Methods Mol Biol.* 2018;**1725**:163-76. [PubMed ID: 29322417]. https://doi.org/10.1007/978-1-4939-7568-6_14.
 40. Oteiza PI, Cremonini E, Fraga CG. Anthocyanin actions at the gastrointestinal tract: Relevance to their health benefits. *Mol Aspects Med.* 2023;**89**:101156. [PubMed ID: 36379746]. <https://doi.org/10.1016/j.mam.2022.101156>.
 41. Nie Y, Sturzenbaum SR. Proanthocyanidins of Natural Origin: Molecular Mechanisms and Implications for Lipid Disorder and Aging-Associated Diseases. *Adv Nutr.* 2019;**10**(3):464-78. [PubMed ID: 30926997]. [PubMed Central ID: PMC6520035]. <https://doi.org/10.1093/advances/nmy118>.
 42. Mahboob A, Samuel SM, Mohamed A, Wani MY, Ghorbel S, Miled N, et al. Role of flavonoids in controlling obesity: molecular targets and mechanisms. *Front Nutr.* 2023;**10**:1177897. [PubMed ID: 37252233]. [PubMed Central ID: PMC10213274]. <https://doi.org/10.3389/fnut.2023.1177897>.
 43. Upadhyaya S, Kadamkode V, Mahammed R, Doraiswami C, Banerjee G. Adiponectin and IL-6: Mediators of inflammation in progression of healthy to type 2 diabetes in Indian population. *Adipocyte.* 2014;**3**(1):39-45. [PubMed ID: 24575367]. [PubMed Central ID: PMC3917930]. <https://doi.org/10.4161/adip.26553>.
 44. Nara H, Watanabe R. Anti-Inflammatory Effect of Muscle-Derived Interleukin-6 and Its Involvement in Lipid Metabolism. *Int J Mol Sci.* 2021;**22**(18). [PubMed ID: 34576053]. [PubMed Central ID: PMC8471880]. <https://doi.org/10.3390/ijms22189889>.
 45. Pedersen BK, Febbraio MA. Muscle as an endocrine organ: focus on muscle-derived interleukin-6. *Physiol Rev.* 2008;**88**(4):1379-406. [PubMed ID: 18923185]. <https://doi.org/10.1152/physrev.90100.2007>.