

Unusual Prevalence of c559T > C Mutation in Patients with Factor XIII deficiency in Southeast of Iran

Ebrahim Miri-Moghaddam^{1,2}; Yasaman Garmie³; Majid Naderi^{1,*}

¹Genetics of Non-Communicable Disease Research Center, Zahedan University of Medical Sciences, Zahedan, IR Iran

²Department of Genetics, Zahedan University of Medical Sciences, Zahedan, IR Iran

³Department of Biology, Faculty of Science, Sistan and Balouchestan University, Zahedan, IR Iran

*Corresponding author: Majid Naderi, Genetics of Non-Communicable Disease Research Center, Ali-Asghar Hospital, Azadi Ave., Zahedan, IR Iran. Tel: +98-5413414567, Fax: +98-5413218998, E-mail: majid_naderi2000@yahoo.com

Received: February 1, 2014; Revised: February 16, 2014; Accepted: February 18, 2014

Background: Congenital factor XIII (FXIII) deficiency is a rare severe autosomal recessive bleeding disorder.

Objectives: The aim of the study was to determine the c559T > C FXIII genotype frequency in patients with FXIII hemophilia who lived in Sistan and Balouchestan province in southeast of Iran.

Patients and Methods: We determined the genotype of 180 patients with Factor XIII hemophilia by tetra-primer amplification refractory mutation system-polymerase chain reaction (T-ARMS-PCR).

Results: The frequency of C559T > C was 96.6% and T-ARMS-PCR was an efficient tool for detecting this genotype. Moreover, the result showed that C559T > C mutation was a risk factor for FXIII deficiency in a sample of Iranian population.

Conclusions: Due to highest prevalence of FXIII deficiency (70 times higher than the global average), FXIII deficiency must be considered as the most underdiagnosed bleeding disorder in this area. Hence, proper screening systems alongside common screening tests to identify carriers seem necessary. Due to the limited mutations, T-ARMS-PCR with sound speed, accuracy, sensitivity, and cheapness can be used to screen for these mutations.

Keywords: Mutation, Factor XIII; Deficiency; Iran; Genotype

1. Background

Blood coagulation disorders are results of deficiency in coagulation factors activity. Coagulation factor XIII (FXIII) has an essential role at the end of coagulation cascade and the stabilization of fibrin clot. FXIII is a zymogen and its active form is a transglutaminase which catalyzes an acyl transfer reaction and cross-links between two polypeptide chains through a γ -glutamyl lysine link (1). FXIII is heterotetramer (A₂B₂) in plasma consisting of two catalytic A subunits and two B subunits; the B subunit acts as a carrier that protects and fixates the A subunits (1). The FXIII-A subunit is encoded by a gene on chromosome 6p24-25 as a protein consisting of 731 amino acid (2). It comprises five domains including the activation peptide (residues of 1-37), β -sandwich (residues of 38-183), central domain or catalytic core region (residues 184-515), and β -barrel 1 and 2 (residues 516-627 and residues 628-731, respectively) (3). The FXIII-B subunit is encoded by a gene located on chromosome 1q31-32.1. This part spans 28 kb and comprises 12 exons and 11 introns (4). FXIII deficiency is a rare bleeding disorder; its frequency is one to three per million populations and is transmitted in an autosomal recessive manner. It is usually more prevalent in

areas with high rate of consanguineous marriages (1, 5). FXIII deficiency is associated with umbilical cord bleeding, prolonged wound healing, recurrent spontaneous miscarriage, and spontaneous intracranial hemorrhage (6). After FVIII deficiency, FXIII is the most common bleeding disorder in Sistan and Balouchestan Province in the southeast of Iran (7). Up to this date, 133 different mutations were reported in FXIII gene in which 114 occurred in FXIII-A subunit and only 19 mutations occurred in FXIII-B subunit. Missense mutations are the most prevalent mutations in FXIII gene (1, 6, 8, 9). Previous studies reported few mutations in patients with FXIII hemophilia in Iranian population and Arg77His and Trp187Arg were the most common mutations (10).

2. Objectives

The aim of this study was to determine the c559T > C FXIII genotype frequency in patients with FXIII hemophilia by T-ARMS-PCR.

3. Patients and Methods

This study was conducted on 180 patients with FXIII hemophilia who were referred to Ali Asghar Hospital with

Implication for health policy/practices/research/medical education:

Careful screening system alongside common screening tests are necessary due to the limited mutations and high frequency in FXIII gene deficiency in this area.

Copyright © 2014, Zahedan University of Medical Sciences. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

clinical symptoms of coagulation disorder and had normal results for prothrombin time (PT), Partial thromboplastin time (PTT), bleeding time (BT), fibrinogen level, and platelet count tests. The most common screening tests for FXIII are based on solubility of fibrin clot in solutions (2% acetic acid or 1% monochloroacetic acid or 5 Mol urea). In abnormal cases, FXIII level was measured by quantitative factor assays and genotyping was performed. This study was approved by the Ethics Committee of Tehran University of Medical Sciences and an informed consent was obtained from each participant. Genomic DNA was extracted from peripheral blood (11) and c559T > C mutation was detected by considering that this mutation was the most common mutation reported by T-ARMS-PCR (9, 12-15). We designed four primers to determine the c559T > C mutation of FXIII. We used, two external primers (forward outer (FO): 5'-TCTTCTCTTCTCATAGGTCGCTACC-3', reverse outer (RO): 5'-CACCACCA-CACCCATCTAATTTTAA-3') and two internal allele-specific primers (forward inner (FI) [C allele]: 5'-AACAGACACGTACATTCTTCAATCATC-3', reverse inner (RI) [T allele]: 5'-AATGCTGCCTCTTACCTTACACAA-3'). The products size were 216 bp for C allele, 286 bp for T allele, and 448 bp for the two outer primers (control band), as is schematically shown in Figure 1.


4. Results

We evaluated 180 patients with hemophilia, 174 (96.6%) patients had substitution of C to T and change of Tryptophan to Arginine in the form of homozygote. The T-ARMS-PCR method was effectively applied for genotyping C559T > C mutation of the FXIII. The genotypes determined by this method is shown in Figure 2, and conformities sequencing are illustrated in Figure 3.

5. Discussion


FXIII deficiency is a rare autosomal recessive disorder with a higher prevalence rate in Sistan and Balouchestan province than the worldwide rate. This high prevalence might be due to the consanguineous marriage as more 70% of marriages are consanguineous in this province (10). In this area, the number of patients diagnosed were significantly increased from 45 in 2006 to 205 cases in 2012, (16). Sistan and Balouchestan province with 2,700,000 population had about 200 FXIII deficient known cases (1 per 13,500 population), which is 70 times higher than the global average. The number of FXIII deficient cases were few in the other parts of the world, for example, in UK it was 26 cases (7) and 72 cases were recognized in a vast study carried out in European Coagulation Disorders Research Center (1996) (17). Frequency ratio of factor XIII deficiency to blood coagulation disorders was reported in 62% Caucasians, 6% in African-America, 19% in Latinos, 3% in Asian (7) whereas Factor XIII deficiency is the second most common blood coagulation disorders

Figure 1. Schematic Illustration of the Tetra Amplification Refractory Mutation System-Polymerase Chain Reaction (T-ARMS-PCR) Assay




The forward outer (FO) and the reverse outer (RO) primers give a 448 bp product (control band). FO and Reverse inner (RI) Primers amplify the T allele, producing a 286 bp product, and primers forward inner (FI) and RO produce a 216 bp product for the C allele.

Figure 2. Electrophoresis Pattern of Tetra Amplification Refractory Mutation System-Polymerase Chain Reaction for Finding FXIII C559T > C mutation



M = DNA marker.

Figure 3. Sequencing Results of FXIII Genomic DNA



A) Homozygous CC genotype; B) Homozygous TT and heterozygous.

in southeast of Iran (10). In the present study, we showed that c.559T>C mutation was the most common mutation (more than 95%) in FXIII gene in Sistan & Balouchestan province. T-ARMS-PCR is a quick and easy technique for determination of the mutation (18). This method was easily applied for genotyping the Trp187Arg in FXIII gene and the determined genotypes were confirmed by the sequencing genotypes. The parents' DNA showed heterozygous pattern. One of the advantages of this method is that efficiently amplifies both wild type and the mutated alleles with a control fragment in a single PCR tube. A large number of mutations have been reported according to different databases for FXIII gene (<http://www.hgmd.cf.ac.uk> and <http://www.f13-database.de>) in different parts of the world. Anwar et al. reported four mutations codon 295; G → A at position -246 upstream of exon 1; T → C and C → T at positions -23 and -24, respectively, in intron nine in patients with FXIII hemophilia who originated from the North of Pakistan, the neighboring country of this

province (19). The common mutations in five different subunits of FXIII-A gene are shown in Table 1. Previous studies reported Arg77His and Trp187Arg as the most common mutations (10).

Considering the high rate of consanguineous marriages, the great homogeneity in the population and mutations was seen in this area. The sequencing change in this mutation is TGG → CGG and is related to a tryptophan to arginine substitution at residue 187 (20). Trp187 in FXIII is conserved in all known transglutaminases (21). In the FXIII subunit, residue of Trp187 is located in the direction to the surface of the protein between the catalytic core and the b-sandwich domain (22) and substitution of this residue with arginine would increase steric clashes between the Arginine side chain atoms and the contiguous atoms from the neighboring residues (20). In sum, C559T > C mutation was a risk factor for predisposition to FXIII deficiency in this part of Iran. It seemed that this province had the highest prevalence rate of FXIII deficiency in the world. With respect to the high incidence of

Table 1. Common Mutations Found in five Different Subunit of FXIII-A Gene (3)

Type of Mutations and Region	Beta Sandwich	Core	Beta Barrel 1	Beta Barrel 2	Activated Peptide
Missense	Asn60Lys, Arg77Cys	Pro186Leu, Trp187Arg, Gly210Arg, Gly215Arg, Leu235Arg, Met242Arg, Arg252Ile, Arg260Cys, Arg260His, Arg260Leu, Gly262Glu, Ser263Phe, Trp283Cys, Pro289Arg, Ser295Arg, Val316Phe, Ala318Val, Arg326Gln, Leu354Pro, Ala378Pro, Arg382Ser, Ala394Val, Thr398Asn, Arg408Gln, Ser413Trp, Val414Phe, Gly420Ser, Leu498Pro, Gly501Arg	Arg540Gln	Gly592Ser, Arg611His	
	Arg77His, Arg78Cys		Asn541Lys	Leu660Pro, Leu667Pro	
	Met159Arg, Tyr167Cys		Gly562Arg	Asp668Gly, Arg703Trp	
				Ser708Asn, His716Arg	
Non sense	c.210T>G, c.514C>T	c.979C>T, c.1201C>T, c.1171G>T, c.1326C>A	-	c.1984C>T, c.1994G>A	
	c.523 C>T			c.2075G>A	
Insertions/deletions	(-7 to -20) insTT	c.599-600delAA, c.689delA, Exon 5 del, ins T866, c.709delG	c.1652del10bp	c.2002-2003delCT	33bpdel
	c.291-296insTCGTCC	c.869insC, c.1030-1032delAAT, c.1201insC, 127067del33		c.2066del C, Exon 15 del	c.27delT
	c.249-261del13bp	c.1392-1395delAATT, c.1405-1408delCAAA, Gross Deletion(> 100 kb)			
	c.397insG				
	c.499-512del14bp				
Splice site	c.131-132delAG	IVS5 -1 G>A, IVS7+1 G>A, c.1111T>G, IVS10+1G, c.2045G>A			
	c.319GNT, IVS3+6 T>C, IVS3+5 G>A				

FXIII gene deficiency, an appropriate screening system to identify the carriers is essential. Due to limited mutations in this area, T-ARMS-PCR with a sound speed, accuracy, sensitivity, as well as cheapness can be used alongside the common screening tests.

Acknowledgements

The author expressed their thanks to the patients who participated in this study and their families.

Authors' Contribution

All authors contributed to this work. Ebrahim Miri-Moghaddam contributed to design, setup, and preparing the manuscript; Majid Naderi contributed to the diagnosing and referring patients with hemophilia; Yasaman Garmei contributed to the preparation of the manuscript and genotyping the patients.

Financial Disclosure

The authors indicated that they had no financial relationships relevant to this article to disclose.

Funding/Support

This study was financially supported by vice president for research and technology, Tehran University of Medical Sciences, by number, 91-4-31-19213.

References

- Muszbek L, Bagoly Z, Cairo A, Peyvandi F. Novel aspects of factor XIII deficiency. *Curr Opin Hematol*. 2011;**18**(5):366-72.
- Board PG, Webb GC, McKee J, Ichinose A. Localization of the coagulation factor XIII A subunit gene (F13A) to chromosome bands 6p24---p25. *Cytogenet Cell Genet*. 1988;**48**(1):25-7.
- Eshghi P, Cohan N, Naderi M, Karimi M. Factor XIII deficiency: a review of literature. *IJBC*. 2012;**4**(2):85-91.
- Ichinose A, McMullen BA, Fujikawa K, Davie EW. Amino acid sequence of the b subunit of human factor XIII, a protein composed of ten repetitive segments. *Biochemistry*. 1986;**25**(16):4633-8.
- Board PG, Losowsky MS, Miloszewski KJ. Factor XIII: inherited and acquired deficiency. *Blood Rev*. 1993;**7**(4):229-42.
- Cummins RO, Chamberlain D, Hazinski MF, Nadkarni V, Kloeck W, Kramer E, et al. Recommended guidelines for reviewing, reporting, and conducting research on in-hospital resuscitation: the in-hospital 'Utstein style'. A statement for healthcare professionals from the American Heart Association, the European Resuscitation Council, the Heart and Stroke Foundation of Canada, the Australian Resuscitation Council, and the Resuscitation Councils of Southern Africa. *Resuscitation*. 1997;**34**(2):151-83.
- Peyvandi F, Duga S, Akhavan S, Mannucci PM. Rare coagulation deficiencies. *Haemophilia*. 2002;**8**(3):308-21.
- Schroeder V, Kohler HP. Factor XIII deficiency: an update. *Semin Thromb Hemost*. 2013;**39**(6):632-41.
- Karimi M, Bereczky Z, Cohan N, Muszbek L. Factor XIII Deficiency. *Semin Thromb Hemost*. 2009;**35**(4):426-38.
- Eshghi P, Abolghasemi H, Sanei-Moghaddam E, Anwar R, Jazebi M, Amid A, et al. Factor XIII deficiency in south-east Iran. *Haemophilia*. 2004;**10**(5):470-2.
- Sambrook J, Russell David W. *Molecular cloning: a laboratory manual. Vol. 3.*: Cold spring harbor laboratory press; 1989.
- Eshghi P, Mahjour SB, Naderi M, Dehbozorgian J, Karimi M. Long-term prophylaxis in patients with factor XIII deficiency complicated by intracranial haemorrhage in Iran. *Haemophilia*. 2010;**16**(2):383-5.
- Lak M, Peyvandi F, Ali Sharifian A, Karimi K, Mannucci PM. Pattern of symptoms in 93 Iranian patients with severe factor XIII deficiency. *J Thromb Haemost*. 2003;**1**(8):1852-3.
- Naderi M, Zarei T, Haghpanah S, Eshghi P, Miri-Moghaddam E, Karimi M. Intracranial hemorrhage pattern in the patients with factor XIII deficiency. *Ann Hematol*. 2013.
- Peyvandi F, Tagliabue L, Menegatti M, Karimi M, Komaromi I, Katona E, et al. Phenotype-genotype characterization of 10 families with severe a subunit factor XIII deficiency. *Hum Mutat*. 2004;**23**(1):98.
- Naderi M, Imani M, Eshghi P, Dorgalaleh A, Tabibian S, Alizadeh S, et al. Factor XIII deficiency in Sistan and Baluchistan province. *Sci J Blood Transfus Organ*. 2013;**10**(3):282-8.
- Seitz R, Duckert F, Lopaciuk S, Muszbek L, Rodeghiero F, Seligsohn U. ETRO Working Party on Factor XIII questionnaire on congenital factor XIII deficiency in Europe: status and perspectives. Study Group. *Semin Thromb Hemost*. 1996;**22**(5):415-8.
- Ye S, Humphries S, Green F. Allele specific amplification by tetra-primer PCR. *Nucleic Acids Res*. 1992;**20**(5):1152.
- Anwar R, Gallivan L, Miloszewski KJ, Markham AF. Factor XIII deficiency causing mutation, Ser295Arg, in exon 7 of the factor XIII gene. *Thromb Haemost*. 2000;**84**(4):591-4.
- Trinh CH, Sh Elsayed W, Eshghi P, Miri-Moghaddam E, Zadeh-Vakili A, Markham AF, et al. Molecular analysis of sixteen unrelated factor XIII deficient families from south-east of Iran. *Br J Haematol*. 2008;**140**(5):581-4.
- Altschul SF, Madden TL, Schaffer AA, Zhang J, Zhang Z, Miller W, et al. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res*. 1997;**25**(17):3389-402.
- Yee VC, Pedersen LC, Le Trong I, Bishop PD, Stenkamp RE, Teller DC. Three-dimensional structure of a transglutaminase: human blood coagulation factor XIII. *Proc Natl Acad Sci U S A*. 1994;**91**(15):7296-300.