



Study of Distribution of Biofilm Producing Genes in *Staphylococcus aureus* Isolated from Local Cheese Samples in Maragheh City

Bahareh Falaki,¹ and Saman Mahdavi^{2,*}

¹Department of Microbiology, Urmia Branch, Islamic Azad University, Urmia, Iran

²Department of Microbiology, Maragheh Branch, Islamic Azad University, Maragheh, Iran

*Corresponding author: Saman Mahdavi, Department of Microbiology, Maragheh Branch, Islamic Azad University, Maragheh, Iran. Tel: +98-9144150454, E-mail: s.mahdavi@iau-maragheh.ac.ir

Received 2017 August 26; Revised 2017 September 25; Accepted 2017 October 22.

Abstract

Background: *Staphylococcus aureus* is one of the main causes of food poisoning in the world. This pathogen has the ability to create biofilms that can lead to food contamination. The presence of biofilm producing genes in this bacterium plays a very important role in its virulence and pathogenicity and even prevents the penetration of antibiotics in pathogenicity time. The aim of this study was to investigate the distribution of biofilm producing genes in *Staphylococcus aureus* isolated from local cheese samples in Maragheh city.

Methods: 20 *Staphylococcus aureus* isolates from local cheese samples in Maragheh city that had been identified phenotypically by biochemical tests for confirmatory diagnosis were identified by multiplication of species-specific thermonuclease gene (*nuc*) by PCR method. Specific primers for each gene were used for detection of biofilm producing genes *icaA*, *icaD*, *fmbA*, and *clfB* by PCR method.

Results: 17 isolates (85%) were identified as *Staphylococcus aureus* by multiplying of thermonuclease (*nuc*) species-specific gene. The frequency of each gene separately in tested isolates was *icaA* (76.47%), *icaD* (58.82%), *fmbA* (64.7%), and *clfB* (76.47%). 5 isolates (29.41%) contained all of the tested genes simultaneously. The most frequency of tested genes combinationally was associated with *icaA*, *icaD*, and *clfB* genes (41.17%). None of the *Staphylococcus aureus* samples was negative regarding the presence of four tested genes.

Conclusions: Molecular confirmation of 85% of *Staphylococcus aureus* strains by PCR method represents more accuracy of this method compared to biochemical methods. High frequency of biofilm producing genes in tested isolates indicates high potential of this bacterium in biofilm forming. The difference in the dispersion of effective genes in biofilm forming among different *Staphylococcus aureus* strains probably originates in geographical differences and differences in ecological origin of isolated bacteria.

Keywords: *Staphylococcus aureus*, Local Cheese, Biofilm, PCR

1. Background

Staphylococcus aureus is one of the most important bacterial species that cause food poisoning throughout the world. The poisoning caused by this bacterium is one of the most widespread types of food poisoning and is among the three first-degree poisonings (1). Different investigations suggest that 1% - 5% of infections caused by food materials are related to the consumption of milk and dairy products and 53% of these infections result from the use of contaminated cheese (2). One factor that plays an influential role in the pathogenicity of this bacterium is its capacity to form biofilms. Biofilm increases the resistance of the bacterium against the prescribed antibiotics and elevates the protective mechanisms of the host. This elevated resistance of the bacterium against the mentioned factors plays a key role in the development of nosocomial infections (3). Infections caused by *Staphylococcus aureus* are recognized

as chronic and treatment-resistant infections. In different studies conducted in body and lab environments, it has been proven that *Staphylococcus aureus* has the capacity to adhere to and penetrate into the epithelial cells of bovine mammary glands (4). Biofilms are structures comprising collections of bacteria; they are surrounded by a polymeric matrix produced by the bacteria themselves and therefore, they can adhere to different surfaces (5). Adhesion of microbial cells to surfaces and their collection to form multilayered cell clusters (biofilm) is a key stage in the development of infections. In this regard, adhesion is considered as one of the most important pathogenic factors in *Staphylococcus aureus* (6). Instruments such as artificial organs, urethral catheters, and inside-vessel catheters are frequently used for patients being treated in special care divisions of hospitals. Furthermore, bacteria that adhere to these surfaces and form biofilms are hardly killed by an-

tibiotics (3). Researchers have shown that the first stage in infections caused by *Staphylococcus aureus* is the adhesion of this bacterium to surfaces such as medical instruments, host tissues, etc. This is attributed to a combination of factors external to the cell (e.g. the ability to adhere and form biofilm) (7). This stage is mediated by PIA (Polysaccharide Intercellular Adhesion) whose production is affected by intercellular adhesion proteins such as *IcaA*, *IcaB*, *IcaC*, and *IcaD* (8, 9). *icaA* and *icaD* genes (which include chromosomal genes) have a more significant role in forming biofilm in *Staphylococcus aureus* and *Staphylococcus epidermidis* (10). Genes such as *fmbA*, *fmbB* (fibronectin binding protein), and *clfB* (clumping factor) (which includes chromosomal genes) also play a considerable role in elevating the possibility of bacterial adhesion to surfaces (11). Formation of biofilms has significant health-related and economic consequences. It is estimated that 65 percent of nosocomial infections in the US are related to the formation of biofilms and the economic loss caused by biofilms equals more than one billion dollars per year (12). Research shows that adhesion to fibronectin through FnBPs is essential for attacking eukaryote cells (13). In this regard, the significant role of adhesion factors has been investigated in a wide range of diseases caused by *Staphylococcus aureus*. It has been proven that *fmbA* and *fmbB* are significantly influential in the formation of tissues in various pathologic conditions such as eye keratitis (14), osteomyelitis and septic arthritis (15) and substitution in the surface of medical instruments (16) in which *Staphylococcus* is seen as being an influential disease-producing germ in infections related to implant (injection of medicine or implanting artificial organs) (17). The aim of the current study was to investigate the biofilm-producing genes (*icaA*, *icaD*, *fmbA*, and *clfB*) in *Staphylococcus aureus* isolated from local cheese in Maragheh city.

2. Methods

2.1. Samples

In this research, 20 strains of *Staphylococcus aureus* bacterium extracted from local cheese (soft salty cheese) in Maragheh (already identified through biochemical experiments) were used as samples.

2.2. Extraction of DNA

To extract DNA, a rapid genomic extraction kit (MBST Company, Iran) was used. After extraction of DNA, the samples were read for determination of DNA concentration in ng/L using a Nanodrop device and on the basis of priority of reading indices, the ratio of 260/280 nm, the ratio of 260/230 nm, and optical density (OD) were read for performing PCR, respectively.

2.3. PCR Test to Detect Thermonuclease Gene (*nuc*)

The polymerase chain reaction was performed in 25 μ L volume, including extracted DNA containing 1 μ L (50 ng) Master mix, 12.5 μ L PCR, 0.4 μ M specific primers (0.5 μ L from each of the forward and reverse primers) (18) (Table 1) and 10.5 μ L of double distilled water. PCR conditions are presented in Table 2. PCR product in 1.5% agarose was electrophoresed and imaged using gel document. *S. aureus* PTCC1112 DNA extract was used as a positive control. Double distilled water was used for negative control. *S. aureus* isolates were considered as *S. aureus* after a definite diagnosis of Thermonuclease gene with a length of approximately 279 bp.

Table 1. The Sequence of Specific Primers Related to the Genes Under Investigation and PCR Products Resulting from Their Proliferation

Gene	Primer Sequence	PCR Product, bp
<i>nuc</i>	5'-GCGATTGATGGTGATACGGTT-3'	279
	5'-AGCCAAGCCTTGACGAACTAAAGC-3'	
<i>fmbA</i>	5'-CATAAATGGGAGCAGCATCA-3'	128
	5'-ATCAGCAGCTGAATTCCTT-3'	
<i>clfB</i>	5'-ACATCAGTAATAGTAGGGGCAAC-3'	203
	5'-TTCGCACGTGTTGTGTTGAC-3'	
<i>icaD</i>	5'-ATGGTCAAGCCAGACAGAG-3'	198
	5'-AGTATTTCAATGTTTAAAGCAA-3'	
<i>icaA</i>	5'-ACACTGCTGGCGCAGTCAA-3'	188
	5'-TCTGGAACCAACATCCAACA-3'	

Table 2. PCR Conditions to Multiply Thermonuclease Gene (*nuc*)

Cycle	Step	Temperature, °C	Time, Min
1	primary denaturation	94	4
	Denaturation	94	1
32	Annealing	55	1
	Extension	72	1
1	Final extension	72	10

2.4. PCR Test to Identify the Biofilm-Producing Genes Under Investigation

To perform PCR, the Taq DNA polymerase kit (QIAGEN Company, Catalogue no. 201223) was utilized. In a 25 μ L volume, this reaction includes 12.5 μ L of double distilled water, 2.5 μ L of buffer, 1 μ L of dNTPs, 3 μ L of primers (1.5 μ L of forward primer and 1.5 μ L of reverse primer), 0.75 μ L

of $MgCl_2$, 0.25 μL of Taq DNA Polymerase, and 5 μL of extracted DNA. The sequence of specific primers used for *icaA*, *icaD*, *clfB*, and *fnbA* genes (19-21) is indicated in Table 1. The proliferation of target genes in thermocycler was carried out through the following timetable (Table 3).

Table 3. PCR Conditions to Multiply *icaA*, *icaD*, *clfB* and *fnbA* Genes

Cycle	Step	Temperature, °C	Time
1	Primary denaturation	94	4 min
	Denaturation	72	30 sec
35	Annealing	55	30 sec
	Extension	72	1 min
1	Final extension	72	10 min

PCR products related to the proliferation of each of the adhesion genes under investigation were studied through electrophoresis on agarose gel, which was observed by gel documentation. In order to show the proliferated pieces, DNA Ladder 100 bp was utilized. Extracted *Staphylococcus aureus* PTCC 1112 DNA was used as positive control. Double distilled water was used as negative control.

3. Results

17 strains (85%) out of 20 bacterial strains were recognized as positive species-specific *Staphylococcus aureus* in terms of thermonuclease (nus). The frequencies of each of the genes of the investigated strains were *icaA* (76.47%), *clfB* (76.47%), *icaD* (58.82%), and *fnbA* (64.7%) (Figure 1). 5 strains (29.41%) simultaneously included all the genes under investigation (Table 2).

4. Discussion

This research reported the frequencies of *icaA* (76.47%), *clfB* (76.47%), *icaD* (58.82%), and *fnbA* (64.7%) genes. 5 strains (29.41%) simultaneously included all the genes under investigation. Cifter et al. (2009) reported that 25.42% of *Staphylococcus aureus* bacteria extracted from mastitis included simultaneously *icaA* and *icaD* genes, 27.11% of them included *icaA* gene, and 64.4% included *icaD* gene (21). The frequency of *icaD* gene is in line with the findings of the present study. Shojaei et al. (2014) reported that the frequencies of *icaA*, *icaD*, *clfB*, and *fnbA* in *Staphylococcus aureus* extracted from raw milk were 59.18%, 38.77%, 32.6%, and 69.38%, respectively (11). This contrasts with the findings of the present study. Basanisi et al. (2016) found that 94.6% of *Staphylococcus aureus* bacteria extracted from dairy products of sheep and goat in the south of Italy included *icaA*

gene (22). The widespread distribution of *Staphylococcus aureus* strains including biofilm genes in milk increases the risk of the presence of these strains in food products. Investigating the capacity of adhesion and formation of biofilm as well as the genes playing a role in the formation of biofilms in different *Staphylococcus aureus* strains can provide invaluable information for better understanding of the complex process of biofilm formation and infections resulting from these organisms (19). This bacterium has a high ability to form biofilms, which can be of significance in the development of chronic infections and the development of strains resistant to antibiotics. Furthermore, through different mechanisms of transferring resistance against medicines, bacteria found in biofilms can elevate the resistance of infection-producing bacteria in special care divisions of hospitals; this phenomenon can be considered as a serious threat for patients being treated in medical centers. When compared to the findings of other studies, the results of the current research show a difference in the distribution of the genes that play a role in the formation of biofilms among the strains of *Staphylococcus aureus*; this could be due to the geographical differences and the differences in ecological origins of the extracted strains (Milk, humans, and other animals).

According to the findings of the current research and other similar investigations, none of the samples of the *Staphylococcus aureus* samples was negative in terms of the presence of the four genes being investigated. This shows the high ability of this bacterium in forming biofilms. It is suggested conducting more research on the precise combination of the structure of the *Staphylococcus aureus* biofilm so that this structure can be more precisely targeted by the medicines. In addition, protein combinations of the structure of biofilm capsule of this bacterium can be used as a vaccine against biofilm infections.

Acknowledgments

We thank all people who contributed to conducting this research.

Footnotes

Authors' Contribution: Saman Mahdavi designed the study, analyzed data, and prepared the manuscript; Bahareh Falaki conducted the experimental studies.

Conflicts of Interest: The authors have no conflicts of interest to declare.




Figure 1. Findings of PCR Related to Strains with *nuc*, *icaD*, *icaA*, *clfB* and *fnbA* Genes

Table 4. Frequencies of Biofilm-Producing Genes and Frequency of Being-Togetherness of these Genes in One Strain

Gene	Number of Isolates	Combination of Genes	Number of Isolates
<i>icaD</i>	10	<i>icaA</i> + <i>clfB</i>	1
<i>icaA</i>	13	<i>icaA</i> + <i>fnbA</i>	1
<i>clfB</i>	13	<i>icaA</i> + <i>icaD</i> + <i>clfB</i>	7
<i>fnbA</i>	11	<i>icaA</i> + <i>icaD</i> + <i>fnbA</i>	1
		<i>icaA</i> + <i>clfB</i> + <i>fnbA</i>	2
		<i>icaD</i> + <i>clfB</i> + <i>fnbA</i>	1
		<i>icaA</i> + <i>icaD</i> + <i>clfB</i> + <i>fnbA</i>	5
Total	17		

References

- Razavilar V. *Pathogenic microorganisms in foods and epidemiology of food poisoning*. 2nd ed. Tehran university press; 2003. p. 128-9.
- Mansuri Najand L, Ghanbarpour R. A study on enteropathogenic *Escherichia coli* isolated from domestic Iranian soft cheese. *Vet Arh*. 2006;**76**(6):531-6.
- Revidwala S, Rajdev BM, Mulla S. Characterization of bacterial etiologic agents of biofilm formation in medical devices in critical care setup. *Crit Care Res Pract*. 2012;**2012**:945805. doi: [10.1155/2012/945805](https://doi.org/10.1155/2012/945805). [PubMed: [22312484](https://pubmed.ncbi.nlm.nih.gov/22312484/)].
- Costerton JW, Stewart PS, Greenberg EP. Bacterial biofilms: a common cause of persistent infections. *Science*. 1999;**284**(5418):1318-22. doi: [10.1126/science.284.5418.1318](https://doi.org/10.1126/science.284.5418.1318). [PubMed: [10334980](https://pubmed.ncbi.nlm.nih.gov/10334980/)].
- Stepanovic S, Vukovic D, Jezek P, Pavlovic M, Svabic-Vlahovic M. Influence of dynamic conditions on biofilm formation by staphylococci. *Eur J Clin Microbiol Infect Dis*. 2001;**20**(7):502-4. doi: [10.1007/s100960100534](https://doi.org/10.1007/s100960100534). [PubMed: [11561809](https://pubmed.ncbi.nlm.nih.gov/11561809/)].
- Donlan RM. Biofilms and device-associated infections. *Emerg Infect Dis*. 2001;**7**(2):277-81. doi: [10.3201/eid0702.700277](https://doi.org/10.3201/eid0702.700277). [PubMed: [11294723](https://pubmed.ncbi.nlm.nih.gov/11294723/)].
- Chaieb K, Mahdouani K, Bakhruf A. Detection of *icaA* and *icaD* loci by polymerase chain reaction and biofilm formation by *Staphylococcus epidermidis* isolated from dialysate and needles in a dialysis unit. *J Hosp Infect*. 2005;**61**(3):225-30. doi: [10.1016/j.jhin.2005.05.014](https://doi.org/10.1016/j.jhin.2005.05.014). [PubMed: [16165246](https://pubmed.ncbi.nlm.nih.gov/16165246/)].
- Atshan SS, Nor Shamsudin M, Sekawi Z, Lung LT, Hamat RA, Karunanidhi A, et al. Prevalence of adhesion and regulation of biofilm-related genes in different clones of *Staphylococcus aureus*. *J Biomed Biotechnol*. 2012;**2012**:976972. doi: [10.1155/2012/976972](https://doi.org/10.1155/2012/976972). [PubMed: [22701309](https://pubmed.ncbi.nlm.nih.gov/22701309/)].
- Foster TJ, Hook M. Surface protein adhesins of *Staphylococcus aureus*. *Trends Microbiol*. 1998;**6**(12):484-8. doi: [10.1016/S0966-842X\(98\)01400-0](https://doi.org/10.1016/S0966-842X(98)01400-0). [PubMed: [10036727](https://pubmed.ncbi.nlm.nih.gov/10036727/)].
- O'Connell DP, Nanavaty T, McDevitt D, Gurusiddappa S, Hook M, Foster TJ. The fibrinogen-binding MSCRAMM (clumping factor) of *Staphylococcus aureus* has a Ca²⁺-dependent inhibitory site. *J Biol Chem*. 1998;**273**(12):6821-9. [PubMed: [9506984](https://pubmed.ncbi.nlm.nih.gov/9506984/)].
- Shojaei M, Karimi Darehabi H, Javadi A. Distribution of genes encoding biofilm production in *S. aureus* isolated from raw milk in Kurdistan [In Persian]. *J Food Hygiene*. 2014;**4**(14):1-8.
- Hauck CR, Ohlsen K. Sticky connections: extracellular matrix protein recognition and integrin-mediated cellular invasion by *Staphylococcus aureus*. *Curr Opin Microbiol*. 2006;**9**(1):5-11. doi: [10.1016/j.mib.2005.12.002](https://doi.org/10.1016/j.mib.2005.12.002). [PubMed: [16406780](https://pubmed.ncbi.nlm.nih.gov/16406780/)].
- Jett BD, Gilmore MS. Internalization of *Staphylococcus aureus* by human corneal epithelial cells: role of bacterial fibronectin-binding protein and host cell factors. *Infect Immun*. 2002;**70**(8):4697-700. [PubMed: [12117986](https://pubmed.ncbi.nlm.nih.gov/12117986/)].

14. Smeltzer MS, Gillaspay AF. Molecular pathogenesis of staphylococcal osteomyelitis. *Poult Sci.* 2000;**79**(7):1042-9. doi: [10.1093/ps/79.7.1042](https://doi.org/10.1093/ps/79.7.1042). [PubMed: [10901208](https://pubmed.ncbi.nlm.nih.gov/10901208/)].
15. Cheung AL, Fischetti VA. The role of fibrinogen in mediating staphylococcal adherence to fibers. *J Surg Res.* 1991;**50**(2):150-5. doi: [10.1016/0022-4804\(91\)90239-1](https://doi.org/10.1016/0022-4804(91)90239-1). [PubMed: [1990220](https://pubmed.ncbi.nlm.nih.gov/1990220/)].
16. Patti JM, Bremell T, Krajewska-Pietrasik D, Abdelnour A, Tarkowski A, Ryden C, et al. The Staphylococcus aureus collagen adhesin is a virulence determinant in experimental septic arthritis. *Infect Immun.* 1994;**62**(1):152-61. [PubMed: [8262622](https://pubmed.ncbi.nlm.nih.gov/8262622/)].
17. Brakstad OG, Aasbakk K, Maeland JA. Detection of Staphylococcus aureus by polymerase chain reaction amplification of the nuc gene. *J Clin Microbiol.* 1992;**30**(7):1654-60. [PubMed: [1629319](https://pubmed.ncbi.nlm.nih.gov/1629319/)].
18. Rohde H, Knobloch JK, Horstkotte MA, Mack D. Correlation of Staphylococcus aureus icaADBC genotype and biofilm expression phenotype. *J Clin Microbiol.* 2001;**39**(12):4595-6. doi: [10.1128/JCM.39.12.4595-4596.2001](https://doi.org/10.1128/JCM.39.12.4595-4596.2001). [PubMed: [11797608](https://pubmed.ncbi.nlm.nih.gov/11797608/)].
19. Vancaeynest D, Hermans K, Haesebrouck F. Genotypic and phenotypic screening of high and low virulence Staphylococcus aureus isolates from rabbits for biofilm formation and MSCRAMMs. *Vet Microbiol.* 2004;**103**(3-4):241-7. doi: [10.1016/j.vetmic.2004.09.002](https://doi.org/10.1016/j.vetmic.2004.09.002). [PubMed: [15504595](https://pubmed.ncbi.nlm.nih.gov/15504595/)].
20. Tristan A, Ying L, Bes M, Etienne J, Vandenesch F, Lina G. Use of multiplex PCR to identify Staphylococcus aureus adhesins involved in human hematogenous infections. *J Clin Microbiol.* 2003;**41**(9):4465-7. doi: [10.1128/JCM.41.9.4465-4467.2003](https://doi.org/10.1128/JCM.41.9.4465-4467.2003). [PubMed: [12958296](https://pubmed.ncbi.nlm.nih.gov/12958296/)].
21. Ciftci A, Findik A, Onuk EE, Savasan S. Detection of methicillin resistance and slime factor production of Staphylococcus aureus in bovine mastitis. *Braz J Microbiol.* 2009;**40**(2):254-61. doi: [10.1590/S1517-83822009000200009](https://doi.org/10.1590/S1517-83822009000200009). [PubMed: [24031354](https://pubmed.ncbi.nlm.nih.gov/24031354/)].
22. Basanisi MG, Nobili G, La Bella G, Russo R, Spano G, Normanno G, et al. Molecular characterization of Staphylococcus aureus isolated from sheep and goat cheeses in southern Italy. *Small Rumin Res.* 2016;**135**:17-9. doi: [10.1016/j.smallrumres.2015.12.024](https://doi.org/10.1016/j.smallrumres.2015.12.024).