Published online 2018 January 31.

Can Vitamin D Prevent Breast Cancer?

Maryam Moossavi,^{1,2} and Milad Mohammadoo-Khorasani^{3,*}

¹Student Research Committee, Birjand University of Medical Sciences, Birjand, Iran
²Department of Molecular Medicine, Birjand University of Medical Sciences, Birjand, Iran
³Department of Clinical Biochemistry, Tarbiat Modares University, Tehran, Iran

^{*} *Corresponding author*: Milad Mohammadoo-Khorasani, Department of Clinical Biochemistry, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, Iran. Tel: +98-9350641028, E-mail: miladkh24@yahoo.com

Received 2017 November 30; Revised 2018 January 02; Accepted 2018 January 15.

Keywords: Vitamin D, Breast Cancer, Prevent

Dear Editor,

Vitamin D (25(OH)D) deficiency is a worldwide health problem. It can increase the risk of many diseases, notably cancers (1-3). More than two billion people have vitamin D deficiency (25(OH)D < 50 nmol/L) around the world (4). Vitamin D is classified in the secosteroids hormone, which is hydroxylated in the liver to form 25-hydroxy vitamin D (25(OH)D). After further hydroxylation, the biologically active metabolite, 1,25(OH)2D, was present in the kidneys (5). Breast cancer is one of the most frequently diagnosed cancers and it is one of the leading causes of cancer-related mortality around world (6). Growing evidences have shown that vitamin D deficiency can lead to breast cancer development (7,8). Animal studies indicated that vitamin D can inhibit tumorigenic effects of fatty diet. Other studies have demonstrated that vitamin D can induce cell cycle arrest in G0/G1. Also, it can induce morphological and biochemical features of apoptosis in breast cancer cells (9).

A meta-analysis conducted by Chen et al., showed that vitamin D level has an effect on breast cancer susceptibility. They reported that vitamin D and calcium had a chemopreventive effect against breast cancer (10). Quite surprisingly in another meta-analysis, Ordonez-Mena et al. reported that increased breast cancer risk was linked with higher amount of 25(OH)D concentration (11). This discrepancy results may be contributed to different situations, different populations, and differences in the adjusted levels.

Stoll et al. recommended that serum level of 25(OH)D, which is obtained through sun exposure, dietary intake, and vitamin D supplementation is more than 400 IU per day, which can decrease breast cancer risk (12). Bilinski et al. showed that lower than 75 nmol/L 25(OH)D concentration was associated with a significantly higher risk of

breast cancer (13). Also, Park et al. reported that serum 25(OH) D below 20 ng/mL was associated with 27% augmentation in the risk of breast cancer (14).

Some studies have shown low levels of vitamin D among breast cancer patients and about 94% of females with vitamin D level of less than 20 ng/mL developed metastases and 73% of them died from the advanced stage of the disease (15).

In conclusion, most vitamin D studies reported an opposite association between vitamin D level and breast cancer risk. However, more studies are needed to detect the optimum level of serum vitamin D as a prophylaxis therapy for breast cancer prevention.

References

- Feldman D, Krishnan AV, Swami S, Giovannucci E, Feldman BJ. The role of vitamin D in reducing cancer risk and progression. *Nat Rev Cancer*. 2014;14(5):342–57. doi: 10.1038/nrc3691. [PubMed: 24705652].
- Klampfer L. Vitamin D and colon cancer. World J Gastrointest Oncol. 2014;6(11):430–7. doi: 10.4251/wjgo.v6.i11.430. [PubMed: 25400874]. [PubMed Central: PMC4229786].
- Jacobs ET, Kohler LN, Kunihiro AG, Jurutka PW. Vitamin D and Colorectal, Breast, and Prostate Cancers: A Review of the Epidemiological Evidence. J Cancer. 2016;7(3):232–40. doi: 10.7150/jca.13403. [PubMed: 26918035]. [PubMed Central: PMC4747876].
- Al-Musharaf S, Fouda MA, Turkestani IZ, Al-Ajlan A, Sabico S, Alnaami AM, et al. Vitamin D Deficiency Prevalence and Predictors in Early Pregnancy among Arab Women. *Nutrients*. 2018;10(4). doi: 10.3390/nu10040489. [PubMed: 29662044]. [PubMed Central: PMC5946274].
- Narooie-Nejad M, Moossavi M, Torkamanzehi A, Moghtaderi A, Salimi S. Vitamin D Receptor Gene Polymorphism and the Risk of Multiple Sclerosis in South Eastern of Iran. *J Mol Neurosci*. 2015;56(3):572–6. doi: 10.1007/s12031-015-0513-x. [PubMed: 25854779].
- 6. Stewart B, Wild CP. World cancer report 2014. NY: Health; 2017.
- Yao S, Kwan ML, Ergas IJ, Roh JM, Cheng TD, Hong CC, et al. Association of Serum Level of Vitamin D at Diagnosis With Breast Cancer Survival: A Case-Cohort Analysis in the Pathways Study. JAMA Oncol.

Letter

Copyright © 2018, Gene, Cell and Tissue. This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (http://creativecommons.org/licenses/by-nc/4.0/) which permits copy and redistribute the material just in noncommercial usages, provided the original work is properly cited

2017;**3**(3):351-7. doi: 10.1001/jamaoncol.2016.4188. [PubMed: 27832250]. [PubMed Central: PMC5473032].

- O'Brien KM, Sandler DP, Taylor JA, Weinberg CR. Serum Vitamin D and Risk of Breast Cancer within Five Years. *Environ Health Perspect*. 2017;**125**(7):77004. doi: 10.1289/EHP943. [PubMed: 28728134]. [PubMed Central: PMC5744694].
- Colston KW, Hansen CM. Mechanisms implicated in the growth regulatory effects of vitamin D in breast cancer. *Endocr Relat Cancer*. 2002;9(1):45-59. doi: 10.1677/erc.0.0090045. [PubMed: 11914182].
- Chen P, Hu P, Xie D, Qin Y, Wang F, Wang H. Meta-analysis of vitamin D, calcium and the prevention of breast cancer. *Breast Cancer Res Treat.* 2010;**121**(2):469–77. doi: 10.1007/s10549-009-0593-9. [PubMed: 19851861].
- Ordonez-Mena JM, Schottker B, Fedirko V, Jenab M, Olsen A, Halkjaer J, et al. Pre-diagnostic vitamin D concentrations and cancer risks in older individuals: an analysis of cohorts participating in the CHANCES consortium. Eur J Epidemiol. 2016;31(3):311–23. doi:

10.1007/s10654-015-0040-7. [PubMed: 25977096].

- Stoll F, Akladios CY, Mathelin C. [Vitamin D and breast cancer: is there a link?]. *Gynecol Obstet Fertil.* 2013;41(4):242–50. doi: 10.1016/j.gyobfe.2013.02.002. [PubMed: 23562418].
- Bilinski K, Boyages J. Association between 25-hydroxyvitamin D concentration and breast cancer risk in an Australian population: an observational case-control study. *Breast Cancer Res Treat*. 2013;**137**(2):599– 607. doi: 10.1007/s10549-012-2381-1. [PubMed: 23239153].
- Park S, Lee DH, Jeon JY, Ryu J, Kim S, Kim JY, et al. Serum 25hydroxyvitamin D deficiency and increased risk of breast cancer among Korean women: a case-control study. *Breast Cancer Res Treat.* 2015;**152**(1):147–54. doi: 10.1007/s10549-015-3433-0. [PubMed: 26037255].
- Imtiaz S, Siddiqui N, Raza SA, Loya A, Muhammad A. Vitamin D deficiency in newly diagnosed breast cancer patients. *Indian J Endocrinol Metab.* 2012;16(3):409–13. doi: 10.4103/2230-8210.95684. [PubMed: 22629509]. [PubMed Central: PMC3354850].