



# Weight-Loss Parameters and Quality of Life in Obese Aged Women Using WATERinMOTION

Mohammadreza Rezaeipour <sup>1, \*</sup>, Gennady Leonidovich Apanasenko <sup>2</sup> and Maryam Banparvari <sup>1</sup>

<sup>1</sup>Sports Sciences Department, University of Sistan and Baluchestan, Zahedan, Iran

<sup>2</sup>Ukrainian Center of Sports Medicine, Kyiv, Ukraine

\*Corresponding author: Department of Sports Sciences, University of Sistan and Baluchestan, Postal Code: 9816745639, Zahedan, Iran. Tel: +98-9153414047; +98-932903270, Email: rezaeipour@ped.usb.ac.ir

Received 2020 October 02; Revised 2021 February 22; Accepted 2021 April 15.

## Abstract

**Background:** There are controversial studies on weight loss, cardiometabolic markers, and quality of life (QOL) in obese aged women participating in water sports.

**Objectives:** This study aimed to determine the effects of the WATERinMOTION exercise program accompanied with no diet on the inactive obese aged women's weight, cardiometabolic markers, and QOL.

**Methods:** Sixty-four inactive obese aged women were recruited to participate in this cross-sectional study at the Ukrainian Sports Center, Kyiv, 2019. They were randomly assigned into two Case (n = 32, with the WATERinMOTION aquatics exercise program) and control (n = 32, held in a sitting position simultaneously) groups using the convenient sampling method. The two programs were run twice a week (55 minutes each time) and lasted for one month. Moreover, anthropometric indices (weight, height, waist circumference), cardiometabolic markers (blood lipid markers and glycaemia), QOL (SF-36 questionnaire), and dietary monitoring were assessed before and after the intervention.

**Results:** The groups did not differ significantly regarding each of the analyzed variables prior to the intervention. Comparing pre- and post-intervention results revealed significant weight loss (-1.3, P = 0.004) and average BMI (-0.4, P = 0.002) in the case group. Moreover, a significant variation was revealed only in terms of weight after the intervention (P = 0.001). Regarding QOL, improvements in health status, vitality, and social aspects were significant in the case group after the intervention.

**Conclusions:** The present study suggests that the WATERinMOTION program accompanied with no diet has a positive effect on weight loss, waist circumference, and metabolic profiles in obese aged women. However, the improvement in the QOL should not be neglected.

**Keywords:** Cross-Sectional Studies, Metabolic Profiles, Obesity, Quality of Life, Resistance Training, Waist Circumference, Water Sports, Women

## 1. Background

Communication, mental health, emotional intelligence, problem-solving ability, and well-being are the most prominent features among the resilient, which help individuals adapt to changes and are also essential for navigating challenging living and working conditions in the 21st century (1). Some factors such as obesity, poor fitness, aging, and neurological and respiratory disorders arouse restrictions, which limit individuals' potentials to enter land-based exercise programs (2). According to Raffaelli et al. (3), aquatic exercise significantly enhance physical exercise, compared to land exercise.

Changing the position of the joints can affect individuals' daily activities and participation in sports (4). Several studies have documented that when the body is immersed

in water up to the shoulders, it loses "90% of gravity" (5, 6). Accordingly, water exercises are particularly beneficial for obese individuals, who are at enhanced risk of orthopedic damage, which is secondary consequences of exercise.

However, the benefits of the aquatics exercise programs in terms of health status are still controversial (5). There is no study on weight loss, cardiometabolic markers, and quality of life (QOL) for obese aged women participating in the WATERinMOTION aquatics exercise program. This physical exercise is highly welcomed by some grownups and aged individuals as it is believed that, in addition to promoting health status, the program is also associated with improvements in physical fitness and weight-related diseases such as cardiovascular and metabolic syndrome. The WATERinMOTION program is a standardized

water sports program consisting of a warm-up, lineal and sidelong motions, teamwork, group dynamic, workouts, upper and lower body suspension, strengthening body core, and flexibility (7). Music is professional, and individuals at different age ranges can participate in this program. In this regard, choreography is correspondingly linked to the music.

Meanwhile, participants in other programs participate in aquatic exercise programs for at least six-week (8) or longer (2, 5, 6, 9). The present study, however, was to detect if the program held two sessions a week is sufficient for a significant increase in health in a month. Waist circumference (WC) was measured by considering internal fat deposits and the likelihood of weight-related diseases (10). Body mass index (BMI) was used to define obesity (11). Personality traits are linked to subjective well-being and described as overall emotional and mental health (12). The QOL questionnaire addresses individuals' emotional and mental health, physical status, functional capacity, general health status, pain, vitality, and social aspects (5, 13). Dietary monitoring was also performed to detect dietary changes (5, 9).

## 2. Objectives

The present study aimed to determine whether the effectiveness of WATERinMOTION as a water sport in helping inactive obese aged women regarding weight loss, WC, and cardiometabolic markers (such as blood lipid markers and glycaemia), and QOL. It was hypothesized that a one-month WATERinMOTION exercise may be an appropriate option to have positive shifts in the participants' weight, cardiometabolic markers, and QOL. The outline of the study is presented in [Figure 1](#).

## 3. Methods

### 3.1. Study Design, Sample, and Randomization

The participants of this cross-sectional study were recruited from inactive obese aged women who had come to the Ukrainian Sports Center (Kyiv, 2019) after reading the ads on the website to participate in the one-month WATERinMOTION exercise program. Given the potential population of 76 individuals and a 95% confidence level with five confidence intervals, the sample size was determined based on the using Creative Research systems not to be < 64 individuals (14). They were randomly assigned into two case (n = 32, with the WATERinMOTION aquatics exercise program) and control (n = 32, held in a sitting position simultaneously) groups using the convenient sampling method. The two one-month programs were run

twice a week (a total of eight 55-minute sessions). A team of two physicians and two instructors explained the study objectives and procedure to the volunteers. Inclusion criteria were inactive old women having 1 - 2 and less exercise sessions per month (9), BMI > 30 kg.m<sup>2</sup>, and 65 years of age (5, 6). Living in self-sufficiency is categorized as degrees to which the person can sustain itself without external support (the average self-sufficiency status in this study was three). The Mini-Mental State examination is the most commonly used test to evaluate memory problems or other mental abilities. The value of the Mini-Mental State examination in our study was considered above 18 for inclusion criteria (15). In addition to a remarkable family history of dyslipidemia and cardiovascular diseases, volunteers with the age-related diseases (e.g., diabetes mellitus, cancer, cardiovascular, Alzheimer's, hypertension, orthopedic restriction, and multiple chronic conditions), hearing loss interfering listening to sports music, tobacco consumption (chronic and acute), transmittable diseases, and a medical WATERinMOTION restriction, and those taking drugs such as thyroxine, progesterin, estrogen, and vitamin E, which affect lipid metabolism, and beta-blockers as well as those who missed three sessions of the program were excluded from the study.

All the participants completed dietary records over four days (3 weekdays and a weekend day) during two weeks before the study, and the same procedure was run after the study for self-reporting the dietetic alteration (5, 6). These data were processed using the EVIDENT II application, which was authorized to determine the quantity of consumed energy/kilocalorie per day (16). Moreover, they were invited to preserve their nutrition habits and physical activities in their leisure time throughout the study period. The follow-up phase was performed for all the participants every other two weeks throughout the study period. The research project and informed consent forms were approved by the Institutional Research Ethics Committee. The volunteers submitted their signed informed consent form to participate in the study one week before the intervention. The study protocol was set in accordance with the Helsinki Declaration and observed all relevant institutional policies and strategies.

### 3.2. Pool Exercises

Like other types of exercises, the main phases of the WATERinMOTION program are warm-up, main set, and cool-down (7). [Table 1](#) demonstrates more aspects of the WATERinMOTION aquatics exercise program.

The WATERinMOTION program was performed in an indoor swimming pool. The water temperature was about 30°C (17), and the pool deepness was 1.30 m (17). The workout intensity ranged from 70% to 85% of the maximum




Figure 1. Paper outline

Table 1. Pool Exercise Program at WATERinMOTION

	Frequency	Intensity	Time (Duration)	Type (Exercise)
Warm-up (5 min)	Twice per week	55% - 70% of HRmax <sup>a</sup>	One month	Stretching
Main exercise (45 min)		70% - 85% of HRmax <sup>b</sup>		Linear and lateral movements; team building; group dynamic exercises; suspension of upper and lower body; core strengthening; flexibility
Cooldown (5 min)		55% - 70% of HRmax		Stretching

Abbreviations: HRmax, maximum heart rate; min, minutes.

<sup>a</sup>Moderate intensity.

<sup>b</sup>High intensity.

heart rate (5). Polar Electro Oy- a waterproof device for measuring heart rate- manufactured by Kempele, Finland was used to control the participants’ heart rate. Furthermore, the music played in the WATERinMOTION program fits everyone in terms of speed and tempo as such everyone could follow the play (7).

3.3. Assessments

Before the first session of the WATERinMOTION program, the researchers assessed the participants’ anthropometric indices, including weight in kilogram (kg) and height in centimeter (cm), according to which the participants’ BMI was calculated. Before and after the intervention, the participants’ weights were calculated with a Scale-Tronix model 5002 manufactured by Wheaton, IL, USA, and their heights were measured with a stadiometer, while considering the exactness of 0.1 kg and 0.1 cm for both weight and height, respectively (5, 6). These calculations were made for the participants dressed in light clothing and no shoes. In this study, BMI and WC were used to study body composition (9, 18). BMI is a pattern considering “an individual’s weight unit (in kg) divided by the height squared (in meters)” (11). Obesity is considered as BMI ≥ 30 (11). Waist circumference (WC) was measured without compressing the skin in the horizontal plate at the end of the natural expiratory in the middle of the iliac protrusion to the lowest ribs (9).

Morning biochemical examinations were accomplished from 9 AM, after an hour-long rest, in a half-lying position (9), and after 12-hour night starvation before and after the intervention (6, 19). Total cholesterol (T-C), high-density lipoprotein cholesterol (HDL-C), and triglyceride (TG) were analyzed by considering the enzymatic manner of COBAS, manufactured by Roche Diagnostics, Germany. Low-density lipoprotein cholesterol (LDL-C) was computed based on Friedewald’s formula: LDL-C = TC - (HDL-C + TG/5). This mode can only be used for TG < 400 mg/dL (20). Fasting blood sugar was checked by a kit to determine the enzymatic glycaemia (Wiener® liquid AA) By oxidation method by Rosario, Argentina.

The SF-36 questionnaire was used to assess the participants’ QOL. This questionnaire was selected as it was versatile, validated, and self-applicable, useful for individual patient assessment, and sensitive to progress in the QOL indices (5, 21).

3.4. Statistical Analysis

The results are reported as mean ± SD. The Shapiro-Wilk test supported the normality of the data distribution. The paired sample t-test was used to compare the groups and time points (pre/post-intervention), and the analysis of covariance (ANCOVA) was also used to compare the results between the two groups. Pearson’s chi-square (χ<sup>2</sup>)

test was applied to investigate the variations in the distribution of categorical variables. In his study,  $P < 0.05$  was set as the significance level, and IBM SPSS software version 24.0 (for Windows) was utilized to analyze the data.

## 4. Results

### 4.1. Demographics, Anthropometric, and Cardiometabolic Markers

The mean age and height of the participants were  $69.7 \pm 4.4$  years and  $167.3 \pm 5.4$  meters in the case group and  $70.5 \pm 4.5$  years and  $167.9 \pm 4.7$  meters in the control group, respectively. Table 2 shows the other demographic information, including weight and BMI, of the participants. No specific event such as refusals, adverse reactions, and drop-out were reported in this study. All the participants completed the study procedure. The completion rate for the groups was estimated by  $\chi^2$ , and no significant variation was noticed ( $P = 0.78$ ). At the beginning of the study (pre-intervention), there was no significant difference between the groups' daily calorie intake, anthropometric indices, and cardiometabolic markers ( $P > 0.05$ ) (Table 2).

Table 2. Variation in daily calorie intake, anthropometric indices, and cardiometabolic markers in case and control groups during the study (pre/post-intervention).

The study results in terms of time (pre-/post-intervention) revealed modest but significant weight loss ( $P = 0.004$ ) and a significant decrease in BMI ( $P = 0.002$ ) in the case group. The case group had mean weight loss of  $-1.3$  kg and BMI of  $-0.4$  kg.m<sup>2</sup>. The case group displayed better, but insignificant, effects regarding cardiometabolic markers (T-C,  $-3.6$  mL/dl,  $P = 0.21$ ; LDL-C,  $-3.7$  mL/dL,  $P = 0.09$ ; HDL-C,  $+1.3$  mL/dl,  $P = 0.1$ ; FBS,  $-1.1$  mL/dl,  $P = 0.12$ ) and WC ( $-0.2$  cm,  $P = 0.14$ ). Furthermore, the comparison of the study groups in the post-intervention phase only revealed a significant variation in weight loss ( $P = 0.001$ ) (Table 2).

### 4.2. QOL

The groups were homogenous in terms of QOL indicators at the beginning of the study ( $P > 0.05$ ). Comparing pre-and post-intervention results discloses significant improvements in health status ( $P = 0.03$ ), vitality ( $P = 0.01$ ), and social aspects ( $P = 0.04$ ) in the case group (Table 3).

## 5. Discussion

The research results showed the effect of the WATERinMOTION approach on the weight loss and cardiometabolic markers of inactive obese aged women. This aquatics exercise program revealed a slight superiority by making some

improvements in QOL; however, physical exertion with a dietary restriction revealed more effectiveness in terms of weight loss after eight weeks in Fett's study (22) on obese men. The present findings were in line with those of the meta-analysis and review reports in dictating the moderate effectiveness of exercises in weight loss, particularly when exercise is not accompanied by dietary monitoring (23).

Meanwhile, aquatic exercises are considered as one of the main calorie burners and as an excellent technique to retain weight under control (24). However, there are still controversies regarding the impact of water exercises on weight (5). In Penaforte et al.'s study (25), a two-month water sports program held three sessions a week for obese aged women revealed a significant drop in body weight and BMI. Rezaeipour and Nychyporuk (9) also had a 12-week aquafit exercise held three sessions a week for postmenopausal women and reported a significant decrease in weight loss markers. In contrast, Charmas and Gromisz (26) found no significant change in weight for young women who swam 12 weeks with three sessions per week.

A calorie-burning exercise raises the likelihood of changes in body weight (27). The Water viscosity is resistant to any physical movement of the WATERinMOTION program and leads to more energy-burning. Preserving a calorie shortage in water sports would have a positive impact on weight loss (28). The moderate weight loss in this study may be caused by a lack of compensatory increase in calorie consumption to adapt to energy expenditure during sports, suggesting that the WATERinMOTION program leads to further weight loss when held two sessions a week per month.

The triglyceride level of blood depends on food intake; hence, it was not assessed in this study (20). The present findings also exhibited positive but not significant changes in the cardiometabolic markers such as blood lipid markers and glycaemia. The findings were in contrast with those reported by Caro et al. (29) and Nakhaei et al. (30), who studied moderate-intensity regular physical exercise.

High WC aroused by visceral fat may be a risk factor resulting in many diseases such as arthritis and cardiometabolic disorders (31). The significant associations between the concerned program with weight and BMI and positive reductions in cardiometabolic markers confirmed the research hypothesis, suggesting that a one-month WATERinMOTION program can be effective in decreasing weight and improving cardiac metabolic markers in inactive obese aged women. The WATERinMOTION exercise program, however, is not suitable for individuals at health risk as it elevates cardiometabolic markers.

The control group revealed an insignificant weight

**Table 2.** Variations in Daily Calorie Intake, Anthropometric Indices, and Cardiometabolic Markers in Case and Control Groups (Pre- and Post-intervention)<sup>a, b</sup>

Variables	Case Group		Control Group	
	Pre-intervention	Post-intervention	Pre-intervention	Post-intervention
Weight, kg	98.7 ± 16.4	97.4 ± 16.7*#	100.4 ± 18.3	100.6 ± 18.2
BMI, kg.m <sup>-2</sup>	35.1 ± 5.1	34.7 ± 5.2*	35.5 ± 5.4	35.5 ± 5.5
T-C, mg/dL	224.9 ± 29.5	221.3 ± 27.6	226.1 ± 25.6	225.7 ± 27.7
LDL-C, mg/dL	146.1 ± 28.7	142.4 ± 25.8	140.1 ± 26.0	139.8 ± 28.1
HDL-C, mg/dL	48.5 ± 5.9	49.8 ± 6.0	51.8 ± 6.7	51.8 ± 7.2
FBS, mg/dL	94.1 ± 16.1	93.0 ± 17.8	95.6 ± 17.3	95.8 ± 16.9
WC, cm	100.6 ± 10.7	100.4 ± 10.1	101.5 ± 11.2	101.5 ± 12.7
Daily calorie intake, Kcal/day	2199 ± 188	2221 ± 167	2218 ± 119	2225 ± 106

Abbreviations: BMI, body mass index; FBS, fasting blood sugar; HDL-C, high-density lipoprotein cholesterol; LDL-C, low-density lipoprotein cholesterol; T-C, total cholesterol; WC, waist circumference.

<sup>a</sup>Values are expressed as mean ± SD.

<sup>b</sup>\*, P < 0.05 in comparison to pre-intervention #, P < 0.05 in comparison to the control group.

**Table 3.** QOL Indicators in Case vs. Control Groups<sup>a, b</sup>

SF-36 Variables	Case Group		Control Group	
	Pre-intervention	POST-intervention	Pre-intervention	Post-intervention
Emotional aspects	60.2 ± 13.1	60.7 ± 11.8	71.2 ± 8.1	69.3 ± 11.6
Mental health	63.4 ± 10.5	66.2 ± 11.3	63.6 ± 8.2	63.9 ± 7.9
Health status	70.1 ± 9.5	80.4 ± 9.7	70.4 ± 6.7	73.2 ± 7.6
Physical aspects	82.6 ± 11.4	88.4 ± 9.6	84.1 ± 10.2	84.9 ± 11.3
Functional capacity	70.7 ± 8.7	78.4 ± 7.9	71.9 ± 9.5	72.3 ± 8.8
Pain	65.7 ± 12.3	61.8 ± 9.7	56.8 ± 10.1	61.3 ± 10.5
Vitality	52.9 ± 7.1	63.6 ± 10.2	57.8 ± 11.5	58.1 ± 7.5
Social aspects	65.6 ± 8.9	76.1 ± 7.8	70.3 ± 6.5	70.3 ± 7.1

<sup>a</sup>Values are expressed as mean ± SD.

<sup>b</sup>\*, P < 0.05.

gain. Although the difference was not statistically meaningful, it may lead to marked weight gain over time, which is associated with a higher risk of different chronic diseases (32). The same can be achieved when analyzing blood glucose. The results of the WC assessment in the case group were positive but insignificant. Another study on the elderly showed that a more significant decrease in WC for walking in water and longer durations (from 24 to 48 weeks) (33), thus indicating better results in comparison to the present findings.

Regarding QOL markers, the case group showed significant improvements in health status, vitality, and social aspects. Research has documented that regular aerobic exercise programs have positive effects on mood and psychological disorders such as depression and anxiety and cognitive aspects (i.e., learning and memory), improve QOL, and hence are crucial therapeutic allies (5, 34). Furthermore,

the increased synthesis of endorphins caused by regular physical activity also leads to progression in individuals' organisms (5, 34).

### 5.1. Conclusions

The study findings demonstrate that the short-term WATERinMOTION exercise program accompanied with no dietetical supervision has a modest effect on individuals' weight loss. Improvements in the participants' health status, vitality, and social aspects this program suggests the positive effects of this type of exercise on individuals' QOL; hence, including such a program in inactive obese aged women's lifestyle is beneficial in terms of WC and cardiometabolic markers. These findings may have some clinical implications. Since the elderly women were included in this study and given the metabolic differences rising from their age and gender, the findings cannot be recom-

mended for other individuals. In this regard, future researchers are recommended to study the proposed protocol for a long period as it would improve the clinical outlook of obese individuals with an inactive lifestyle. Furthermore, further studies are suggested to investigate males and other age groups.

## Footnotes

**Authors' Contribution:** Mohammadreza Rezaeipour wrote the paper. Gennady Leonidovich Apanasenko examined the cardiometabolic markers and participated in the data collection. Maryam Banparvari and Mohammadreza Rezaeipour performed the statistical analysis. Mohammadreza Rezaeipour performed the assays for metabolic profiles and weight loss. Maryam Banparvari assessed the QOL indicators. Gennady Leonidovich Apanasenko and Mohammadreza Rezaeipour conceived the study, participated in the research design and coordination, and helped in drafting the manuscript. All authors studied and approved the final manuscript.

**Conflict of Interests:** The authors declared no conflict of interest.

**Ethical Approval:** This study complies with all relevant national statutes, institutional policies, and Helsinki Declaration Principles. The samples participated in this study after receiving their informed consent and with the approval of the Institutional Research Ethics Committee [NMAPE, (2019) no.: 04112-06]. The participants presented their physicians' permission for their participation in the offered intervention.

**Funding/Support:** The authors declare no funding or support.

**Informed Consent:** The volunteers submitted their signed informed consent form to participate in the study one week before the intervention.

## References

- Gamble B, Crouse D. Strategies for Supporting and Building Student Resilience in Canadian Secondary and Post-Secondary Educational Institutions. *Sci Med J*. 2020;**2**(2):70–6. doi: [10.28991/SciMedJ-2020-0202-4](https://doi.org/10.28991/SciMedJ-2020-0202-4).
- Pereira Neiva H, Brandao Fail L, Izquierdo M, Marques MC, Marinho DA. The effect of 12 weeks of water-aerobics on health status and physical fitness: An ecological approach. *PLoS One*. 2018;**13**(5). e0198319. doi: [10.1371/journal.pone.0198319](https://doi.org/10.1371/journal.pone.0198319). [PubMed: [29851998](https://pubmed.ncbi.nlm.nih.gov/29851998/)]. [PubMed Central: [PMC5978883](https://pubmed.ncbi.nlm.nih.gov/pmc/articles/PMC5978883/)].
- Raffaelli C, Lanza M, Zanolla L, Zamparo P. Exercise intensity of head-out water-based activities (water fitness). *Eur J Appl Physiol*. 2010;**109**(5):829–38. doi: [10.1007/s00421-010-1419-5](https://doi.org/10.1007/s00421-010-1419-5). [PubMed: [20229021](https://pubmed.ncbi.nlm.nih.gov/20229021/)].
- Paranjape S, Singhania N. Effect of Body Positions on Quadriceps Angle Measurement. *Sci Med J*. 2019;**1**(1):20–4. doi: [10.28991/SciMedJ-2019-0101-3](https://doi.org/10.28991/SciMedJ-2019-0101-3).
- Rezaeipour M. Investigation of Pool Workouts on Weight, Body Composition, Resting Energy Expenditure, and Quality of Life among Sedentary Obese Older Women. *Montenegrin J Sports Sci Med*. 2020;**9**(1):67–72. doi: [10.26773/mjssm.200309](https://doi.org/10.26773/mjssm.200309).
- Rezaeipour M, Apanasenko GL. Effects of Waterobics Programs on Body Mass, Body Composition, and Coronary Risk Profile of Sedentary Obese Middle-aged Women. *Women's Health Bul*. 2019;**6**(4):13–7.
- Darley T. *Effectiveness of an Aquatic Exercise Program for Reducing Weight, Body Fat and Chronic Low Back and Joint Pain*. 2020.
- Ferrigan K, Hice J, Leemkuil K, Singer S, Charles D, Michaels NN, et al. Aquatic exercise for weight reduction in middle-aged adults: a pilot study. *J Aquat Phys Ther*. 2017;**25**(2):16–21.
- Rezaeipour M, Nychporuk VI. Study of Weight Loss Parameters Among Sedentary, Overweight Postmenopausal Females Using Different Time Models of Aquafit. *Hormozgan Med J*. 2019;**In Press**(In Press). doi: [10.5812/hmj.96378](https://doi.org/10.5812/hmj.96378).
- Baran M, Celikkalkan K, Cagan Appak Y, Karakoyun M, Bozkurt M, Kocyigit C, et al. Body Fat Mass is Better Indicator than Indirect Measurement Methods in Obese Children for Fatty Liver and Metabolic Syndrome. *Sci Med J*. 2019;**1**(4):168–75. doi: [10.28991/SciMedJ-2019-0104-2](https://doi.org/10.28991/SciMedJ-2019-0104-2).
- Weight H. *About Adult BMI*. 2017.
- An L, Liu C, Zhang N, Chen Z, Ren D, Yuan F, et al. GRIK3 rs490647 is a Common Genetic Variant between Personality and Subjective Well-being in Chinese Han Population. *Emerg Sci J*. 2019;**3**(2). doi: [10.28991/esj-2019-01171](https://doi.org/10.28991/esj-2019-01171).
- Ciconelli RM, Ferraz MB, Santos W, Meinão I, Quesada MR. Tradução para a língua portuguesa e validação do questionário genérico de avaliação de qualidade de vida SF-36 (Brasil SF-36). *Rev Bras Reumatol*. 1999;**39**(3):143–50.
- Systems CR. *Sample size calculator*. 2012. Available from: <https://www.surveysystem.com/sscalc.htm>.
- Ogden CL, Carroll MD, Curtin LR, McDowell MA, Tabak CJ, Flegal KM. Prevalence of overweight and obesity in the United States, 1999–2004. *JAMA*. 2006;**295**(13):1549–55. doi: [10.1001/jama.295.13.1549](https://doi.org/10.1001/jama.295.13.1549). [PubMed: [16595758](https://pubmed.ncbi.nlm.nih.gov/16595758/)].
- Recio-Rodriguez JJ, Rodriguez-Martin C, Gonzalez-Sanchez J, Rodriguez-Sanchez E, Martin-Borras C, Martinez-Vizcaino V, et al. EVIDENT Smartphone App, a New Method for the Dietary Record: Comparison With a Food Frequency Questionnaire. *JMIR Mhealth Uhealth*. 2019;**7**(2). e11463. doi: [10.2196/11463](https://doi.org/10.2196/11463). [PubMed: [30735141](https://pubmed.ncbi.nlm.nih.gov/30735141/)]. [PubMed Central: [PMC6384535](https://pubmed.ncbi.nlm.nih.gov/pmc/articles/PMC6384535/)].
- Bure A. *Physical Activity In The Water And The Human Body*. 2019.
- Purath J, Buchholz SW, Kark DL. Physical fitness assessment of older adults in the primary care setting. *J Am Acad Nurse Pract*. 2009;**21**(2):101–7. doi: [10.1111/j.1745-7599.2008.00391.x](https://doi.org/10.1111/j.1745-7599.2008.00391.x). [PubMed: [19228248](https://pubmed.ncbi.nlm.nih.gov/19228248/)].
- Rezaeipour M. Comparison of Two Types of Diets on Losing Weight and Lipid Profile of Overweight/Obese Middle-Aged Women Under Exercise Condition. *Health Scope*. 2014;**3**(3). doi: [10.17795/jhealthscope-15611](https://doi.org/10.17795/jhealthscope-15611).
- Rezaeipour M. Study of exercise time models on weight loss and coronary risk panel in inactive middle-aged men by overweight or obesity. *J Urmia Univ Med Sci*. 2018;**29**(5):389–97.
- de Castro M, Caiuby AV, Draibe SA, Canziani ME. [Quality of life in chronic renal disease patients submitted to hemodialysis evaluated with SF-36 instrument]. *Rev Assoc Med Bras (1992)*. 2003;**49**(3):245–9. Portuguese. [PubMed: [14666346](https://pubmed.ncbi.nlm.nih.gov/14666346/)].
- Fett CA. *Avaliação metabólica nutricional de obesas no basal e após tratamento com dieta hipocalórica e treinamento em circuito ou caminhada*. Universidade de São Paulo; 2005.
- Khachatoorian Y, Samara A. Differential effects of dietary restriction combined with exercise vs dietary restriction alone on visceral and subcutaneous adipose tissues: A systematic review. *Obes Med*. 2018;**9**:7–17. doi: [10.1016/j.obmed.2017.11.002](https://doi.org/10.1016/j.obmed.2017.11.002).

24. Belza B, Topolski T, Kinne S, Patrick DL, Ramsey SD. Does adherence make a difference? Results from a community-based aquatic exercise program. *Nurs Res.* 2002;**51**(5):285–91. doi: [10.1097/00006199-200209000-00003](https://doi.org/10.1097/00006199-200209000-00003). [PubMed: [12352776](https://pubmed.ncbi.nlm.nih.gov/12352776/)].
25. Penaforte FRO, Calhau R, Mota GR, Chiarello PG. Impact of short-term water exercise programs on weight, body composition, metabolic profile and quality of life of obese women. *J Hum Sport Exercise.* 2015;**10**(4). doi: [10.14198/jhse.2015.104.07](https://doi.org/10.14198/jhse.2015.104.07).
26. Charmas M, Gromisz W. Effect of 12-Week Swimming Training on Body Composition in Young Women. *Int J Environ Res Public Health.* 2019;**16**(3). doi: [10.3390/ijerph16030346](https://doi.org/10.3390/ijerph16030346). [PubMed: [30691153](https://pubmed.ncbi.nlm.nih.gov/30691153/)]. [PubMed Central: [PMC6388213](https://pubmed.ncbi.nlm.nih.gov/PMC6388213/)].
27. Petridou A, Siopi A, Mougios V. Exercise in the management of obesity. *Metabolism.* 2019;**92**:163–9. doi: [10.1016/j.metabol.2018.10.009](https://doi.org/10.1016/j.metabol.2018.10.009). [PubMed: [30385379](https://pubmed.ncbi.nlm.nih.gov/30385379/)].
28. Layne MR. Water Exercise. *Hum Kinet.* 2015.
29. Caro J, Navarro I, Romero P, Lorente RI, Priego MA, Martinez-Hervas S, et al. Metabolic effects of regular physical exercise in healthy population. *Endocrinol Nutr.* 2013;**60**(4):167–72. doi: [10.1016/j.endonu.2012.11.004](https://doi.org/10.1016/j.endonu.2012.11.004). [PubMed: [23517693](https://pubmed.ncbi.nlm.nih.gov/23517693/)].
30. Nakhaei H, Mogharnasi M, Fanaei H. Effect of swimming training on levels of asprosin, lipid profile, glucose and insulin resistance in rats with metabolic syndrome. *Obes Med.* 2019;**15**. doi: [10.1016/j.obmed.2019.100111](https://doi.org/10.1016/j.obmed.2019.100111).
31. Carr DB, Utzschneider KM, Hull RL, Kodama K, Retzlaff BM, Brunzell JD, et al. Intra-abdominal fat is a major determinant of the National Cholesterol Education Program Adult Treatment Panel III criteria for the metabolic syndrome. *Diabetes.* 2004;**53**(8):2087–94. doi: [10.2337/diabetes.53.8.2087](https://doi.org/10.2337/diabetes.53.8.2087). [PubMed: [15277390](https://pubmed.ncbi.nlm.nih.gov/15277390/)].
32. Elisha B, Azar M, Taleb N, Bernard S, Iacobellis G, Rabasa-Lhoret R. Body Composition and Epicardial Fat in Type 2 Diabetes Patients Following Insulin Detemir Versus Insulin Glargine Initiation. *Horm Metab Res.* 2016;**48**(1):42–7. doi: [10.1055/s-0035-1554688](https://doi.org/10.1055/s-0035-1554688). [PubMed: [26340704](https://pubmed.ncbi.nlm.nih.gov/26340704/)].
33. Naylor LH, Maslen BA, Cox KL, Spence AL, Robey E, Haynes A, et al. Land- versus water-walking interventions in older adults: Effects on body composition. *J Sci Med Sport.* 2020;**23**(2):164–70. doi: [10.1016/j.jsams.2019.08.019](https://doi.org/10.1016/j.jsams.2019.08.019). [PubMed: [31506246](https://pubmed.ncbi.nlm.nih.gov/31506246/)].
34. Anish EJ. Exercise and its effects on the central nervous system. *Curr Sports Med Rep.* 2005;**4**(1):18–23. doi: [10.1097/01.csmr.0000306066.14026.77](https://doi.org/10.1097/01.csmr.0000306066.14026.77). [PubMed: [15659274](https://pubmed.ncbi.nlm.nih.gov/15659274/)].