Published online 2022 May 17.

Research Article

Understanding Household Catastrophic Health Expenditures and Fairness of Financing for Cancer Treatment: A Cross-sectional Case Study in West of Iran

¹Health Management and Economics Research Center, Iran University of Medical Sciences, Tehran, Iran

²Department of Public Health, Lorestan University of Medical Sciences, Khorramabad, Iran

³Department of Health Management & Economics, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

⁴Department of Health Services Management, School of Health Management and Information Sciences, Iran University of Medical Sciences, Tehran, Iran

⁵Department of Health Economics, School of Health Management and Information Sciences, Iran University of Medical Sciences, Tehran, Iran

^{*} Corresponding author: Department of Health Economics, School of Health Management and Information Sciences, Iran University of Medical Sciences, Tehran, Iran. Email: payam_haghighatfard@yahoo.com

Received 2021 October 04; Revised 2022 March 16; Accepted 2022 April 06.

Abstract

Background: Direct out-of-pocket (OOP) and indirect healthcare payments can limit the household budget and cause several financial problems for the household.

Objectives: This study aimed to measure the financial protection and determinants of catastrophic health expenditures (CHEs) for cancer treatment in Shahid Rahimi Hospital, Khorramabad, located in western Iran.

Methods: This study was conducted on 220 households of cancer patients in Lorestan Province, Iran. The framework of data collection was based on the World Health Organization (WHO) Global Health Survey. Interviews were conducted with individuals who met the inclusion criteria of this study; they were selected using a simple random sampling method. Data were analyzed using Chi-square test in STATA.

Results: The present results showed that the incidence of CHEs and the fair financial contribution index (FFCI) were 70% and 86%, respectively. There was a significant relationship between the household CHEs and variables, such as supplementary insurance coverage, household income status, educational level of the household head, household size, age of the household head, type of cancer, and type of cancer treatment.

Conclusions: More financial protection should be provided for the families of cancer patients due to the high incidence of CHEs and unfair financing of cancer care services. Moreover, healthcare systems should consider supportive policies for cancer patients and their household members by increasing the insurance coverage and expanding service packages to reduce cancer treatment expenditures.

Keywords: Iran, Catastrophic Health Expenditure, Out-of-Pocket, Cancer

1. Background

The financial burden of non-communicable diseases is high in most countries around the world, and more and more families seem to fall into poverty due to chronic diseases (1). The out-of-pocket payment (OOP) of healthcare services can limit the household budget and expose the household members to several financial problems. Each year, millions of people face financial difficulty meeting their living expenses due to high healthcare expenditures, and vulnerable populations are forced to spend their income on health services (2). The costs of diagnosis and treatment of diseases can increase the rate of poor households, especially in rural areas; accordingly, human rights organizations, including the World Health Organization (WHO), have emphasized the need for universal health coverage. This goal cannot be achieved unless access to healthcare services is available for all people, without the incidence of catastrophic expenditures (3, 4).

Catastrophic health expenditures (CHEs) and the fair financial contribution index (FFCI), as major indices for evaluating the performance of healthcare systems, represent the level of financial support for people facing health expenditures (5). These indices are considered unfair when

Aziz Rezapour ¹, Soraya Norayi Motlagh ², Banafsheh Darvishi Teli³, Negar Yousefzadeh⁴ and Payam Haghighatfard ^{5,*}

Copyright © 2022, Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (http://creativecommons.org/licenses/by-nc/4.0/) which permits copy and redistribute the material just in noncommercial usages, provided the original work is properly cited.

the household payments for health services significantly exceed the household budget (6). CHEs occur when OOPs exceed 10% of the total annual income or 40% of the household payment capacity. The FFCI indicates the level of inequality in OOPs for health, with scores ranging from 0 (perfect inequality) to 1 (perfect equality). According to some studies, about two-thirds of chronically ill patients and 42% of chronically ill households face CHEs due to insufficient financial support (7, 8).

Cancer, as a chronic non-communicable disease, can increase the risk of CHEs (9). This disease is caused by uncontrolled cell proliferation, which may extend to other body organs (10). More than 100 types of cancer have been identified so far, which are treated using various methods, including surgery, chemotherapy, and radiotherapy, depending on the type of disease. According to the global report of the International Cancer Agency in 2020, the number of new cases and deaths caused by cancer was estimated at 19.3 and 10 million people, respectively (11, 12), suggesting a significant increase compared to previous global cancer studies (13, 14).

In 2010, nearly 808 million people around the world faced CHEs. Due to a rapid increase in the number of new cancer cases over the past decade, the financial burden of this disease has increased. Research suggests that any type of expenditure imposed directly on households reduces their tendency to use health services. Therefore, the compulsory prepayment system (insurance), funded by tax credits or compulsory insurance premiums, is the most effective and fair approach to finance the treatment expenditures of cancer patients (15, 16).

In a cross-sectional study, Rezapour et al. examined the effects of direct OOPs on 2,200 households in Tehran, Iran, using two approaches of CHE and impoverishing health expenditure. According to their results, although the need for healthcare services was strongly felt in poor households, inequality in access to these services was evident, and healthcare utilization was higher in affluent groups and households. Their findings showed that the educational level of the household head, household size, and frequency of inpatient care service utilization had positive relationships with the possibility of facing catastrophic expenditures, while the presence of a preschool child in the household reduced the likelihood of these expenditures. However, health insurance did not show a significant relationship with the rate of exposure to catastrophic expenditures (17).

Additionally, in a descriptive analytical study by Kavosi et al., CHEs and coping strategies were investigated in the households of cancer patients in Namazi Hospital of Shiraz, Iran. Their results showed that 67% of families of cancer patients experienced CHEs. There was a significant relationship between CHEs and type of insurance, history of outpatient services, type of cancer treatment, and household members refusing to receive health services (18). Moreover, Piroozi et al. investigated the determinants and inequality of CHEs in diabetes mellitus treatment in Iran. The results of their study showed that 57.5% of the patients were exposed to CHEs. The socioeconomic status, sex, and marital status were the main determinants of CHE inequality among patients with type 2 diabetes mellitus (19).

Kasahun et al. investigated the rate of exposure to catastrophic expenditures among cancer patients in Ethiopia's capital and found that 74.4% of cancer patient households faced catastrophic expenditures. Based on the results of their study on the type of treatment, chemotherapy imposed more catastrophic expenditures than radiotherapy. Moreover, households with lower income were more exposed to the catastrophic expenditures of cancer. With advancing age and increasing frequency of chemotherapy, exposure to the catastrophic expenditures of cancer has also increased (20). Moreover, the incidence and mortality rates of cancer have increased in Iran in recent years (21). Therefore, the current study was carried out, as little research has been conducted on cancerrelated CHEs in Iran.

2. Objectives

The present study aimed to investigate the equality of financing for cancer treatment, to estimate CHEs, and to determine the influential factors for presenting policymaking solutions to reduce CHEs for the households of cancer patients, referred to Shahid Rahimi Hospital in Khorramabad, as the main public center for cancer patients in Lorestan Province, Iran.

3. Methods

3.1. Study Setting

This study was conducted in Khorramabad, located in western Iran, with 18 public hospitals and five private medical centers. Shahid Rahimi Educational and Medical Center, as a referral hospital for cancer patients, was selected as the research environment. There are wards in this center designated to cancer diagnosis and treatment.

3.2. Study Design

This hospital-based, cross-sectional study was conducted to investigate the catastrophic expenditures and fairness of financing for cancer treatment.

3.3. Study Population

The statistical population of this study included all households of cancer patients, referred to the radiotherapy and chemotherapy wards of Shahid Rahimi Hospital in June 2017.

3.4. Inclusion Criteria

The inclusion criteria were cancer patients, referred to the radiotherapy or chemotherapy wards of the hospital (or both) in Lorestan Province, Iran.

3.5. Exclusion Criteria

The exclusion criteria were other acute cancer patients, who usually received treatment in other provinces of Iran.

3.6. Sample Size

According to a previous study, 67.9% of cancer patient households faced CHEs [18]. With almost 2,500 registered cancer patients in Lorestan Province, a sample size of 220 was measured at P = 0.679 (q = 1-p = 0.321, error = 0.05) and an accuracy level of e = 0.06 (z = 1.96), based on the following formula:

Sample size
$$= \frac{\frac{z^2 p(1-p)}{e^2}}{1 + \frac{z^2 p(1-p)}{e^2 N}}$$

By visiting the hospital wards, the researchers collected the data through interview, using a simple random sampling method. A face-to-face household interview conducted for each household randomly and each interview lasted about 30 minutes.

3.7. Study Variables

In this study, the dependent variable was household CHE, and the independent variables were characteristics, including the educational level of the household head, sex, age, and other variables, such as type of cancer, insurance type, supplementary insurance coverage, treatment type, utilization of inpatient services, residency and household size.

3.8. Data Collection

In the present study, a WHO questionnaire, called the Global Health Survey, was used to collect data. This questionnaire was developed in 2003 to measure the performance of health systems, based on the three objectives of the health system. This questionnaire was translated into Persian in a previous study (18). The results showed that the translated version had adequate reliability and validity. Generally, it consists of three main parts, including the demographic information, household questionnaire, and

individual questionnaire. In the present study, the identification information and household questionnaires were used. The time allowed for recalling information related to the total household expenditures, consumption expenditures, and outpatient treatment expenditures was one month (one year for hospitalized cases). Regarding the socioeconomic status, information was collected on the total expenditure of each household (22, 23).

3.9. Data Analysis

Based on the WHO approach, to determine the CHEs and fair financing of cancer treatment, the income and total household expenditures spent on food and non-food items (e.g., housing, education, transportation, and other expenditures) were determined. In this study, a two-stage method was applied for data analysis. The first step was understanding the household CHEs, and the second step was determining the fairness of financial contributions for cancer treatment expenditures. To estimate the CHEs, food expenditure and household size were converted using the following formula (β = 0.56):

$Equisize_h = hhsize_h^\beta$

To determine the poverty line (PL), first, the percentage of food expenditure to total expenditure was calculated for each household by dividing the food expenditures to the equivalent household size calculated in the previous step to determine the PL. The average food expenditure of households within the 45th and 55th percentiles was considered as PL. The subsistence expenditures (SEs) were also calculated for each household using the following formula:

$SE = equisiz_h \times PL$

The criterion for determining the household status regarding PL is to compare the calculated SEs with the total household expenditures. If the total household expenditures are lower than the SEs, the household is below PL (poor). In the next step, capacity to pay (CTP) was measured using the following formula:

$CTP_i = EXP_iSE_i$

where CTP_i is the capacity to pay, EXP_i is the household expenditure, and SE_i is the subsistence expenditure. Finally, by calculating the ratio of health expenditures to CTP for each household and comparing it with the reference threshold (ratio ≥ 0.40), the percentage of households exposed to CHEs was determined (24, 25). Besides, to determine the factors affecting CHEs, a Chi-square test was used, and a significance level of 0.05 was considered (P < 0.05). Data were analyzed in STATA version 14. In this study, to measure the financial protection of households, the WHO approach was applied. The following formula was used for estimating the fairness of financing for cancer treatment:

$$FFCI = 1 - \frac{\sqrt{\sum_{h}^{n} w_{h} |oopctp_{h} - ooptp_{o}|^{3}}}{\sum w_{h}}$$

Where w_h is the household weight $oopctp_h = oop_h ctp_h$ and $oopctp_o = \sum_{k} \frac{w_h oop_h}{w_h ctp_h}$, and OOP represents the proportion of OOP to CTP for each household. Further details on the calculation of the above-mentioned index are discussed in the literature (17, 26). The acceptable range of FFCI was 0 (perfect inequality in financing for cancer care) to 1 (perfect equality in financing for cancer care).

This article is part of a research project (grant No: 95-04-163-29776; ethics code: IR.IUMS.REC1395.95-04-163-29776). The research team would like to thank the financial support of the research deputy, affiliated to Iran University of Medical Sciences, Tehran, Iran.

4. Results

A total of 220 households were included in the study, about 70% of whom were exposed to CHEs. The FFCI was estimated at 0.857 for the study population. The rate of exposure to CHEs was calculated for different educational groups, income levels, age groups, and other independent variables; the results are presented in Table 1.

As shown in Table 1, the rate of exposure to catastrophic expenditures was lower in households whose head had academic education compared to other educational groups. Also, the rate of exposure to CHEs was higher in younger age groups (head of the household), and the difference was statistically significant. Considering the exposure of each household quintile or income level, the percentage of exposure to CHEs was higher in high-income groups compared to low-income groups. Moreover, the results showed that younger household heads were more exposed to CHEs compared to other age groups.

In Figure 1, the distribution of households exposed to CHEs at different income levels (quintiles) is presented. The first quintile is the poorest household, while the fifth quintile is the wealthiest. The highest percentage of households exposed to CHEs was observed in the richest group (the fifth quintile); in other words, 27.92% of households with CHEs belonged to this group.

As shown in Table 1, women-headed households were more likely to face CHEs, and this relationship was statistically significant. Besides, households with supplemental insurance coverage were less exposed to CHEs compared to other households, although this relationship was statistically insignificant. Households living in Khorramabad faced relatively lower catastrophic expenditures compared to households living in other cities; however, the difference was not statistically significant.

The household size had a significant reducing effect on the household CHEs; in other words, by increasing the household size, the likelihood of catastrophic expenditures decreased. The relationship between the incidence of CHEs for households of cancer patients and the type of cancer showed that the financial burden of cancer treatment was high in households with breast and lung cancer patients. Besides, exposure to these expenditures was higher in households with patients receiving chemotherapy and radiotherapy simultaneously as compared to households with patients receiving radiotherapy or chemotherapy separately. In terms of the type of insurance, households covered by social security and medical service companies were more exposed to CHEs compared to households covered by the Relief Foundation and rural insurance companies. Also, inpatient service utilization increased the likelihood of exposure to CHEs.

5. Discussion

In many countries, especially less developed and developing ones, the rate of exposure to CHEs is high for cancer treatment (27). Studies show that CHE is proportionally high in Iran (28). In this study, the number of households facing CHEs and the FFCI were measured as two key indices. The results regarding the financial burden of cancer imposed on patients and their households in Lorestan Province were exposed to CHEs, and the FFCI of healthcare financing for the studied households was estimated at 0.857. Nonetheless, the values of these indices vary in previous studies (24, 26).

The incidence of CHEs in the current study was proportionally higher than other similar studies in Iran. In this regard, the results of a study by Kavosi et al. in Shiraz, Iran, indicated that 67.9% of households of cancer patients were exposed to CHEs (18). Moreover, Kasahun et al. found that the incidence of CHE for cancer patient households was about 75% (20). According to general health policies, the FFCI, as a national indicator, must be 0.9 at minimum (26); however, this requirement was not met in our study; in other words, inequity in financing for cancer treatment was significant in the study population. In a similar study, Rezapour et al. (17) reported a lower FFCI (0.68) in Iran. Another study from Kermanshah Province, Iran, measured this index to be 0.57(29). Consistent with the current study, Murray reported the value of this index to range from 0.74 in Brazil to 0.941 in Slovakia (30). In recent years, although

Exposure to CHEs Pelale Kex No Pelale Sex of the household head 67(81.63) 38 (19.36) 98 Male 67(81.63) 28 (35.24) 0.0387 Female 87(65.75) 28 (35.24) 0.0387 Mate 66(67.27) 74 (32.73) 0.0387 Illterate 36 (67.27) 74 (32.73) 1 Primary school 15 (70) 35 (30) 1 Secondary school 7(70.83) 17 (27.16) 1 Secondary school 7(70.83) 17 (27.16) 1 Academic 35 (37.14) 4 (42.86) 1 Quintile 35 (37.2) 24 (17.24) 1 Quintile 3 35 (32.27) 14 (47.36) 1 Quintile 4 32 (52.27) 14 (47.30) 1 Quintile 5 33 (37.51) 12 (27.27) 1 Quintile 4 32 (52.27) 12 (47.33) 2 Quintile 5 43 (97.33) 12 (27.27) 2 Quintile 5	Table 1. The Relationship Between Exposure to Catastrophic Health Expenditures (CHEs) and the Study Variables ^a				
Vec No Sex of the household head 0.0495 Male 67 (81.63) 38 (9.36) Female 87 (65.75) 28 (35.24) Educational level of the household head 87 (65.77) 74 (32.73) Illiterate 36 (67.27) 74 (32.73) Primary school 15 (70) 35 (30) Secondary school 7 (70.83) 17 (27.16) Secondary school 7 (70.83) 17 (27.16) Oppiona 6 (82.76) 24 (172.16) Academic 3 (57.14) 4 (42.80) Academic 3 (57.14) 4 (24.80) Quintile 33 (0.75) 11 (25) Quintile 3 23 (52.27) 21 (47.73) Quintile 3 23 (52.27) 21 (47.73) Quintile 4 32 (52.73) 12 (27.27) Quintile 5 43 (97.73) 12 (27.27) Quintile 4 32 (52.73) 12 (27.27) Quintile 5 43 (97.73) 12 (27.27) Quintile 5 43 (97.73) 12 (27.27) Quintile 5 <th>Characteristics</th> <th colspan="2">Exposure to CHEs</th> <th>P-Value</th>	Characteristics	Exposure to CHEs		P-Value	
Male 67 (81.63) 38 (19.36) Female 87 (65.75) 28 (35.24) Educational level of the household head 0.0387 0.0387 Illiterate 36 (67.27) 74 (32.73) Primary school 15 (70) 35 (30) Secondary school 7(70.33) 17 (27.16) Diploma 5 (82.76) 24 (17.24) Academic 3 (57.14) 4 (42.86) Quintile 1 33 (0.75) 11 (25) Quintile 2 23 (52.27) 21 (47.73) Quintile 3 23 (52.27) 21 (47.73) Quintile 4 32 (72.73) 12 (27.27) Quintile 3 23 (52.27) 12 (47.73) Quintile 4 32 (72.73) 12 (27.27) Quintile 5 43 (97.73) 12 (27.27) Quintile 4 32 (72.73) 12 (27.27) Quintile 5 41 (85.42) 7 (14.58) 30 - 60 57 (67.06) 28 (32.94) < 60 57 (67.06) 28 (32.94) < 60 56 (63.71) 10.61	Saw of the household head	Yes	No	0.0105	
Nat. O (0.03) G (0.03) G (0.03) Female 87 (65.75) 28 (35.4) Female 86 (67.27) 74 (32.73) Illiterate 36 (67.27) 74 (32.73) Primary school 15 (70) 35 (30) Secondary school 7(70.83) 17 (27.16) Diploma 5 (82.76) 24 (17.24) Academic 3 (57.14) 4 (42.86) Income quintile 3 (0.75) 11 (25) Quintile 1 33 (0.75) 11 (25) Quintile 2 23 (52.27) 21 (47.73) Quintile 3 23 (52.27) 21 (47.73) Quintile 4 32 (07.73) 12 (27.27) Quintile 5 30 (97.3) 12 (27.27) Quintile 4 32 (07.73) 12 (27.27) Quintile 5 4 (85.42) 7(14.58) 30 - 60 57 (67.06) 28 (32.94) > 60 56 (64.37) 31 (35.63) Supplementary insurance 0.493 Covered 8 (54.61) 5 (46.38) Not	Sex of the household head	67 (81 62)	28 (10.26)	0.0495	
Fende 6 (95.75) 26 (35.24) Educational level of the household head 0.0387 Illiterate 36 (67.27) 74 (32.73) Primary school 15 (70) 35 (30) Secondary school 7 (70.83) 17 (27.16) Diploma 5 (82.76) 24 (17.24) Academic 3 (57.14) 4 (42.86) Income quintile < < 0.001 Quintile 1 33 (0.75) 11 (25) Quintile 2 23 (52.27) 21 (47.73) Quintile 3 23 (52.27) 21 (47.73) Quintile 4 32 (72.73) 12 (27.27) Quintile 5 43 (97.37) 12 (27.27) Quintile 5 41 (85.42) 7 (14.58) 30 - 60 57 (67.06) 28 (32.94) > 60 56 (64.37) 31 (35.63) Supplementary i	Famala	07 (81.03)	38 (19.30)		
Read 0.055/ Illierate 36 (67.27) 74 (32.73) Primary school 15 (70) 35 (30) Secondary school 7(70.83) 17 (27.16) Diploma 5 (82.76) 24 (17.24) Academic 3 (57.14) 4 (42.86) Income quintile 3 (0.75) 11 (25) Quintile 1 33 (0.75) 11 (25) Quintile 2 23 (52.27) 21 (47.73) Quintile 3 23 (52.27) 21 (47.73) Quintile 4 32 (72.73) 12 (27.27) Quintile 5 43 (97.3) 12 (27.27) Quintile 4 32 (72.73) 12 (27.27) Quintile 5 43 (97.3) 12 (27.27) Quintile 5 43 (97.3) 12 (27.27) Quintile 5 41 (85.42) 7 (14.58) 30 - 60 57 (67.66) 28 (32.94) 30 - 60 56 (64.37) 31 (35.63) Supplementary insurance 0.493 Covered 8 (54.61) 5 (46.38) Not covered 14 (53.70)	Female	87(05.75)	28 (33.24)	0.0287	
Illiterate 36 (67.27) 74 (32.73) Primary school 15 (70) 35 (30) Secondary school 7 (70.83) 17 (27.16) Diploma 5 (82.76) 24 (17.24) Academic 3 (57.14) 4 (42.86) Income quintile 3 (57.14) 4 (42.86) Quintile1 33 (0.75) 11 (25) Quintile2 23 (52.27) 21 (47.73) Quintile3 23 (52.27) 21 (47.73) Quintile4 32 (52.27) 21 (47.73) Quintile3 23 (52.27) 21 (47.73) Quintile4 32 (52.73) 12 (27.27) Quintile5 43 (97.73) 12 (27.27) Quintile5 41 (85.42) 7 (14.58) 30 - 60 57 (67.06) 28 (32.94) 30 - 60 56 (64.37) 31 (35.63) Supplementary insurace 0.493 Covered <td< td=""><td>head</td><td></td><td></td><td>0.0387</td></td<>	head			0.0387	
Primary school 15 (70) 35 (30) Secondary school 7 (70.83) 17 (27.16) Diploma 5 (82.76) 24 (17.24) Academic 3 (57.14) 4 (42.86) Income quintile 33 (0.75) 11 (25) Quintile 1 33 (0.75) 11 (25) Quintile 2 23 (52.27) 21 (47.73) Quintile 3 23 (52.27) 21 (47.73) Quintile 4 32 (72.73) 12 (27.27) Quintile 5 43 (97.3) 12 (27.27) Quintile 5 43 (97.3) 12 (27.27) Ace of the household head (y) 7 (14.58) 0.029 < 30	Illiterate	36 (67.27)	74 (32.73)		
Secondary school $7(70.83)$ $17(27.16)$ Diploma $5(82.76)$ $24(17.24)$ Academic $3(57.14)$ $4(42.86)$ Income quintile $33(0.75)$ $11(25)$ Quintile 1 $33(0.75)$ $11(25)$ Quintile 2 $23(52.27)$ $21(47.73)$ Quintile 3 $23(52.27)$ $21(47.73)$ Quintile 4 $32(72.73)$ $12(27.27)$ Quintile 5 $43(97.73)$ $1(2.27)$ Age of the household head (y) 0.029 < 30 $41(85.42)$ $7(14.58)$ $30 - 60$ $57(67.06)$ $28(32.94)$ > 60 $56(64.37)$ $31(35.63)$ Supplementary insurance 0.493 $Covered$ $8(54.61)$ $5(46.38)$ Not covered $146(53.00)$ $61(47.29)$	Primary school	15 (70)	35 (30)		
Diploma 5 (82.76) 24 (17.24) Academic 3 (57.14) 4 (42.86) Income quintile < 0.001 Quintile 1 33 (0.75) 11 (25) Quintile 2 23 (52.27) 21 (47.73) Quintile 3 23 (52.27) 21 (47.73) Quintile 4 32 (72.73) 12 (27.27) Quintile 5 43 (97.3) 12 (27.27) Quintile 5 43 (97.32) 12 (27.27) Quintile 5 43 (97.73) 12 (27.27) Quintile 5 41 (85.42) 7 (14.58) 30 - 60 57 (67.06) 28 (32.94) > 60 56 (64.37) 31 (35.63) Supplementary insurance 0.493 Covered 8 (54.61) 5 (46.38) Not covered 146 (53.70) 61 (47.29)	Secondary school	7 (70.83)	17 (27.16)		
Academic 3 (57.14) 4 (42.86) Income quintile < 0.001 Quintile 1 33 (0.75) 11 (25) Quintile 2 23 (52.27) 21 (47.73) Quintile 3 23 (52.27) 21 (47.73) Quintile 4 32 (72.73) 12 (27.27) Quintile 5 43 (97.73) 12 (27.27) Sologe of the household head (y) 7 (14.58) 0.029 30 - 60 57 (67.06) 28 (32.94) 0.493 5 60 56 (64.37) 31 (35.63) 0.493 Govered 8 (54.61) 5 (46.38) Not covered 14 (6 (53.70)	Diploma	5 (82.76)	24 (17.24)		
Income quintile < 0.001 Quintile 1 33 (0.75) 11 (25) Quintile 2 23 (52.27) 21 (47.73) Quintile 3 23 (52.27) 21 (47.73) Quintile 4 32 (72.73) 12 (27.27) Quintile 5 43 (97.73) 12 (27.27) Solog 57 (67.06) 28 (32.94) 30 - 60 57 (67.06) 28 (32.94) 5 60 56 (64.37) 31 (35.63) Supplementary insurance 0.493 Covered 8 (54.61) 5 (46.38) Not covered 146	Academic	3 (57.14)	4 (42.86)		
Quintile 1 33 (0.75) 11 (25) Quintile 2 23 (52.27) 21 (47.73) Quintile 3 23 (52.27) 21 (47.73) Quintile 4 32 (72.73) 12 (27.27) Quintile 5 43 (97.73) 12 (27.27) Quintile 5 43 (97.73) 12 (27.27) Age of the household head (y) 0.029 <30	Income quintile			< 0.001	
Quintile 2 23 (52.27) 21 (47.73) Quintile 3 23 (52.27) 21 (47.73) Quintile 4 32 (72.73) 12 (27.27) Quintile 5 43 (97.73) 12 (27.27) Quintile 5 43 (97.73) 12 (27.27) Age of the household head (y) 12 (27.27) 0.029 < 30	Quintile 1	33 (0.75)	11 (25)		
Quintile 3 23 (52.27) 21 (47.73) Quintile 4 32 (72.73) 12 (27.27) Quintile 5 43 (97.73) 12 (27.27) Age of the household head (y) 12 (27.27) 0.029 < 30	Quintile 2	23 (52.27)	21 (47.73)		
Quintile 4 32 (72.73) 12 (27.27) Quintile 5 43 (97.73) 1(2.27) Age of the household head (y) 0.029 < 30	Quintile 3	23 (52.27)	21 (47.73)		
Quintile 5 43 (97.73) 1(2.27) Age of the household head (y) 0.029 < 30 41 (85.42) 7 (14.58) 30 - 60 57 (67.06) 28 (32.94) > 60 56 (64.37) 31 (35.63) Supplementary insurance 0.493 Covered 8 (54.61) 5 (46.38) Not covered 146 (53.70) 61 (47.29)	Quintile 4	32 (72.73)	12 (27.27)		
Age of the household head (y) 0.029 < 30	Quintile 5	43 (97.73)	1(2.27)		
< 30	Age of the household head (y)			0.029	
30-60 57 (67.06) 28 (32.94) > 60 56 (64.37) 31 (35.63) Supplementary insurance 0.493 Covered 8 (54.61) 5 (46.38) Not covered 146 (53.70) 61 (47.29)	< 30	41(85.42)	7(14.58)		
> 60 56 (64.37) 31 (35.63) Supplementary insurance 0.493 Covered 8 (54.61) 5 (46.38) Not covered 146 (53.70) 61 (47.29)	30 - 60	57 (67.06)	28 (32.94)		
Supplementary insurance 0.493 Covered 8 (54.61) 5 (46.38) Not covered 146 (53.70) 61 (47.29)	> 60	56 (64.37)	31 (35.63)		
Covered 8 (54.61) 5 (46.38) Not covered 146 (53.70) 61 (47.29)	Supplementary insurance	, ,		0.493	
Not covered 146 (53.70) 61 (47.29)	Covered	8 (54.61)	5(46.38)		
	Not covered	146 (53.70)	61(47.29)		
Insurance type 0.033	Insurance type	(55/14)	()	0.033	
Relief Foundation 28 (56) 22 (44)	Relief Foundation	28 (56)	22 (44)		
Social Security 64 (01.79) 17 (99.20)	Social Security	64 (01.79)	17 (99.20)		
Health Service 25 (76.75) 8 (24.24)	Health Service	25 (76.75)	8(24.24)		
Residence 0.575	Residence			0.575	
Rural areas 37(07.66) 19(93.33)	Rural areas	37 (07.66)	19 (93.33)		
Khorramabad 81 (5.67) 39 (5.32)	Khorramabad	81 (5.67)	39 (5.32)		
Other cities 73 (73) 27 (27)	Other cities	73 (73)	27 (27)		
Treatment type 0.57	Treatment type		. ,	0.57	
Chemotherapy 108 (79.68) 49 (21.31)	Chemotherapy	108 (79.68)	49 (21.31)		
Radiotherapy 16(66.66) 8(33.33)	Radiotherapy	16 (66.66)	8 (33.33)		
Both 30 (92.76) 9 (08.23)	Both	30 (92.76)	9(08.23)		
Utilization of inpatient services 0.0421	Utilization of inpatient services	- ()	- ()	0.0421	
Yes 77 (78.77) 22 (22.22)	Yes	77 (78.77)	22 (22.22)		
No 77(64.63) 44(36.36)	No	77(64.63)	44 (36.36)		
Type of cancer 0.12	Type of cancer			0.12	
Breast cancer 30 (17.73) 11 (83.26)	Breast cancer	30 (17.73)	11 (83.26)		
Gastrointestinal cancer 53(23.60) 35(77.39)	Gastrointestinal cancer	53(23.60)	35(77.39)		
Liver cancer 8(73 72) 3(27 27)	Liver cancer	8(73.72)	3(27.27)		
Ling cancer 14 (78 77) 4 (22 22)		14 (78 77)	4(22.22)		
Others 49 (03 79) 13 (07 20)	Others	49 (03 79)	13 (97 20)		
Honsehold size	Household size	(61,0)	5(57.20)	< 0.001	
1-3 68(77.95) 3(23.4)	1-3	68 (77.95)	3(23.4)		
4-6 77(6.61) 48(4.38)	4 - 6	77 (6.61)	48 (4.38)		
> 6 9(5.37) 15(5.62)	> 6	9 (5.37)	15 (5.62)		

^aValues are expressed as No. (%).



Figure 1. Catastrophic health expenditures (CHEs, %) for cancer treatment across the quintiles of household income.

the healthcare system has improved healthcare services, it has not been successful in reaching the goal discussed in upstream documents related to health in iran

Generally, high-cost cancer treatment services can impose a heavy financial burden on households and push them into poverty (31). Therefore, improved protection of patients and their families against financial difficulties is essential, especially for incurable diseases, such as cancer. Regarding the determinants of CHEs for cancer treatment, the current study showed that supplementary health insurance had an insignificant protective effect on CHEs and that the risk of catastrophic expenditures was lower in households with supplementary insurance (61.54%) compared to households without this type of insurance (70.53%). Moreover, Lee and Yoon indicated that the probability of facing CHEs among households without this insurance was higher than those covered by this insurance plan (32). Overall, the success of such plans in preventing CHEs depends on their sufficient coverage or effective definition of cancer care.

The current study revealed that the type of insurance is a significant determinant of exposure to CHEs. Accordingly, the incidence of CHEs was higher in households covered by social security and medical service insurance compared to those covered by the Relief Foundation insurance. Since people with the Relief Committee insurance are normally in the low-income group, they ignore receiving healthcare services due to their low payment capacity. In this regard, a previous study reported inconsistent results with the current study, and the CHEs were found to be higher in people covered by the Relief Foundation insurance because of their lower CTP (18).

Moreover, there was a significant association between the incidence of household CHEs and the type of cancer treatment. The results showed that the rate of exposure to CHEs was higher in households with breast cancer and lung cancer. Patients who received radiotherapy were less likely to be exposed to CHEs compared to patients receiving chemotherapy and radiotherapy simultaneously. In the study by Kavosi et al., it was found that 74.1% of patients undergoing chemotherapy were exposed to CHEs, while the rate of exposure to CHEs was 54.5% in patients undergoing radiotherapy. In line with other studies on household CHEs for cancer treatment (18, 33), a significant relationship was found between catastrophic expenditures and the sex of the household head.

The present study indicated that the rate of exposure to catastrophic expenditures was lower among households

whose head had academic education compared to other educational levels; this study confirmed the results of previous studies (34, 35). According to our findings, younger household heads were more exposed to CHEs compared to other age groups. This finding is in line with the results of previous studies (35, 36), as younger household heads often have a lower economic status, which is a good explanation for our findings.

The current study also revealed the positive effect of inpatient service utilization on the exposure of households to CHEs, as inpatient service utilization can increase the use of services provided for patients and consequently, increase the expenditure of medical services. Kavosi et al. (23) and Cruz Rivero et al. (36) also found that increased inpatient service utilization was associated with the frequency of hospitalization and the risk of CHEs. The household size is another factor, which had a negative significant relationship with the possibility of CHEs in the study population; in other words, CHEs decrease by increasing the household size.

The negative impact of increasing the household size on the likelihood of exposure to catastrophic expenditures may be due to the lower expectations of larger households to utilize health services and more attention to health by small-sized households. Finally, the CHEs increase in higher income households, because they allocate a larger share of their CTP to health expenditures. However, poorer households with lower income and payment capacity ignore receiving health services and prefer meeting other needs to health needs; a similar study in China confirmed our finding (37).

This study, similar to other studies, had some limitations. Due to differences in the socioeconomic status, the results of the present study may be only generalizable to other regions of the country. Besides, this study was conducted through interviews and had limitations, such as cost reporting bias and recall bias. Another limitation of the current study was restriction of the FFCI to distinguish between the progressive and regressive trends of the financing system; to resolve this issue, other indices are needed.

5.1. Conclusions

The present results showed that the CHEs were high for cancer treatment. Despite the efforts made so far, the Iranian health system has failed in preventing the catastrophic expenditures of cancer treatment and has not been successful in protecting households from the financial burden. Also, the recent health system transformation plan and health financing system reforms in Iran have not had significant effects on improving this index for these households. Therefore, a large number of households with cancer patients face financial difficulties due to health expenditures. Accordingly, more financial support and effective insurance plans need to be provided for these households, and the health system should not only improve supportive policies by expanding the insurance coverage, but also introduce cancer benefit packages to reduce cancer treatment expenditures.

Footnotes

Authors' Contribution: Study concept and design: A. R., and S. N.; analysis and interpretation of data: A. R., and P. H.; drafting of the manuscript: P. H.; critical revision of the manuscript for important intellectual content: B. D., N. Y., and P. H.; statistical analysis: P. H.

Conflict of Interests: The authors declare that they have no competing interests.

Ethical Approval: This article is part of a research project (Grant No: 95-04-163-29776, Ethical Code: IR.IUMS.REC1395.95-04-163-29776).

Funding/Support: The research team appreciates the financial support of the research deputy affiliated to Iran University of Medical Sciences.

References

- Vandenberghe D, Albrecht J. The financial burden of noncommunicable diseases in the European Union: a systematic review. *Eur J Public Health*. 2020;**30**(4):833–9. doi: 10.1093/eurpub/ckz073. [PubMed: 31220862].
- Buigut S, Ettarh R, Amendah DD. Catastrophic health expenditure and its determinants in Kenya slum communities. *Int J Equity Health*. 2015;14:46. doi: 10.1186/s12939-015-0168-9. [PubMed: 25971679]. [PubMed Central: PMC4438568].
- Kolesar RJ, Erreygers G, van Dam W, Chea V, Choeurng T, Leng S. Hardship Financing, Productivity Loss, and the Economic Cost of Illness and Injury in Cambodia [Preprint]. HAL Open Science. 2021. Available from: https://hal.archives-ouvertes.fr/hal-03437399/.
- Zhou Y, Wushouer H, Vuillermin D, Guan X, Shi L. Does the universal medical insurance system reduce catastrophic health expenditure among middle-aged and elderly households in China? A longitudinal analysis. *Eur J Health Econ*. 2021;22(3):463–71. doi: 10.1007/s10198-021-01267-3. [PubMed: 33582893].
- Khammarnia M, Setoodehzadeh F, Ansari-Moghaddam A, Barfar E, Zanganeh Baygi M, Peyvand M. Household financial contribution to the health system after Iran's Health Transformation Plan. *Rural Remote Health*. 2020;20(1):5495. doi: 10.22605/RRH5495. [PubMed: 32069064].
- Haakenstad A, Coates M, Marx A, Bukhman G, Verguet S. Disaggregating catastrophic health expenditure by disease area: cross-country estimates based on the World Health Surveys. *BMC Med.* 2019;**17**(1):36. doi: 10.1186/s12916-019-1266-0. [PubMed: 30755209]. [PubMed Central: PMC6372998].
- Shumet Y, Mohammed SA, Kahissay MH, Demeke B. Catastrophic Health Expenditure among Chronic Patients Attending Dessie Referral Hospital, Northeast Ethiopia. *Clinicoecon Outcomes Res.* 2021;**13**:99– 107. doi: 10.2147/CEOR.S291463. [PubMed: 33568923]. [PubMed Central: PMC7868221].

- Swetha NB, Shobha S, Sriram S. Prevalence of catastrophic health expenditure and its associated factors, due to out-of-pocket health care expenses among households with and without chronic illness in Bangalore, India: a longitudinal study. *J Prev Med Hyg.* 2020;61(1):E92-7. doi: 10.15167/2421-4248/jpmh2020.61.1.1191. [PubMed: 32490274]. [PubMed Central: PMC7225654].
- 9. Puteh SEW, Almualm Y. Catastrophic health expenditure among developing countries. *Health Syst Policy Res*. 2017;4(1):0.
- Singh PK. Histone methyl transferases: A class of epigenetic opportunities to counter uncontrolled cell proliferation. *Eur J Med Chem.* 2019;166:351–68. doi: 10.1016/j.ejmech.2019.01.069. [PubMed: 30735901].
- Alamzadeh Z, Beik J, Mirrahimi M, Shakeri-Zadeh A, Ebrahimi F, Komeili A, et al. Gold nanoparticles promote a multimodal synergistic cancer therapy strategy by co-delivery of thermo-chemo-radio therapy. *Eur J Pharm Sci.* 2020;**145**:105235. doi: 10.1016/j.ejps.2020.105235. [PubMed: 31991226].
- Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin.* 2018;**68**(6):394–424. doi: 10.3322/caac.21492. [PubMed: 30207593].
- 13. Boyle P, Levin B. *World cancer report 2008*. IARC Press, International Agency for Research on Cancer; 2008.
- Dye C, Reeder JC, Terry RF. Research for universal health coverage. *Sci Transl Med.* 2013;5(199):199ed13. doi: 10.1126/scitranslmed.3006971. [PubMed: 23966297].
- Bauer UE, Briss PA, Goodman RA, Bowman BA. Prevention of chronic disease in the 21st century: elimination of the leading preventable causes of premature death and disability in the USA. *Lancet.* 2014;**384**(9937):45–52. doi: 10.1016/S0140-6736(14)60648-6. [PubMed: 24996589].
- Wagstaff A, Flores G, Hsu J, Smitz MF, Chepynoga K, Buisman LR, et al. Progress on catastrophic health spending in 133 countries: a retrospective observational study. *Lancet Glob Health*. 2018;6(2):e169–79. doi: 10.1016/S2214-109X(17)30429-1. [PubMed: 29248367].
- Rezapour A, Ghaderi H, Azar FE, Larijani B, Gohari MR. Effects of health out-of-pocket payment on households in iran; catastrophic and impoverishment: Population based study in Tehran (2012). *Life Sci J.* 2013;10(3):1457-69.
- Kavosi Z, Delavari H, Keshtkaran A, Setoudehzadeh F. Catastrophic health expenditures and coping strategies in households with cancer patients in Shiraz Namazi hospital. *Middle East J Cancer.* 2014;5(1):13– 22.
- Piroozi B, Mohamadi-Bolbanabad A, Moradi G, Safari H, Ghafoori S, Zarezade Y, et al. Incidence and Intensity of Catastrophic Healthcare Expenditure for Type 2 Diabetes Mellitus Care in Iran: Determinants and Inequality. *Diabetes Metab Syndr Obes*. 2020;**13**:2865-76. doi: 10.2147/DMSO.S263571. [PubMed: 32922053]. [PubMed Central: PMC7450415].
- Kasahun GG, Gebretekle GB, Hailemichael Y, Woldemariam AA, Fenta TG. Catastrophic healthcare expenditure and coping strategies among patients attending cancer treatment services in Addis Ababa, Ethiopia. *BMC Public Health*. 2020;20(1):984. doi: 10.1186/s12889-020-09137-y. [PubMed: 32571275]. [PubMed Central: PMC7310089].
- Roshandel G, Ghanbari-Motlagh A, Partovipour E, Salavati F, Hasanpour-Heidari S, Mohammadi G, et al. Cancer incidence in Iran in 2014: Results of the Iranian National Population-based Cancer Registry. *Cancer Epidemiol.* 2019;**61**:50–8. doi: 10.1016/j.canep.2019.05.009. [PubMed: 31132560].
- Braveman P, Gruskin S. Poverty, equity, human rights and health. Bull World Health Organ. 2003;81(7):539–45. [PubMed: 12973647]. [PubMed

Central: PMC2572503].

- 23. Kavosi Z, Rashidian A, Pourmalek F, Majdzadeh S, Pourreza AG, Mohammad K, et al. [Measuring household exposure to catastrophic health care expenditures: A Longitudinal study in Zone 17 of Tehran]. *Hakim Res J.* 2009;**12**(2):38–47. Persian.
- Ravangard R, Jalali FS, Bayati M, Palmer AJ, Jafari A, Bastani P. Household catastrophic health expenditure and its effective factors: a case of Iran. *Cost EffResourAlloc*. 2021;**19**(1):59. doi: 10.1186/s12962-021-00315-2. [PubMed: 34530840]. [PubMed Central: PMC8444555].
- 25. Xu K. Distribution of health payments and catastrophic expenditures methodology. World Health Organization; 2005.
- Darvishi A, Amini-Rarani M, Mehrolhassani MH, Yazdi-Feyzabadi V. Fairness in household financial contribution to the Iran's healthcare system from 2008 to 2018. *BMC Res Notes*. 2021;**14**(1):190. doi: 10.1186/s13104-021-05606-8. [PubMed: 34001249]. [PubMed Central: PMC8130119].
- Doshmangir I, Hasanpoor E, Abou Jaoude GJ, Eshtiagh B, Haghparast-Bidgoli H. Incidence of Catastrophic Health Expenditure and Its Determinants in Cancer Patients: A Systematic Review and Metaanalysis. *Appl Health Econ Health Policy*. 2021;**19**(6):839–55. doi: 10.1007/s40258-021-00672-2. [PubMed: 34318445].
- Aryankhesal A, Etemadi M, Mohseni M, Azami-Aghdash S, Nakhaei M. Catastrophic Health Expenditure in Iran: A Review Article. *Iran J Public Health*. 2018;47(2):166–77. [PubMed: 29445626]. [PubMed Central: PMC5810379].
- Daneshkohan A, Karami M, Najafi F, Matin BK. Household catastrophic health expenditure. *Iran J Public Health*. 2011;40(1):94–9. [PubMed: 23113061]. [PubMed Central: PMC3481728].
- Murray CJ, Knaul F, Xu K, Musgrove P, Kawabata K. Defining and Measuring Fairness in Financial Contribution to the Health System. In: Murray CJL, Knaul F, Xu K, Musgrove P, Kawabata K, editors. *Defining and Measuring Fairness of Financial Contribution*. 24. World Health Organization; 2000.
- Hoang VM, Pham CP, Vu QM, Ngo TT, Tran DH, Bui D, et al. Household Financial Burden and Poverty Impacts of Cancer Treatment in Vietnam. *Biomed Res Int.* 2017;2017:9350147. doi: 10.1155/2017/9350147.
 [PubMed: 28904976]. [PubMed Central: PMC5585616].
- Lee M, Yoon K. Catastrophic Health Expenditures and Its Inequality in Households with Cancer Patients: A Panel Study. *Processes*. 2019;7(1). doi: 10.3390/pr7010039.
- Kimman M, Jan S, Yip CH, Thabrany H, Peters SA; Action Study Group, et al. Catastrophic health expenditure and 12-month mortality associated with cancer in Southeast Asia: results from a longitudinal study in eight countries. *BMC Med*. 2015;**13**:190. doi: 10.1186/s12916-015-0433-1. [PubMed: 26282128]. [PubMed Central: PMC4539728].
- Ghiasvand H, Sha'baninejad H, Arab M, Rashidian A. Hospitalization and Catastrophic Medical Payment: Evidence from Hospitals Located in Tehran. Arch Iran Med. 2014;17(7):0.
- Moradi Y, Nasehi M, Asadi-Lari M, Khamseh ME, Baradaran HR. The relationship between social capital components and control of type 2 diabetes: A path analysis model. *Med J Islam Repub Iran*. 2017;31(1):119– 23. doi: 10.18869/mjiri.31.21.
- Cruz Rivero C, Luna Ruiz GA, Morales Barrera R, Coello Levet CG. Catastrophic expenditure in health and income elasticities by item of expenditure in health services in Mexico. *Well-Being and Social Policy*. 2006;2(1):47–69.
- You X, Kobayashi Y. Determinants of out-of-pocket health expenditure in China: analysis using China Health and Nutrition Survey data. *Appl Health Econ Health Policy*. 2011;9(1):39-49. doi: 10.2165/11530730-00000000-00000. [PubMed: 21174481].