



# Dengue Fever: The Threat of Emerging Diseases Coinciding the Corona Crisis in Southeastern Iran

Jalil Nejati <sup>1</sup>, Mehdi Zanganeh Baygi <sup>2</sup> and Ruben Bueno-Mari <sup>3,4,\*</sup>

<sup>1</sup>Health Promotion Research Center, Zahedan University of Medical Sciences, Zahedan, Iran

<sup>2</sup>Department of Public Health, Health Promotion Research Center, School of Health, Zahedan University of Medical Sciences, Zahedan, Iran

<sup>3</sup>Department of Research and Development, Laboratorios Lokimica, Paterna, Spain

<sup>4</sup>Parasite and Health Research Group, Department of Pharmacy, Pharmaceutical Technology and Parasitology, Faculty of Pharmacy, University of Valencia, Burjassot, Spain

\*Corresponding author: Department of Research and Development, Laboratorios Lokimica, Paterna, Spain. Email: [rbueno@lokimica.es](mailto:rbueno@lokimica.es)

Received 2022 January 11; Revised 2022 March 05; Accepted 2022 April 27.

**Keywords:** Dengue, *Aedes*, Iran

## Dear Editor,

Dengue fever is one of the main mosquito-borne diseases worldwide, the preferred regions for which are the warm and tropical climates. This viral infection encompasses a broad spectrum of symptoms, from extremely mild to severe cases requiring medical intervention and hospitalization. Although it has no specific treatment, the symptoms can be managed. According to the WHO, dengue fever accounted for 5.2 million persons in 2019. Its prevalence rate increased as much as eightfold over the last two decades. The reported deaths increased fourfold in 2015 (4032) in comparison with 2000 (960) (1).

*Aedes aegypti* (Linnaeus, 1762) and *Ae. albopictus* (Skuse, 1895) are considered the main vectors of dengue fever. Because of choosing small artificial habitats for egg-laying, they are known as container-breeding mosquitos. This behavioral feature enables them to reproduce even in arid areas. This virus can be transmitted to the next generation through the eggs of an infected adult mosquito. In other words, by the transovarial transmission, several consecutive generations are infected if the female parent blood-feeds from the infected host. Moreover, unlike malaria vectors, their eggs are drought resistant, allowing them to move around easily (2). The detection of *Aedes albopictus* in the United States, followed by the transported cargo ships of discarded automobile tires from eastern Asia had occurred based on this mentioned biological characteristic (3). In the case of exposure to water as a stimulus, the eggs are hatched, and the larval development is done quickly (2). The aforementioned pointscan further highlights the importance of dengue fever and its vectors.

It is concluded that increasing global trade and trans-

port can facilitate the global distribution of dengue fever and its vectors from the native areas in Southeast Asia. Due to the potential of both species to adapt to new environments, they could establish themselves in many countries.

Several countries in the Eastern Mediterranean region have reported imported and indigenous cases of dengue fever. The first collection of adult *Ae. albopictus* was recorded in Southeastern Iran in 2013. However, subsequent studies have not confirmed its establishment (4). Although the likelihood of its presence in the study area has been predicted based on modelling, this species seemingly have failed to adapt to the climatic conditions of these areas (5).

Similarly, *Ae. aegypti* was collected from Western Iran (Khuzistan) in 1921 and Southern Iran (Bushehr) in 1953; however, the species has not been recorded in Iran for above 70 years (6). Although some studies still confirm the absence of this species in Southern Iran, several studies have reported its collection (7, 8). In this regard, the misidentification of the species is possible, mainly if a valid key is not used, concerns have raised about the presence and establishment of this important vector in the southern parts of the country.

Sistan and Baluchestan Province is located in Southeastern Iran, the south parts of which have a climatic condition similar to the Hormazagn Province, where monsoon streams cause heavy precipitation (4). According to some documents, dengue epidemics and transmission often peak after rainfall (1). This province has a long shared border with Pakistan and Afghanistan in the east and the Oman Sea in the south (4). The east neighbor, Pakistan, has been facing dengue outbreaks for years. It was first reported in Punjab in 1982 with 12 dengue seropositive pa-

tients. The last notable dengue outbreak in this country, accounting for 47,120 confirmed dengue cases and 75 deaths, was reported in 2019. The published papers indicate the establishment of *Ae. aegypti* and *Ae. albopictus* in this country (9).

On the other hand, there is international trade with Southeast Asia via through Chabahar Port. As another trading partner of Iran, India is tackling with dengue fever, and its goods are being exchanged via this port. These cases highlights the significance of the south of the province regarding the presence and occurrence of dengue and arouse concerns in the national health system.

Sistan and Baluchestan province is involved with many vector-borne diseases such as malaria, leishmaniasis, and Crimean-Congo hemorrhagic fever, leading to the establishment of a vector surveillance system, equipment, and trained personnel (4). In this regard, the use of a malaria entomological surveillance platform can improve dengue entomological surveillance. According to some modeling studies, the northern areas of the country with vast forests and similarities to the natural biotopes of *Ae. albopictus* are more prone to the establishment of this virus (10). Nevertheless, it seems that there is not an extensive and capable infrastructure for entomological surveillance in this area like southern parts. Under such a critical conditions, the widespread distribution of the coronavirus crisis has involved all staff, even medical entomology technicians.

Like other countries, Iran is involved in the COVID-19 pandemic. This crisis has imposed a lot of costs, logistics, and efforts; hence, the vector surveillance system as well as quick intervention and vector control response can be affected in such cases. This virus can also reduce the quality and quantity of entomological field studies. Moreover, due to physicians and health workers' involvement in the treatment, control, and vaccination program of COVID-19, most training topics were related to this disease as a priority. At the executive level of healthcare, less attention has been paid to the issue of dengue since the beginning of the coronavirus epidemic across the country.

Accordingly, future researchers are recommended to improve the detection of dengue fever cases, assess the knowledge and attitudes of physicians, health officers, and entomological technicians, and hold training and retraining courses on dengue fever and its vector. Moreover, reinforcing the dengue vectors surveillance system and their control equipment is suggested.

## Footnotes

**Authors' Contribution:** All authors have contributed to preparing this manuscript. They read the manuscript and

confirmed the final version.

**Conflict of Interests:** The authors declare that they have no conflicts of interest.

**Funding/Support:** This study received no funding.

## References

1. World Health Organization. *Dengue and severe dengue*. World Health Organization; 2022. Available from: <https://www.who.int/news-room/fact-sheets/detail/dengue-and-severe-dengue>.
2. Faraone J, Fischer S, Aponte CA, Etchepare E, Stechina OS, Stein M. Hatching pattern and coexistence of *Aedes aegypti* and *Aedes albopictus* (Culicidae) in a subtropical city, Argentina, after three decades of coexistence. *Acta Trop*. 2021;**218**:105885. [PubMed ID: 33713628]. <https://doi.org/10.1016/j.actatropica.2021.105885>.
3. Smith GC, Bozz ICJ, Moore CG. Importation of *Aedes Albopictus* and Other Exotic Mosquito Species into The United States in Used Tires from Asia. *J Am Mosq Control Assoc*. 1988;**4**(2):138–42.
4. Nejati J, Zaim M, Vatandoost H, Moosa-Kazemi SH, Bueno-Mari R, Azari-Hamidian S, et al. Employing Different Traps for Collection of Mosquitoes and Detection of Dengue, Chikungunya and Zika Vector, *Aedes albopictus*, in Borderline of Iran and Pakistan. *J Arthropod Borne Dis*. 2020;**14**(4):376–90. [PubMed ID: 33954211]. [PubMed Central ID: PMC8053067]. <https://doi.org/10.18502/jad.v14i4.5275>.
5. Nejati J, Bueno-Mari R, Collantes F, Hanafi-Bojd AA, Vatandoost H, Charrahy Z, et al. Potential Risk Areas of *Aedes albopictus* in South-Eastern Iran: A Vector of Dengue Fever, Zika, and Chikungunya. *Front Microbiol*. 2017;**8**:1660. [PubMed ID: 28928720]. [PubMed Central ID: PMC5591785]. <https://doi.org/10.3389/fmicb.2017.01660>.
6. Azari-Hamidian S. Checklist of Iranian mosquitoes (Diptera: Culicidae). *J Vector Ecol*. 2007;**32**(2):235–42. [PubMed ID: 18260513]. [https://doi.org/10.3376/1081-1710\(2007\)32\[235:coimdc\]2.0.co;2](https://doi.org/10.3376/1081-1710(2007)32[235:coimdc]2.0.co;2).
7. Jaberhashemi SA, Azari-Hamidian S, Soltani A, Azizi K, Dorzaban H, Norouzi M, et al. The Fauna, Diversity, and Bionomics of Culicinae (Diptera: Culicidae) in Hormozgan Province, Southern Iran. *J Med Entomol*. 2022;**59**(3):987–96. [PubMed ID: 35134206]. <https://doi.org/10.1093/jme/tjac003>.
8. Dorzaban H, Soltani A, Alipour H, Hatami J, Hashemi SAJ, Shahriari-Namadi M, et al. Morphological and molecular-based identification of *Aedes aegypti* (Diptera: Culicidae), a main vector of Dengue Fever, the first record in Iran after decades. *Research Square*. 2020:1–16. <https://doi.org/10.21203/rs.3.rs-20236/v1>.
9. Mukhtar MU, Han Q, Liao C, Haq F, Arslan A, Bhatti A. Seasonal Distribution and Container Preference Ratio of the Dengue Fever Vector (*Aedes aegypti*, Diptera: Culicidae) in Rawalpindi, Pakistan. *J Med Entomol*. 2018;**55**(4):1011–5. [PubMed ID: 29462424]. <https://doi.org/10.1093/jme/tjy010>.
10. Kamal M, Kenawy MA, Rady MH, Khaled AS, Samy AM. Mapping the global potential distributions of two arboviral vectors *Aedes aegypti* and *Ae. albopictus* under changing climate. *PLoS One*. 2018;**13**(12):e0210122. [PubMed ID: 30596764]. [PubMed Central ID: PMC6312308]. <https://doi.org/10.1371/journal.pone.0210122>.