



Long-Lasting Symptoms in COVID-19 Patients After Hospitalization

Sayyed Gholam Reza Mortazavi Moghaddam^{1,*}, Sara Soltani² and Gholamreza Sharifzadeh³

¹Division of Pulmonary, Department of Internal Medicine, Vali-e-Asre Hospital, Birjand University of Medical Sciences, Birjand, Iran

²Department of Internal Medicine, Vali-e-Asre Hospital, Birjand University of Medical Sciences, Birjand, Iran

³Epidemiology Department of Social Medicine, School of Health and Nutrition, Birjand University of Medical Sciences, Birjand, Iran

*Corresponding author: Division of Pulmonary, Department of Internal Medicine, Vali-e-Asre Hospital, Birjand University of Medical Sciences, Birjand, Iran. Email: gmortazavi@yahoo.com

Received 2022 November 14; Revised 2023 January 15; Accepted 2023 March 19.

Abstract

Background: Patients with COVID-19 may experience symptoms for a long time.

Objectives: The aim of this study is to determine the prevalence of COVID-19 related problems after discharge from the hospital.

Methods: This cross-sectional study was conducted on 194 hospitalized COVID-19 patients (110 [56.7%] men and 84 [43.3%] women) using the census method in 2021-2022. The patients were followed up for 4 weeks.

Results: The mean age and body mass index (BMI) of the patients were 57.57 ± 16.79 years and 25.9 ± 4.64 kg/m². The major complaint was fever (75.3%), followed by dyspnea (62%), general weakness (60.8%), cough (59.3%), and anorexia (49%). On admission, lung high-resolution computed tomography (HRCT) scans were normal in 51 patients (26.3%). Regardless of the underlying disease, at least 1 symptom was present in 63 (32%) of patients at the end of the study. Increased appetite was observed in 16 (8.2%) patients at discharge and thereafter. Hair loss was reported in 2% at the beginning and 8% at the end of the study.

Conclusions: Prolonged symptoms in COVID-19 patients are common and require long-term care.

Keywords: COVID-19, Fatigue, Dyspnea

1. Background

COVID-19 was first reported in Wuhan (China) (1). Respiratory and non-respiratory symptoms of COVID-19 patients may not subside during hospitalization. The duration of symptoms in COVID-19 patients is usually 2 weeks for mild cases and 3-6 weeks for severe cases (2).

The clinical problems may continue long after the acute stage of the disease. Depending on the severity of the disease, they are called "long-COVID-19 syndrome" (symptomatic for 4 to 12 weeks) or "post-COVID-19 syndrome" (symptomatic for more than 12 weeks) (3).

2. Objectives

Studying COVID-19 patients after discharge can help better manage post-COVID-19 or long-COVID-19 syndrome. In addition, COVID-19 imposes costs on the healthcare system, which can be better managed by identifying the conditions and consequences of the disease (4).

3. Methods

This cross-sectional study was conducted on 194 COVID-19 patients hospitalized in Vali-e-Asr Hospital (Birjand Uni-

versity of Medical Sciences, Birjand, Iran) during 2021 - 2022. This study was approved by the Ethics Committee of Birjand University of Medical Sciences (code: IR.BUMS.REC.1399.529). All hospitalized COVID-19 patients with positive polymerase chain reaction (PCR) tests were eligible for the study. Exclusion criteria were death during hospitalization, unwillingness to participate, and transfer to the intensive care unit (ICU) due to critical conditions. A questionnaire was prepared by the researcher and approved by experts. After discharge, patients were followed up weekly for 4 weeks. The data were recorded weekly, entered into SPSS version 19 (SPSS Inc, Chicago, IL, USA), and analyzed using descriptive statistics.

4. Results

A total of 194 COVID-19 patients (110 [56.7%] men and 84 [43.3%] women) were enrolled in this study from 2021 to 2022. The mean age and body mass index (BMI) of the patients were 57.57 ± 16.79 years and 25.9 ± 4.64 kg/m². There was a history of inhalant addiction in 8 (4.1%), oral addiction in 3 (1.5%), and smoking in 3 (1.5%) patients. The most common underlying diseases were hypertension (64; 33%), diabetes mellitus (44; 22.7%), dyslipidemia (30; 15.5%),

Table 1. The Prevalence of COVID-19-Related Symptoms in Sequential Weekly Evaluation ^a

Clinical Presentation	On Admission	At Discharge	1 Week Later	2 Weeks Later	3 Weeks Later	4 Weeks Later
Asthenia	118 (60.8)	103 (53.1)	75 (37.8)	47 (24.2)	19 (10)	16 (8.3)
Anorexia	95 (49)	55 (28.4)	36 (18.6)	20 (10.3)	4 (2.1)	6 (3.1)
Myalgia	63 (32.5)	18 (9.3)	10 (5.2)	4 (2.1)	4 (2.1)	4 (2.1)
Nausea, vomiting, diarrhea	36 (18.5)	8 (4.1)	8 (4.1)	2 (1)	1 (0.5)	1 (0.5)
Headache	24 (12.4)	8 (4.1)	5 (2.6)	2 (1)	1 (0.5)	1 (0.5)
Anosmia	7 (3.6)	3 (1.5)	3 (1.5)	0 (0)	0 (0)	0 (0)
Sore throat	6 (3.1)	1 (0.5)	1 (0.5)	1 (0.5)	1 (0.5)	1 (0.5)
Dizziness	6 (3.1)	4 (2.1)	3 (1.5)	1 (0.5)	1 (0.5)	1 (0.5)
Hair loss	2 (1)	3 (1.5)	7 (3.6)	9 (4.6)	5 (2.6)	8 (4.1)
Forgetfulness and delirium	2 (1)	3 (1.5)	4 (2.1)	2 (1)	1 (0.5)	0 (0)
Chillness	48 (24.7)	4 (2.1)	1 (0.5)	0 (0)	1 (0.5)	0 (0)
Dyspnea	120 (62)	89 (46)	65 (33.5)	45 (23.2)	21 (10.8)	12 (6.2)
Cough	113 (59.3)	87 (44.8)	62 (32)	43 (22.2)	20 (10.3)	11 (5.7)
Chest pain	54 (27.8)	34 (17.5)	15 (7.7)	6 (3.1)	4 (2.1)	2 (1)

^a Values are expressed as No. (%).

cardiovascular diseases (13; 6.7%), and pulmonary disorder (14; 6.6%). Less common disorders were thyroid disease (7; 3.6%), epilepsy/multiple sclerosis/convulsions (5; 2.6%), kidney disease (7; 3.6%), and thalassemia (0.5%), lymphoma (0.5%), psoriasis (0.5%), tuberculosis (0.5), and prostate disease (0.5%). The most common complaints of COVID-19 patients in sequential weekly follow-ups are presented in [Table 1](#). Increased appetite after recovery from COVID-19 was reported in 9.7% of patients. On admission, lung high-resolution computed tomography (HRCT) scans were normal in 51 patients (26.3%). The percentage of lung engagement was 5%-25%, 25%-50%, 50%-75%, and more than 75% in 37 (19%), 40 (20%), 43 (22%), and 23 (12%) of patients, respectively. Dexamethasone and remdesivir were prescribed for 166 (85.6%), beta interferon for 119 (61.3), and favipiravir for 30 (15.5%) patients.

5. Discussion

Thirty-two percent of patients were symptomatic at the end of 4 weeks of follow-up. The most common residual complaint was asthenia, followed by dyspnea, cough, anorexia, headache, smell or taste disturbance, throat discomfort, hair loss, and chest pain.

COVID-19-related symptoms may continue long after the acute stage of the disease, which are classified as post-COVID-19 syndrome or long-COVID-19 syndrome (3). Typical clinical symptoms of long-COVID-19 syndrome are cough, shortness of breath, fatigue/weakness, headache,

autonomic disorder, loss of smell or taste, confusion, muscle pains, and sometimes low fever. In general, the most common remaining symptoms after the acute stage of COVID-19 are dyspnea and weakness (5).

One of the symptoms that increased in prevalence in each follow-up round was hair loss compared to the previous round. In a study conducted in Brazil, hair loss was the most common symptom in post-COVID-19 syndrome, especially in men. It was related to the duration of the disease in the acute stage and its severity (6).

In our study, 44 patients (22.7%) had diabetes. The prevalence of diabetes among patients with COVID-19 in China was reported to be 20% (7). In Italy, the prevalence of diabetes in hospitalized COVID-19 patients was 8.9%, which was relatively higher than the overall local diabetes prevalence (6.2%) (8).

There were few patients with a history of lung diseases, such as asthma and chronic obstructive pulmonary disease (COPD), in our study. In a study conducted by Vimercati et al. on hospital staff at the University Hospital of Bari, it was found that obesity and underlying respiratory diseases (such as asthma and COPD) were important risk factors in the persistence of symptoms after 35 days (9).

One of the valuable tools to evaluate COVID-19 patients is the HRCT severity score. Vijayakumar et al. did not report a difference between the rate of dyspnea 3 months and 1 year after COVID-19, according to initial CT abnormalities. They also stated that there was no correlation between the mMRC (Modified Medical Research Council) dyspnea score

and HRCT severity score 3 months after the acute phase of COVID-19 (10).

5.1. Conclusions

Since prolonged COVID-19-related symptoms are relatively common; it is recommended that COVID-19 patients be followed up after hospital discharge. This follow-up should be done regardless of an underlying disease.

Acknowledgments

The authors would like to thank the Research Committee of Birjand University of Medical Sciences, all participants, and Vali-e-Asr Hospital staff.

Footnotes

Authors' Contribution: Sayyed Gholam Reza Mortazavi-Moghaddam: Planning, project, observer of the process, writing the article; Gholam Reza Sharifzadeh: Statistic analysis and writing the article; Sara Soltani: Data sampling and processing.

Conflict of Interests: Funding or Research support: The authors are not financially supported by any organization. Employment: The authors have not been employed in an institution that earns points from the researched topic in the last 5 years. Personal financial interests: The authors of the article do not have a financial or non-financial motivation for the research. Stocks or shares in companies: The authors of the article are not involved in any kind of company that benefits from the topic. Consultation fees: The authors of the article have not received any consulting rights from any institution. Patents: The authors have not received any patent rights. Personal or professional relations with organizations and individuals (parents and children, wife and husband, family relationships, etc.): The authors' family members are not related to this research. Unpaid membership in a government or non-governmental organization; The authors do not have any membership

in the Unpaid government or non-governmental organization related to the research. Are you one of the editorial board members or a reviewer of this journal? No.

Ethical Approval: IR.BUMS.REC.1399.529.

Funding/Support: This study did not receive financial support.

References

1. Cevik M, Kuppalli K, Kindrachuk J, Peiris M. Virology, transmission, and pathogenesis of SARS-CoV-2. *BMJ*. 2020;**371**:m3862. [PubMed ID: 33097561]. <https://doi.org/10.1136/bmj.m3862>.
2. Wu L, Wu Y, Xiong H, Mei B, You T. Persistence of Symptoms After Discharge of Patients Hospitalized Due to COVID-19. *Front Med (Lausanne)*. 2021;**8**:761314. [PubMed ID: 34881263]. [PubMed Central ID: PMC8645792]. <https://doi.org/10.3389/fmed.2021.761314>.
3. No authors listed. Long COVID: let patients help define long-lasting COVID symptoms. *Nature*. 2020;**586**(7828):170. [PubMed ID: 33029005]. <https://doi.org/10.1038/d41586-020-02796-2>.
4. Zafar M. Impact of the COVID-19 on the Health System and Healthcare Workers: A Systematic Review. *Health Scope*. 2022;**11**(3). <https://doi.org/10.5812/jhealthscope-123211>.
5. Yong SJ. Long COVID or post-COVID-19 syndrome: putative pathophysiology, risk factors, and treatments. *Infect Dis (Lond)*. 2021;**53**(10):737-54. [PubMed ID: 34024217]. [PubMed Central ID: PMC8146298]. <https://doi.org/10.1080/23744235.2021.1924397>.
6. Muller-Ramos P, Ianhez M, Silva de Castro CC, Talhari C, Criado PR, Amante Miot H. Post-COVID-19 hair loss: prevalence and associated factors among 5,891 patients. *Int J Dermatol*. 2022;**61**(5):e162-4. [PubMed ID: 35080250]. <https://doi.org/10.1111/ijd.16041>.
7. Guan WJ, Zhong NS. Clinical Characteristics of Covid-19 in China. Reply. *N Engl J Med*. 2020;**382**(19):1861-2. [PubMed ID: 32220206]. <https://doi.org/10.1056/NEJMc2005203>.
8. Fadini GP, Morieri ML, Longato E, Avogaro A. Prevalence and impact of diabetes among people infected with SARS-CoV-2. *J Endocrinol Invest*. 2020;**43**(6):867-9. [PubMed ID: 32222956]. [PubMed Central ID: PMC7103097]. <https://doi.org/10.1007/s40618-020-01236-2>.
9. Vimercati L, De Maria L, Quarato M, Caputi A, Gesualdo L, Migliore G, et al. Association between Long COVID and Overweight/Obesity. *J Clin Med*. 2021;**10**(18). [PubMed ID: 34575251]. [PubMed Central ID: PMC8469321]. <https://doi.org/10.3390/jcm10184143>.
10. Vijayakumar B, Tonkin J, Devaraj A, Philip KEJ, Orton CM, Desai SR, et al. CT Lung Abnormalities after COVID-19 at 3 Months and 1 Year after Hospital Discharge. *Radiology*. 2022;**303**(2):444-54. [PubMed ID: 34609195]. [PubMed Central ID: PMC8515207]. <https://doi.org/10.1148/radiol.202111746>.