



Determination of Influencers' Characteristics in the Health Sector

Mohammadreza Pourkarim ¹, Shahnaz Nayeبزadeh ^{1,*}, Seyed Moayed Alavian ² and Seyyed Hassan Hataminasab ¹

¹Department of Business Management, Yazd Branch, Islamic Azad University, Yazd, Iran

²Middle East Liver Diseases Center, Tehran, Iran

*Corresponding author: Department of Business Management, Yazd Branch, Islamic Azad University, P. O. Box: 89195/155, Yazd, Iran. Email: snayebzadeh@iauyazd.ac.ir

Received 2023 August 29; Revised 2023 September 14; Accepted 2023 September 17.

Abstract

Background: Modern marketing strategies, accompanied by tools such as social media, mobile platforms, and applications, play a significant role in discovering the healthcare sector's needs and desires of target audiences. Additionally, the emergence of influential individuals, known as influencers, in domains such as marketing, public health promotion, physical and mental well-being, sexual health, beauty, and lifestyle is expanding daily. The inspiration and motivation generated by such influencers tend to offer an attractive opportunity for effective communication with the audience.

Objectives: This study aims to extract and identify indicators and factors of influencer marketing in the health domain.

Methods: This study employed a mixed-methods approach, combining qualitative and quantitative analyses. Literature in the field was reviewed, coded, and subjected to analysis. Subsequently, influencer marketing indicators were extracted and screened. Based on the identified factors, a questionnaire consisting of 26 questions was developed and distributed as a survey among a study population of 1126 individuals related to medical and non-medical sciences. The extracted factors were then examined through exploratory factor analysis and confirmatory factor analysis. A structural factor model was developed, and the fit of the model and the number of factors were assessed.

Results: A total of 26 influencer indicators were identified and categorized into 5 factors: Active and skill-related features, infrastructure actions, consequential features, inherent and intrinsic features, and behavioral characteristics in the health domain. The identified factors were validated through exploratory factor analysis (EFA) and confirmed through confirmatory factor analysis. In addition, the fit model was approved.

Conclusions: The study revealed that establishing international and local influencers within the health domain, incorporating influencer indicators and features alongside an analysis of environmental conditions, plays a pivotal role in shaping the healthcare system and fostering public health promotion across countries.

Keywords: Influencer, Marketing, Social Media, Public Health, World Health Organization

1. Background

1.1. Internet and Social Media

The ever-increasing use of the internet and its penetration rate, alongside the emergence of digital innovations, has transformed human societies and lifestyles (1). The development of the internet, computers, and smartphones has led to substantial changes in audiences' and customers' expectations, preferences, and behaviors. Additionally, the emergence of advanced technologies such as artificial intelligence, blockchain, and virtual reality has automated systems, fostered integration, and ultimately brought about further transformations in human behavior (2).

Social media and their role in fostering communication have transformed various sectors, markets, and segments of societies, including the healthcare sector (3). Using diverse social media platforms such as Facebook, Twitter, Telegram, WhatsApp, and TikTok and incorporating innovations like artificial intelligence into the abovementioned platforms has created new opportunities within societies. Factors such as content sharing, engagement rates, and individuals' interactions with posts, images, videos, and animations have influenced education and awareness among intended audiences (4).

The power and cost-effectiveness of social media seem to be the important advantages of this type of

communication media. Recent research indicates that over 75% of adults use various social media platforms to access health-related information (5). Existing studies have explored the role of social media in disease management, increasing interpersonal communication and interaction, social learning, and various forms of social support (6). Social media can be a crucial and strategic factor in raising awareness, behavior control, reinforcement, and behavior modification by establishing new behavioral models (7, 8). Shaping new behaviors is important in knowledge dissemination and sharing experiences within these media (9). Moreover, countries' use of social media platforms enables the exploitation of vast communication networks, which can, in turn, enhance the effectiveness and cost moderation of delivering messages (6). Examining the advantages and challenges of using social media in different countries can lead to further optimization. For instance, the role social media played during the recent pandemic in achieving successful patterns, including prevention, vaccination, screening, and education, exemplifies the effective use of this platform (10).

1.2. Social Media and the Health Sector

Countries have targeted the formulation of investment and research and development models, utilizing new approaches in community education. Correct utilization of interactive tools (social media) has been considered a successful example in strengthening healthcare systems, enhancing social and political commitments, community mobilization, and coordinating public programs (11). Online relationships through social networks have evolved and have been capable of fostering mutual interests, trust, and reinforcement of behavioral norms (12). International research shows that children, adolescents, and young adults engage with health information through their favorite websites and social networks (13). The future of many sectors in the health domain relies heavily on the direct integration of new technologies in hardware and software, as well as the application of these digital innovations. Proper use of these technologies, alongside experts and decision-makers in the realm of prevention, diagnosis, and treatment, plays a significant role in the performance of public health care (14). Recently, public health institutions have repeatedly employed social media platforms to advertise educational resources and content for public awareness. For example, social media platforms promote healthy dietary regimes, lifestyle improvements, preventing epidemics, blood donation and other campaigns, vaccination event promotions, and other health-related initiatives (15). In addition, identifying

existing channels for information gathering, sharing, and exchange plays a key role in implementing health promotion activities (16). Campaigns can also be used to determine target audiences and tailor messages to their specific characteristics. It must be noted that campaigns with defined topics and objectives have the potential to alter attitudes (17). Governments can significantly drive controlling factors and influence individuals' health through online tools, particularly social media platforms and public campaigns. Stakeholders and decision-makers aim to target the dynamism and interactions of individuals in these platform communities by disseminating accurate content and establishing new behavioral patterns (10). It is worth mentioning that the structure of social networks and the nature of these media can serve as information providers and advertisers or create competition regarding transmission, learning, and dissemination capabilities (18).

1.3. Social Media Influencers and the Health Domain

The community of social media influencers expands and possesses considerable potential and influential power over their audience (19). Influencers can be described as thought leaders in the media (20). Due to their credibility and distinct position, influencers or influential individuals serve as supporters and trusted figures in social media, acting as progressive agents (21). Influencers in social networks, channels, platforms, and applications, utilizing the power of media, play a significant role in human societies. Utilizing these individuals can contribute to conveying health-centered messages (22). The success of worldwide health programs relies on aligning goals, implementing strategies, and using tools correctly and innovatively. Coordination and alignment of strategies with culture, social norms, public awareness levels, financial resources, and the levels of engagement with actors and stakeholders are vital for any initiative. The emergence of influencer phenomena and influencer marketing has created attractive opportunities for accessing and engaging with audiences (23). Influencer marketing is a form of native advertising that can promote various communication techniques to create promotional messages (24). Identifying influencers on social media can significantly affect target audiences (21). The interest of advertisers in influencer marketing is steadily growing, giving rise to increased budgeting and expenses for investing in influencers (25). Influencers can stimulate, inspire, and motivate audiences by engaging them with posts, comments, likes, hashtags, etc. (26). Influencer marketing is a strategic marketing process that involves researching, identifying, engaging, and supporting individuals who can generate influential conversations

and interactions with audiences about a brand or behavior on social media (27). Collaborating with these influencers provides new opportunities for interacting with audiences and offers useful applications for modeling health behaviors, promoting proper behavioral norms, and eliminating risky behaviors or breaking taboos (23). Most healthcare brands can effectively engage their audience by providing education, findings, news, and new preventive and therapeutic techniques in an attractive and informative manner. It should be noted that establishing and facilitating cultural aspects appears to be a significant challenge in this field (28). Since the internet inhibits many influencers, celebrities, bloggers, athletes, musicians, etc., it is quintessential to reduce this clutter and categorize and identify suitable influencers at the right time (29). In the realm of interdisciplinary studies and the integration of sciences, there has been no effort to identify influencers in specific medical areas or explore the characteristics of influencers globally and in Iran, particularly in healthcare and medical services. Since countries have diverse cultures and population segments, it is crucial to properly examine geographical regions, residents' awareness levels, cultures, religions, and traditions to develop successful strategies through influencers and regional and global programs and strategies. Influencers can play a pivotal role in the success or failure of such programs (30). The presence of influencers in international programs, including those designed by the World Health Organization, can significantly contribute to cost reduction.

2. Objectives

This study aimed to identify specific characteristics of influencer marketing in the healthcare sector and categorize the most important ones.

3. Methods

This study was approved by the Ethical Committee of Yazd Branch, Islamic Azad, University, Yazd, Iran (IR.IAU.YAZD.REC.1401.065. Approval date: 2022-12-28).

The methodology of this study involved both qualitative and quantitative analysis. Initially, a total of available articles published between 2015 and 2022 encompassing marketing and influencer keywords were reviewed. Subsequently, to extract influencer and influencer marketing characteristics, specifically in the healthcare field, the texts were coded and analyzed using NVIVO software (version 12). The identified factors were then used to develop a questionnaire with 26

questions. The questionnaire was distributed among 1,126 individuals through online Google Forms and paper-based surveys. Purposeful sampling was used to select responses, followed by simple random sampling. The study population included individuals in both medical and non-medical sciences (Appendix 1 and Figure 1). To categorize the indicators and identify the factors, factor analysis (FA), specifically exploratory factor analysis (EFA), was performed using SPSS software (version 27).

Exploratory factor analysis is employed to identify and validate the main dimensions within the proposed constructs. In this analysis, researchers aim to examine experimental data to discover and identify indicators and their relationships without imposing any specific model. In other words, exploratory analysis is used when sufficient prior evidence does not exist to form a hypothesis about the number of underlying factors in the data. It allows researchers to explore the data to determine the number and nature of factors that explain the interrelationships among variables.

Subsequently, using SPSS AMOS software (version 23), confirmatory factor analysis (CFA) and structural equation modeling (SEM) was conducted to assess the factor structure validation, the number of factors, and the fit of the factor structures. In confirmatory factor analysis, researchers aim to validate a specific factor structure. Hypotheses are clearly defined regarding the number of factors, and the fit of the hypothesized factor structure with the covariance structure of the measured variables is rigorously tested. In confirmatory factor analysis, the goal is to create a model that describes, explains, or justifies experimental data using a relatively small number of parameters. This model is developed based on prior knowledge and empirical information about the data structure.

3.1. Descriptive Statistics

This section of the statistical analysis focuses on examining the distribution of the sample in terms of variables such as gender, marital status, age, work experience, and education level (Figure 1).

4. Results

In the present study, 1126 individuals replied to our request and completed the questionnaire. Of these, 58.3% were female, and 41.7% were male. Notably, more than half of the participants (656) were female. Furthermore, 56% were unmarried. Regarding age, 31% of the sample was between 26 and 35. Regarding age, 31% of the sample were between 26 and 35. Regarding the education




Figure 1. Demographic profile of study participants: A, gender distribution; B, marital status; C, different age groups of participants; D, educational level of individuals; E, work experience related to the academic degree; F, daily usage; and G, type of social media activities of participants.

level, 507 respondents (45%) held a bachelor's degree, 342 (30.4%) held a master's degree, 146 individuals (13%) held a PhD, 104 respondents (9.2%) were general practitioners, 24 (2.1%) were specialists, and 3 respondents (0.3%) were

subspecialists. In terms of work experience, more than half of the study population, i.e., 582 subjects (52%) had less than 5 years of work experience, 190 (17%) between 5 to 10 years, 122 individuals (11%) between 10 and 15 years, and 232

(20%) had more than 15 years of relevant work experience in their field of study.

Moreover, approximately 41% of the participants ($n = 459$) used social media platforms for two to four hours per day, and approximately 64% ($n = 717$) of the respondents used social media as entertainment and for leisure activities in cyberspace.

4.1. Exploratory Factor Analysis

Based on the results of the initial phase of the research, a set of 26 questions was extracted and formulated into a structured questionnaire. An exploratory factor analysis was conducted, and it was observed that all 26 questions had factor loadings greater than 0.40 for each factor. Accordingly, none of the items were excluded from the analysis, and the exploratory factor analysis was performed, yielding the following results.

The first step in exploratory factor analysis is assessing the sample adequacy. The results of the Kaiser-Meyer-Olkin (KMO)-Bartlett test indicated that the KMO measure (0.902) exceeded 0.60, indicating the adequacy of the samples. The significance level of zero for Bartlett's test also suggested the appropriateness of the factor model. Based on these results, the exploratory factor analysis can be conducted on the collected data. Additionally, the principal component analysis with the varimax rotation method was employed in all stages of the analysis. The extracted variance showed that the eigenvalue for 5 factors was greater than 1.0. Therefore, the proposed factor structure would consist of 5 factors, which account for more than 60.82% of the cumulative variance.

Next, the final factor structure was presented in the rotated matrix, which included the contribution of variables in explaining the factors. Each element in this table indicates the partial correlation between the respective structure (row) and the rotated factor (column) (Appendix 2).

As observed, the factor loading values in the final rotated matrix indicate that the items are loading significantly on one factor only, with each item having a factor loading above 0.40. Therefore, based on the results obtained from the rotated matrix, the factors can be labeled as shown in Table 1.

4.2. Confirmatory Factor Analysis

Although exploratory data analysis has hitherto been conducted successfully, confirmatory factor analysis needs to be undertaken to ensure the validity and meaningfulness of the analysis. The software AMOS was used for this purpose (Figure 2).

The significance of the estimated standardized coefficients in the final model is presented in Table 2.

As shown in Table 2, all relationships are statistically significant (the significance level of the test statistic for each relationship was less than 0.05). Furthermore, since all factor loadings in Table 2 are greater than 0.40, it can be inferred that the items effectively explain the construct, and the construct (variables) exhibit factorial and construct validity.

Various fit indices were calculated to assess the model fit, indicating a close fit between the model and the data. The chi-square value divided by the degrees of freedom was close to 3. Additionally, the RMSEA value was 0.045, which was less than 0.08, and indices such as Tucker Lewis index (TLI), incremental fit index (IFI), relative fit index (RFI), normed fit index (NFI), and comparative fit index (CFI) were all above 0.90. Therefore, the model demonstrated a good fit and was hence validated.

Furthermore, influencer attributes such as expressive ability (62.3%), being honest with the audience (60.2%), creativity (57.9%), content production (53.8%), self-confidence (52.8%), adherence to professional principles (51.9%), physical appearance (44.4%), work experience in a specific field (38%), were identified as having the highest frequency in the influential factor of health domain (with a wide spectrum) examined by the statistical population in this study.

Additionally, having relationships with stakeholders (47.9%), noticeable presence in the community (43.9%), accessibility (43%), being relatable to the audience (41.6%), expertise in a specific field (41.3%), educational level (40.7%), acceptable work resume (40.6%), number of followers (40.1%), presence on media platforms (38.3%), uniformity of profile and coloring (37.7%), number of views (37.5%), participation in various events (37.3%), number of likes (35.3%), scientific resume (34.6%), were extracted with the highest frequency and a wide range of responses.

Additionally, gender (36%) and number of comments (35.1%) were extracted with a moderate spectrum of responses in terms of audience perception. A flowchart illustrating the entire study package is included as a new figure in the manuscript (Figure 3).

5. Discussion

Social media has become increasingly popular as a tool for digital marketing. Identifying and communicating with active individuals on social media, often called 'influencers,' in various fields, especially the health sector, requires a series of tips and advice for ordinary people. On the other hand, health organizations employ their influencers to disseminate health messages and implement their programs. Our study aimed to discover a more effective method of identifying genuine influencers

Table 1. Number of Determined Factors

Agents	Name	Symbol
Factor 1	Active and skillful characteristic	SKIL
Factor 2	Infrastructure actions	INFR
Factor 3	Consequence	CONS
Factor 4	Intrinsic and latent characteristic	ATRI
Factor 5	Behavioral characteristic	BEHA

Table 2. The Meaningfulness of Confirmatory of the Model Coefficients

Variables	Coefficient Standard	Coefficient Non-standard	SE	CR	P-Value
Intrinsic and latent characteristic					
Q1	0.722	1.000	-	-	-
Q2	0.434	0.601	0.048	12.388	0.000
Q3	0.685	0.993	0.051	19.355	0.000
Q4	0.723	1.035	0.052	20.057	0.000
Q5	0.520	0.769	0.051	15.205	0.000
Active and skillful characteristic					
Q6	0.696	1.000	-	-	-
Q7	0.760	1.005	0.045	22.206	0.000
Q8	0.758	0.948	0.043	21.824	0.000
Q9	0.810	1.079	0.045	23.809	0.000
Q10	0.694	0.894	0.044	20.527	0.000
Q11	0.568	0.764	0.045	17.072	0.000
Behavioral characteristic					
Q12	0.460	1.000	-	-	-
Q13	0.561	1.210	0.075	16.169	0.000
Q14	0.695	1.663	0.127	13.074	0.000
Q15	0.727	1.741	0.132	13.152	0.000
Q16	0.585	1.364	0.112	12.191	0.000
Infrastructure actions					
Q17	0.524	1.000	-	-	-
Q18	0.553	0.991	0.062	16.025	0.000
Q19	0.887	1.782	0.097	18.361	0.000
Q20	0.797	1.609	0.090	17.878	0.000
Q21	0.739	1.516	0.091	16.707	0.000
Q22	0.740	1.702	0.099	17.229	0.000
Consequence					
Q23	0.842	1.000	-	-	-
Q24	0.894	1.111	0.030	36.871	0.000
Q25	0.892	1.138	0.033	33.985	0.000
Q26	0.847	1.065	0.031	34.292	0.000

Abbreviations: SE, standard; CR, composite reliability.




Figure 2. The structural model in standard (corrected) form

in the health sector. We conducted surveys among both specialized and non-specialized individuals in the health sector. The collected data were meticulously analyzed, leading to significant results. To discuss our objectives and findings, it is important to address some preliminary points.

Different regions worldwide, including the Middle East, increasingly embrace healthcare organizations

(31). Generating innovative ideas and providing novel services in healthcare institutions and sectors are both important and essential. Creating a value chain in health organizations and developing a strategy is important. They are also decisive in evaluating the services of organizations in the public and private sectors (32). In preparing, implementing, and evaluating strategic plans, a competitive advantage is formed through close




Figure 3. Flowchart illustrating the step-by-step methodology employed in the study for data collection and analysis.

communication with contacts and efforts to activate social programs in providing services to patients and contacts, which secures the success of strategic plans (33). These institutions utilize digital marketing platforms and tools to establish their brand in the minds of their audience. Through this digital approach, social media platforms contribute to managing communication with healthy audiences and patients by providing online services and disseminating medical information to healthcare institutions. Topics such as sexual health, physical fitness, and mental well-being are abundant on these social media platforms, particularly in generation Z (born after 1981). Therefore, having a roadmap and planning for applying these tools seems crucial (34). YouTube, Instagram, Twitter, TikTok, etc., contain influencers who express positive and negative experiences, plans and personal goals, and even self-disclosures of their actions among their followers. Disclosure of personal life and experiences by influencers can build a strong relationship with followers and subsequently lead to a change in the attitudes of their audience (35).

Since the official COVID-19 pandemic declaration by the World Health Organization, public announcements such as daily notifications, prevention tips, vaccination offers, contact tracing, and screening have unprecedentedly highlighted the role of experts in the media (36). In this context, the presentation of textual and video content on topics such as COVID and statements such as “I have also been vaccinated” among influencers, celebrities,

and opinion leaders with their followers has become significantly widespread and effective (37). The adaptation of health subjects in social media and the enhancement of the effectiveness of social media in health management and disease prevention still require further improvement and particular caution (38, 39).

The necessity of hiring specialized influencers at the community health level is crucial because the abundant non-specialized and unofficial influencers can pose a threat to public health. Therefore, categorizing and screening influencers based on their expertise and target communities can lead to effective and safe practices (40). Influencers with marketing strategies, especially social marketing, can play a directing role in health (41). The main and fundamental factor in the presence of influencers is the ease of spreading and transferring their perspectives and attitudes to their audience (42, 43).

The impact of influencers on individuals' attitudes, with additional studies in psychology, anthropology, and cultural studies, can result in synergistic effects. The combination of digital tools, textual elements, visual effects, and learning prepares the audience for the persuasion phase to receive the message (44). Unfortunately, many influencers with dubious and unofficial sources, with a lack of scientific and specialized qualifications, or to profit through platforms and media, are spreading incorrect knowledge in various health fields (45). This issue has a visible, devastating impact on global and national public health management (46). Qualified

and approved health experts, on the other hand, are the most reliable and accurate information sources in the health field. Influencers, along with the adaptation of an influencer marketing approach in the health domain, can provide the ability to disseminate scientifically based information correctly to the general public. To address the importance of the topics mentioned above, creating and introducing approved influencers in various specialized health domains requires an understanding of influencer characteristics, factors, and indicators.

This study presents a general perspective on influencer marketing in the health field. Contrary to frequently available articles, we have focused on the preferred characteristics of health influencers based on input from experts and non-experts (47). Our research indicates that the most important priority for a health influencer is possessing skills and expertise in communication, education, creativity, confidence, and content creation. This result aligns with recent studies emphasizing the importance of education and related skills (45, 48). It is worth mentioning that factors such as the honesty of the influencer with the followers, the appearance and compliance with the professional work principles, and having relevant work experience in the specific field are desirable and accepted among the audience. Factors like honesty must be highlighted as part of ethics on social media (49), although the relevance of influencers' work experience continues to spark debates (50).

As the next priority, the communication and interaction of health influencers with stakeholders, including the trustees and decision-makers, and communication with organizations, managers, and experts is important. Also, influencers' active and dynamic availability, together with creating and inducing a sense of companionship and being an audience, are important factors in establishing a relationship with followers. This part of the results, which depicts a dynamic relation of influencers and bi-directional interactions with the audience, was already expected and discussed on different occasions in the literature (51, 52).

It is worth mentioning that health influencers should pay special attention to improving the level of education, work resumes, and scientific expertise in their activities. In addition, the use of correct techniques to retain and attract new followers and order in the integrity and coloring of the profile, the effort to increase the number of visits, and the techniques of receiving likes are also important factors in the characteristics of the influencer (52). These findings shed light on the necessity of strategic activities of influencers on social media (53, 54). It should be noted that presence in different media and platforms and participation in various meetings are effective factors for

being up-to-date and efficient in communicating with the audience (55). As discussed in the literature (56, 57), gender is a factor that can be further analyzed and investigated in completing the influencer indicators. For example, the field of sexual health, emerging diseases in societies, behavioral diseases, and the increasing trend of trans subjectivity can be discussed and investigated in more depth; more research should be done on the appropriate gender of the influencers in these fields to attract their target audience.

Based on the previous research and considering the need for expert influencers in the field of health, creating regional or international influencers with the discussed characteristics can assist health systems, including the World Health Organization, and the health systems of countries can benefit from them in influencing, transferring norms, publishing guidelines, training of desirable behaviors and warnings used in public health-related issues (15). Since different countries have various restrictions for accessing social media platforms, domestically tailored strategies by influencers on accessible platforms can assist the audience.

5.1. Conclusions

our study highlighted certain characteristics that influencers in public health should have. An influencer's honesty with their audience, physical appearance, adherence to professional principles, and relevant work experience in the field can lead to greater acceptance and approval among the target audience. Additionally, influencers' communication and interaction with stakeholders, including managers and health policy decision-makers, as well as engagement with organizations and experts concerned, were crucial points spotlighted by our study. The application of our study results in ongoing WHO initiatives, such as eliminating viral hepatitis, can be extremely influential. Accordingly, recruiting skillful, honest, and responsible influencers with relevant education in viral hepatitis, including virologists, gastroenterologists, and physicians, represents the most effective approach in this crucial worldwide health program. This approach can enhance the capabilities of health organizations to implement their programs in various societies by utilizing social media platforms to reach a wider audience.

Footnotes

Authors' Contribution: Study concept and design: Influencers in the health sector; acquisition of data: Mohammadreza Pourkarim; analysis and interpretation

of data: Mohammadreza Pourkarim; drafting of the manuscript: Mohammadreza Pourkarim; critical revision of the manuscript for important intellectual content: Shahnaz Nayebyzade and Seyed Moyed Alavian; statistical analysis: Mohammadreza Pourkarim; study supervision: Shahnaz Nayebyzade, Seyed Moayed Alavian, and Seyed Hassan Hataminasab.

Conflict of Interests: The authors declare no relevant conflict of interest.

Data Reproducibility: The dataset presented in the study is available on request from the corresponding author during submission or after publication.

Ethical Approval: This study was approved by the Ethical Committee of Yazd Branch, Islamic Azad, University, Yazd, Iran (IR.IAU.YAZD.REC.1401.065. Approval date: 2022-12-28).

Funding/Support: The authors received no financial support for this article's research, authorship, and publication.

References

- Gagnon K, Sabus C. Professionalism in a digital age: Opportunities and considerations for using social media in health care. *Phys Ther.* 2015;**95**(3):406-14. [PubMed ID: 24903111]. <https://doi.org/10.2522/ptj.20130227>.
- Kannan PK, Li H*. Digital marketing: A framework, review and research agenda. *Int J Res Mark.* 2017;**34**(1):22-45. <https://doi.org/10.1016/j.ijresmar.2016.11.006>.
- Zhang C, Gotsis M, Jordan-Marsh M. Social media microblogs as an HPV vaccination forum. *Hum Vaccin Immunother.* 2013;**9**(11):2483-9. [PubMed ID: 23842072]. [PubMed Central ID: PMC3981860]. <https://doi.org/10.4161/hv.25599>.
- Pourkarim M, Van Espen L, Thijssen M, Van Ranst M, Pourkarim MR. How adequate social media management supports the viral hepatitis elimination program. *Hepat Mon.* 2018;**18**(5):e69791. <https://doi.org/10.5812/hepatmon.69791>.
- Wong CA, Merchant RM, Moreno MA. Using social media to engage adolescents and young adults with their health. *Healthc (Amst).* 2014;**2**(4):220-4. [PubMed ID: 25984444]. [PubMed Central ID: PMC4433153]. <https://doi.org/10.1016/j.hjdsi.2014.10.005>.
- Lin X, Kishore R. Social media-enabled healthcare: A conceptual model of social media affordances, online social support, and health behaviors and outcomes. *Technol Forecast Soc Change.* 2021;**166**:120574. <https://doi.org/10.1016/j.techfore.2021.120574>.
- Tang Z, Miller AS, Zhou Z, Warkentin M. Does government social media promote users' information security behavior towards COVID-19 scams? Cultivation effects and protective motivations. *Gov Inf Q.* 2021;**38**(2):101572. [PubMed ID: 35719729]. [PubMed Central ID: PMC9188430]. <https://doi.org/10.1016/j.giq.2021.101572>.
- Gever VC, Talabi FO, Adelabu O, Sanusi BO, Talabi JM. Modeling predictors of COVID-19 health behaviour adoption, sustenance and discontinuation among social media users in Nigeria. *Telemat Inform.* 2021;**60**:101584. [PubMed ID: 36569993]. [PubMed Central ID: PMC9758446]. <https://doi.org/10.1016/j.tele.2021.101584>.
- Zhao J, Han H, Zhong B, Xie W, Chen Y, Zhi M. Health information on social media helps mitigate Crohn's disease symptoms and improves patients' clinical course. *Comput Hum Behav.* 2021;**115**:106588. <https://doi.org/10.1016/j.chb.2020.106588>.
- Abbas J, Wang D, Su Z, Ziapour A. The role of social media in the advent of COVID-19 pandemic: Crisis management, mental health challenges and implications. *Risk Manag Healthc Policy.* 2021;**14**:1917-32. [PubMed ID: 34012304]. [PubMed Central ID: PMC8126999]. <https://doi.org/10.2147/RMHP.S284313>.
- Howell J, Pedrana A, Schroeder SE, Scott N, Aufegger L, Atun R, et al. A global investment framework for the elimination of hepatitis B. *J Hepatol.* 2021;**74**(3):535-49. [PubMed ID: 32971137]. [PubMed Central ID: PMC7505744]. <https://doi.org/10.1016/j.jhep.2020.09.013>.
- Paige SR, Stellefson M, Chaney BH, Chaney JD, Alber JM, Chappell C, et al. Examining the relationship between online social capital and ehealth literacy: Implications for instagram use for chronic disease prevention among college students. *Am J Health Educ.* 2017;**48**(4):264-77. [PubMed ID: 29152031]. [PubMed Central ID: PMC5687578]. <https://doi.org/10.1080/19325037.2017.1316693>.
- Ahola Kohut S, LeBlanc C, O'Leary K, McPherson AC, McCarthy E, Nguyen C, et al. The internet as a source of support for youth with chronic conditions: A qualitative study. *Child Care Health Dev.* 2018;**44**(2):212-20. [PubMed ID: 29082537]. <https://doi.org/10.1111/cch.12535>.
- Fang C, An J, Bruno A, Cai X, Fan J, Fujimoto J, et al. Consensus recommendations of three-dimensional visualization for diagnosis and management of liver diseases. *Hepatol Int.* 2020;**14**(4):437-53. [PubMed ID: 32638296]. [PubMed Central ID: PMC7366600]. <https://doi.org/10.1007/s12072-020-10052-y>.
- Grajales FJ, Sheps S, Ho K, Novak-Lauscher H, Eysenbach G. Social media: a review and tutorial of applications in medicine and health care. *J Med Internet Res.* 2014;**16**(2):e13. [PubMed ID: 24518354]. [PubMed Central ID: PMC3936280]. <https://doi.org/10.2196/jmir.2912>.
- Green AM, Innes-Hughes C, Rissel C, Mitchell J, Milat AJ, Williams M, et al. Codesign of the population health information management system to measure reach and practice change of childhood obesity programs. *Public Health Res Pract.* 2018;**28**(3):e2831822. [PubMed ID: 30406261]. <https://doi.org/10.17061/phrp.2831822>.
- Perloff RM. Social media effects on young women's body image concerns: Theoretical perspectives and an agenda for research. *Sex Roles.* 2014;**71**(11-12):363-77. <https://doi.org/10.1007/s11199-014-0384-6>.
- Centola D. The spread of behavior in an online social network experiment. *Science.* 2010;**329**(5996):1194-7. [PubMed ID: 20813952]. <https://doi.org/10.1126/science.1185231>.
- Childers CC, Lemon LL, Hoy MG. #Sponsored #Ad: Agency perspective on influencer marketing campaigns. *J Curr Issues Res Advert.* 2018;**40**(3):258-74. <https://doi.org/10.1080/10641734.2018.1521113>.
- De Veirman M, Cauberghe V, Hudders L. Marketing through Instagram influencers: the impact of number of followers and product divergence on brand attitude. *Int J Advert.* 2017;**36**(5):798-828. <https://doi.org/10.1080/02650487.2017.1348035>.
- Lou C, Yuan S. Influencer marketing: How message value and credibility affect consumer trust of branded content on social media. *J Interact Advert.* 2019;**19**(1):58-73. <https://doi.org/10.1080/15252019.2018.1533501>.
- Krisam M, Altendorfer LM. [influencer marketing in healthcare: A review] [Bundesverband der Ärzte des Öffentlichen Gesundheitsdienstes]. *Gesundheitswesen.* 2023;**85**(2):100-2. Germany. [PubMed ID: 33706391]. <https://doi.org/10.1055/a-1377-6478>.
- Lutkenhaus RO, Jansz J, Bouman MP. Tailoring in the digital era: Stimulating dialogues on health topics in collaboration with social media influencers. *Digit Health.* 2019;**5**. [PubMed ID: 30729023]. [PubMed Central ID: PMC6350129]. <https://doi.org/10.1177/2055207618821521>.
- Wojdyski BW, Evans NJ. Going native: Effects of disclosure position and language on the recognition and evaluation of online native advertising. *J Advert.* 2015;**45**(2):157-68. <https://doi.org/10.1080/00913367.2015.1115380>.
- MoShi Q. *The impact of influencers on young people's buying decisions.* Oulu University of Applied Science; 2020.

26. Boerman SC, van Reijmersdal EA. Informing consumers about "hidden" advertising: A literature review of the effects of disclosing sponsored content. *Advertising in New Formats and Media*. 2016. p. 115–46. <https://doi.org/10.1108/978-1-78560-313-620151005>.
27. Condello I, Santarpino G. Considerations for influencer marketing in cardiac surgery and interventional cardiology. *Ann Thorac Surg*. 2021;**112**(2):689. [PubMed ID: 33352175]. <https://doi.org/10.1016/j.athoracsur.2020.09.076>.
28. Groeneveld PW, Yang L, Segal AG, Karaca-Mandic P, Kanter GP. The effects of market competition on cardiologists' adoption of transcatheter aortic valve replacement. *Med Care*. 2020;**58**(11):996–1003. [PubMed ID: 32947511]. [PubMed Central ID: PMC7572719]. <https://doi.org/10.1097/MLR.0000000000001391>.
29. Gillin P. New media, new influencers and implications for the public relations profession. *J new commun res*. 2008;**2**(2):1–10.
30. Pourkarim M, Nayebedeh S, Alavian SM, Hataminasab SH. Digital marketing: A unique multidisciplinary approach towards the elimination of viral hepatitis. *Pathogens*. 2022;**11**(6):626. [PubMed ID: 35745480]. [PubMed Central ID: PMC9228079]. <https://doi.org/10.3390/pathogens11060626>.
31. Agha A, Rashid A, Rasheed R, Khan S, Khan U. Antecedents of customer loyalty at telecomm sector. *Turk Online J Qual Inq*. 2021;**12**:1352–74.
32. Elrod JK, Fortenberry Jr JL. Formulating productive marketing communications strategy: a major health system's experience. *BMC Health Serv Res*. 2018;**18**(Suppl 3):926. [PubMed ID: 30545343]. [PubMed Central ID: PMC6293486]. <https://doi.org/10.1186/s12913-018-3676-7>.
33. Mubarik MS, Naghavi N, Mubarik M, Kusi-Sarpong S, Khan SA, Zaman SI, et al. Resilience and cleaner production in industry 4.0: Role of supply chain mapping and visibility. *J Clean Prod*. 2021;**292**:126058. <https://doi.org/10.1016/j.jclepro.2021.126058>.
34. Pfender EJ, Devlin MM. What do social media influencers say about birth control? A content analysis of YouTube vlogs about birth control. *Health Commun*. 2023;**38**(14):3336–45. <https://doi.org/10.1080/10410236.2022.2149091>.
35. Ferchaud A, Grzeslo J, Orme S, LaGroue J. Parasocial attributes and YouTube personalities: Exploring content trends across the most subscribed YouTube channels. *Comput Hum Behav*. 2018;**80**:88–96. <https://doi.org/10.1016/j.chb.2017.10.041>.
36. Zhou SL, Jia X, Skinner SP, Yang W, Claude I. Lessons on mobile apps for COVID-19 from China. *J Saf Sci Resil*. 2021;**2**(2):40–9. <https://doi.org/10.1016/j.jnlssr.2021.04.002>.
37. Bonnevie E, Smith SM, Kummeth C, Goldbarg J, Smyser J. Social media influencers can be used to deliver positive information about the flu vaccine: findings from a multi-year study. *Health Educ Res*. 2021;**36**(3):286–94. [PubMed ID: 34252187]. [PubMed Central ID: PMC8411386]. <https://doi.org/10.1093/her/cyab018>.
38. Zou W, Zhang WJ, Tang L. What do social media influencers say about health? A theory-driven content analysis of top ten health influencers' posts on sina weibo. *J Health Commun*. 2021;**26**(1):1–11. [PubMed ID: 33372857]. <https://doi.org/10.1080/10810730.2020.1865486>.
39. Heldman AB, Schindelar J, Weaver JB. Social media engagement and public health communication: Implications for public health organizations being truly "social". *Public Health Rev*. 2013;**35**(1):13. <https://doi.org/10.1007/bf03391698>.
40. Mathew V, Soliman M. Does digital content marketing affect tourism consumer behavior? An extension of technology acceptance model. *J Consum Behav*. 2020;**20**(1):61–75. <https://doi.org/10.1002/cb.1854>.
41. Uzir MUH, Al Halbusi H, Thurasamy R, Thiam Hock RL, Aljaberi MA, Hasan N, et al. The effects of service quality, perceived value and trust in home delivery service personnel on customer satisfaction: Evidence from a developing country. *J Retail Consum Serv*. 2021;**63**:102721. <https://doi.org/10.1016/j.jretconser.2021.102721>.
42. Breves P, Amrehn J, Heidenreich A, Liebers N, Schramm H. Blind trust? The importance and interplay of parasocial relationships and advertising disclosures in explaining influencers' persuasive effects on their followers. *Int J Advert*. 2021;**40**(7):1209–29. <https://doi.org/10.1080/02650487.2021.1881237>.
43. Johnson BK, Bradshaw AS, Davis J, Diegue V, Frost L, Hinds J, et al. Credible Influencers: Sponsored YouTube personalities and effects of warranting cues. *J Media Psychol*. 2021;**34**(4). <https://doi.org/10.1027/1864-1105/a000310>.
44. Farivar S, Wang F, Turel O. Followers' problematic engagement with influencers on social media: An attachment theory perspective. *Comput Hum Behav*. 2022;**133**:107288. <https://doi.org/10.1016/j.chb.2022.107288>.
45. Hendry NA, Hartung C, Welch R. Health education, social media, and tensions of authenticity in the 'influencer pedagogy' of health influencer Ashy Bines. *Learn Media Technol*. 2021;**47**(4):427–39. <https://doi.org/10.1080/17439884.2021.2006691>.
46. Byrne E, Kearney J, MacEvilly C. The role of influencer marketing and social influencers in public health. *Proc Nutr Soc*. 2017;**76**(OCE3):E103. <https://doi.org/10.1017/s002966517001768>.
47. Ouvrein G, Pabian S, Giles D, Hudders L, De Backer C. The web of influencers. A marketing-audience classification of (potential) social media influencers. *J Mark Manag*. 2021;**37**(13-14):1313–42. <https://doi.org/10.1080/0267257x.2021.1912142>.
48. Okuah O, Scholtz B, Snow B. A grounded theory analysis of the techniques used by social media influencers and their potential for influencing the public regarding environmental awareness. Conference of the South African Institute of Computer Scientists and Information Technologists (SAICSIT); 2019.
49. Wellman ML, Stoldt R, Tully M, Ekdale B. Ethics of authenticity: Social media influencers and the production of sponsored content. *J Media Ethics*. 2020;**35**(2):68–82. <https://doi.org/10.1080/23736992.2020.1736078>.
50. Gräve J. What kpis are key? Evaluating performance metrics for social media influencers. *Soc Media Soc*. 2019;**5**(3). <https://doi.org/10.1177/2056305119865475>.
51. Veale HJ, Sacks-Davis R, Weaver ER, Pedrana AE, Stooze MA, Hellard ME. The use of social networking platforms for sexual health promotion: identifying key strategies for successful user engagement. *BMC Public Health*. 2015;**15**:85. [PubMed ID: 25884461]. [PubMed Central ID: PMC4340797]. <https://doi.org/10.1186/s12889-015-1396-z>.
52. Hudders L, De Jans S, De Veirman M. The commercialization of social media stars: a literature review and conceptual framework on the strategic use of social media influencers. *Int J Advert*. 2020;**40**(3):327–75. <https://doi.org/10.1080/02650487.2020.1836925>.
53. Charest F, Bouffard J, Zajmovic E. Public relations and social media: Deliberate or creative strategic planning. *Public Relat Rev*. 2016;**42**(4):530–8. <https://doi.org/10.1016/j.pubrev.2016.03.008>.
54. Le K, Aydin G. Impact of the pandemic on social media influencer marketing in fashion: a qualitative study. *Qual Market Res Int J*. 2022;**26**(4):449–69. <https://doi.org/10.1108/qmr-11-2021-0133>.
55. Gupta S, Dash SB, Mahajan R. The role of social influencers for effective public health communication. *Online Inf Rev*. 2021;**46**(5):974–92. <https://doi.org/10.1108/oir-01-2021-0012>.
56. Rowley J, Johnson F, Saffi L. Gender as an influencer of online health information-seeking and evaluation behavior. *J Assoc Inf Sci Technol*. 2015;**68**(1):36–47. <https://doi.org/10.1002/asi.23597>.
57. Kawachi I, Berkman LF. Social ties and mental health. *J Urban Health*. 2001;**78**(3):458–67. [PubMed ID: 11564849]. [PubMed Central ID: PMC3455910]. <https://doi.org/10.1093/jurban/78.3.458>.