

The impact of Hepatitis C virus infection on kidney transplantation outcomes: A systematic review of 18 observational studies

Zohreh Rostami ^{1*}, Mohammad Hossien Nourbala ¹, Seyed Moayed Alavian ², Fatemeh Bieragh-dar ¹, Yunes Jahani ¹, Behzad Einollahi ¹

¹ Nephrology and Urology Research Center, Baqiyatallah University of Medical Sciences, Tehran, IR Iran

² Baqiyatallah Research Center for Gastroenterology and Liver Disease, Baqiyatallah University of Medical Sciences, Tehran, IR Iran

ARTICLE INFO

Article Type:
Review Article

Article history:
Received: 26 Apr 2010
Revised: 31 Jan 2011
Accepted: 28 Feb 2011

Keywords:
Hepatitis C infection
Kidney transplantation
Graft survival
Patient survival
Mortality
Natural history
Outcome assessment

ABSTRACT

Background: Hepatitis C virus (HCV) infection occurs in 0% to 51% of dialysis patients, and many HCV-positive patients are urged to undergo kidney transplantation. However, the outcome of renal transplantation in HCV-positive recipients is unknown.

Objectives: Our review aimed to address the outcomes of renal transplantation recipients (RTRs) following kidney transplantation.

Materials and Methods: We selected studies that used the adjusted relative risk (aRR) and 95% CI of all-cause mortality and graft loss in HCV-positive compared with HCV-negative RTRs as study endpoints. Cox proportional hazard analysis was used in all studies to calculate the independent effects of HCV infection on RTR outcomes. Sixteen retrospective cohort studies and 2 clinical trials were selected for our review. Sixteen studies were related to patient survival, and 12 examined graft survival.

Results: The combined hazard ratio in HCV-infected recipients was 1.69-fold (1.33-1.97, $p < 0.0001$) and 1.56 times (1.22-2.004, $p < 0.0001$) greater than that of HCV-negative recipients for mortality and graft loss, respectively.

Conclusions: Although HCV-infected RTRs have worse outcomes than HCV-negative RTRs, kidney transplantation is the preferred treatment for patients with HCV infection and end-stage renal disease.

© 2011 Kowsar M.P.Co. All rights reserved.

► Implication for health policy/practice/research/medical education:

HCV infection may negatively interfere on final outcomes of kidney transplantation. We strongly recommend reading this interesting article to all general practitioners, surgeons, nephrologists and urologists.

► Please cite this paper as:

Rostami Z, Nourbala MH, Alavian SM, Bieragh-dar F, Jahani Y, Einollahi B. The impact of hepatitis C virus infection on kidney transplantation outcome: A systematic review of 18 observational studies. *Hepat Mon.* 2011;11(4):247-54.

Background

Hepatitis C virus (HCV) infection is a common problem among dialysis patients and kidney transplant recipients (1). The Centers for Disease Control and Prevention (CDC) detects HCV infection by enzyme linked immunosorbent

assay (ELISA) in 8.1% (range 0% to 51%) of ESRD patients in large dialysis centers (2). Additionally, many HCV-positive patients are urged to undergo kidney transplantation (3). The major cause of mortality due to liver failure in kidney transplant recipients is HCV infection (4). The outcome of renal transplantation in HCV-positive recipients is unknown (2, 5); some studies have reported better survival in HCV-positive ESRD patients compared with those remaining on dialysis (1, 4, 6, 7).

A rise in viral load following immunosuppression in

* Corresponding author at: Zohreh Rostami, Nephrology and Urology Research Center, Baqiyatallah University of Medical Sciences, MollaSadra Ave., Vanak Sq., Tehran, IR Iran. Tel: +98-2181262073, Fax: +98-2188067114.
E-mail: rostami@ijnu.ir

HCV-positive kidney transplant recipients was suggested to be a significant cause of poor outcome (1, 4, 6, 8). Also, viral load and liver deterioration are related (8). Conversely, several surveys did not observe worse outcomes in HCV-positive renal transplant recipients (RTRs) when HCV infection was acquired before kidney transplantation, especially during the first 5-8 years (7).

However, a recent study from a US registry evaluated the effect of immunosuppressive regimens on survival in HCV-positive RTRs, demonstrating that antibody induction does not adversely affect patient survival (1, 7, 9). Moreover, cyclosporine (10) and mycophenolate mofetil (MMF) may have protective effects (1, 6) and inhibit HCV replication in renal transplant patients with HCV infection. Whether hepatitis virus infected-patients should stay on dialysis or be referred for kidney transplantation remains unknown.

Objectives

We performed a meta-analysis to determine the effects of HCV infection on outcomes in RT patients.

Materials and Methods

Search strategy

We searched electronic databases, including PubMed, the Cochrane Database of Systematic Reviews, EMBASE, and CINAHL, for studies from Jan 1981 to Jan 2010 to identify those that reported the effect of HCV infection on RTR outcomes. Our keywords included "hepatitis C," "HCV infection," "kidney transplantation," "graft survival," "patients survival," "mortality," "natural history," "outcome," and their synonyms. Two authors independently developed a search strategy to identify randomized trials and cohort studies that investigated the effect of HCV on patients and graft survival after kidney transplantation. To identify additional relevant articles, reference lists from qualitative topic reviews and the identified articles were also searched. Duplicate publications were discarded. We restricted our search to human studies and placed no restrictions on language.

Study Selection

The electronic and manual searches yielded 1137 papers by title and abstract, of which 149 were considered relevant and selected for a full text review. 131 irrelevant reports were excluded (Figure 1). After a full text review, 16 retrospective cohort studies (1, 11-25), and 2 clinical trials (26, 27) were selected for our review (Table 1). Sixteen studies were related to patient survival, and 12 examined graft survival. Study characteristics are summarized in Table 1.

Criteria for inclusion

Two independent reviewers assessed with a standard

method each included trial about adult kidney transplant recipients with HCV infection, defined as testing positive for anti-HCV or HCV RNA by polymerase chain reaction (PCR) in serum at the time of enrollment. Also participants were evaluated with regard to patient and kidney outcomes, which were defined as liver-related death and return to dialysis due to HCV infection. Discrepancies were resolved in conference. Other criteria for inclusion were controlled trials and cohort studies that reported patient and graft survival among HCV-infected RTRs. Table 1 shows the characteristics of the studies in this review. Studies that included HCV-infected donors were excluded. Between the trials included in our meta-analysis, there are a few differences in patients and graft outcome (Table 2). Thus, we decided to pool these data for evaluation.

Review questions and endpoints of interest

Our review aimed to answer two specific questions:

1. What is the effect of HCV infection on renal graft survival?
2. What is the effect of HCV infection on renal recipient survival?

All selected studies used the adjusted relative risk (aRR) and 95% CI of all-cause mortality and graft loss in HCV-positive versus -negative RTRs as study endpoints. Cox proportional hazard 5) (we have converted HR to RR with a formula) analysis was used in all studies to calculate independent effects of HCV infection on RTR outcomes after adjustments for potentially contributing factors, such as age, gender, follow-up period, type of transplant, diabetes mellitus, post-transplant plasma creatinine, race, duration of dialysis, donor death etiology, and proteinuria. First-generation enzyme-linked immunosorbent assay test before 1991, second generation until 1997 and third generation until now were used to detect HCV




Figure 1. Summary of literature search and study selection

Table 1. Study characteristics

Authors	country	Number of all patients/ HCV+ RTRs	Patient/graft survival reports (Yes/No)	Type of Study
Einollahi et al. (2003)(11)	Iran	1006/45	Yes/yes	Retrospective cohort
Luan et al. (2008)(1)	U.S	79337/3708	Yes/no	Retrospective cohort
Aroldi et al. (2005)(12)	Italy	541/209	Yes/yes	Retrospective cohort
Legendre et al. (1998)(13)	France	499/112	Yes/no	Retrospective cohort
Gentil et al. (1999)(14)	Spain	320/85	Yes/yes	Retrospective cohort
Lee et al. (2001)(15)	Taiwan	477/136	Yes/yes	Retrospective cohort
Breitenfeldt et al. (2002)(16)	Germany	927/123	Yes/yes	Retrospective cohort
Bruchfeld et al. (2004)(17)	Sweden	571/51	Yes/no	Retrospective cohort
Morales et al. (2004)(18)	Spain	3365/488	Yes/yes	Retrospective cohort
Ingsathit et al. (2007)(19)	Thailand	346/22	Yes/no	Retrospective cohort
Batty et al. (2001)(20)	U.S	28692/1624	Yes/no	Retrospective cohort
Mahmoud et al. (2004)(21)	Egypt	133/80	Yes/yes	Retrospective cohort
Lin et al. (2004)(22)	Taiwan	299/129	Yes/yes	Retrospective cohort
Ridruejo et al. (2007)(23)	Argentina	396/155	Yes/yes	Retrospective cohort
Gentil Govantes et al. (2009)(24)	Spain	5693/1053	No/yes	Retrospective cohort
Mitwalli et al. (2006)(25)	Saudi Arabia	448/286	No/yes	Retrospective cohort
Pereira et al. (1995)(26)	U.S	75/19	Yes/yes	Clinical trial
Pereira et al. (1998)(27)	U.S	103/23	Yes/yes	Clinical trial

infection. Further, serum HCVRNA (PCR) was examined in anti-HCV-positive patients for confirmation of HCV infection in 6 studies.

Statistical analysis

We pooled outcomes (mortality rates, renal allograft failure), which had been expressed as relative risk (RR) with 95% confidence intervals (CI), using STATA 8. The results of each outcome were analyzed for heterogeneity by Q test (the random effects method of Der Simonian-Laird). Funnel plots, Begg's rank correlation test, and Egger's regression asymmetry test were used to assess the existence of publication bias. The Forest plot was used to demonstrate the details of pooled analysis. Combined hazard ratios were assessed by sensitivity analysis.

Results

Description of Included Trials

The included studies are summarized in Table 1. Follow up duration and adjusted variables for each study shown in table 2 and adjusted relative risk for mortality and graft loss also presented in table 3. A total of 8348 HCV-infected RTRs before or after kidney transplantation were identified from 123,228 living and deceased RTRs, as reported in 18 studies. Pereira BJ *et al.* had 2 studies in different years. Data on 8 studies (Pereira BJ *et al.* study 1, Pereira BJ *et al.* study 2, Legendre C *et al.*, Gentil MA *et al.*, Lee WC *et al.*, Breitenfeldt *et al.*, Bruchfeld *et al.*, and Morales *et al.*) that were reported before 2005 were also used in a meta-analysis by Fabrizi *et al.* and Gentil MA *et al.* confirmed HCV infection detection by immunoblotting, and Bruchfeld (71%), Ridruejo (33.54%), Ingsathit (100%), Mitwalli (100%),

Table 2. Follow up and adjusting variables of included articles

Authors	year	Adjusted variables
Einollahi <i>et al.</i> (2003)(11)	1995-2001	Donor characteristic (age, source, gender, blood group) and recipient characteristic (age, gender, ESRD etiology, history of diabetes, blood group)
Luan <i>et al.</i> (2008)(1)	1995-2004	Recipient characteristics (age, sex, race, diabetes, renal diagnosis, time on dialysis, panel reactive antibody level, availability of private insurance) and donor characteristics (age, living donor, extended criteria donor, cold ischemia time, presence of hypertension, creatinine level, and cause of death)
Aroldi <i>et al.</i> (2005)(12)	1972-1989	age
Gentil <i>et al.</i> (1999)(14)	1986-1997	Donor characteristic (age, gender, time on dialysis, ESRD etiology, number of transplant, pre-transplant transfusion, peak and immediate pre-transplant immunization, number of HLA A+B and HLA DR mismatches, years of transplant, cold ischemia time, anti HCV Ab, pre-transplant clinical liver disease
Lee <i>et al.</i> (2001)(15)	1984-1999	Sex, mode of dialysis, duration of dialysis, diabetes, hypertension, HBV infection, HCV infection, liver function impairment, hepatoma
Breitenfeldt <i>et al.</i> (2002)(16)	1978-1994	HBeAg, HCV infection after transplantation, con-comitant HBV and HCV infection, occurrence of acute rejection, age at transplantation and time on dialysis, HBeAg positivity, HBsAg positivity, HCV infection after transplantation, age at transplantation and occurrence of acute rejection.
Bruchfeld <i>et al.</i> (2004)(17)	1989-1997	age, sex, diabetes, previous transplantations, type of transplant, and time in RRT for death, HCV, diabetes
Morales <i>et al.</i> (2004)(18)	1990-1994	year of transplant, recipient age, Last panel reactive antibodies, acute rejection, triglycerides, Creatinine, proteinurea
Ingsathit <i>et al.</i> (2001)(19)	3.7 year	acute rejection episode, recipient age, long duration of dialysis; diabetes mellitus, delayed graft function, and sex mismatch, Creatinine
Batty <i>et al.</i> (2001)(20)	1994-1997	age, race, gender, end-stage renal disease due to diabetes, weight, year of transplant, duration of pre-transplant dialysis, previous transplant, donor and recipient age, donor and recipient race, donor and recipient gender, delayed graft function, antibody induction therapy (combined and also analyzed separately for OKT3 and ALG), and allograft rejection
Mahmoud <i>et al.</i> (2004)(21)	1993-1995	donor and recipient age and sex, primary cause of ESRD, HLA mismatch, number of transplants, time on dialysis therapy, number of acute rejection episodes, presence of persistent proteinuria, and year of transplantation.
Lin <i>et al.</i> (2004)(22)	1981-2000	Recipient age and sex, donor age and sex, anti HCV Ab, chronic hepatitis, pre-transplant diabetes, , pre-transplant hypertension, pre-transplant coronary artery disease, HLA DR mismatch
Ridruejo <i>et al.</i> (2007)(23)	1991-2004	Age, anti-HCV, traditional immunosuppression, rejection
Gentil Govantes <i>et al.</i> (2009)(24)	1984-1989 1990-1995 1996-2001 2002-2007	sex and age of the recipient, diabetes as ESRD cause, retransplant status, duration of previous RRT, and transplant year, Transplant time period
Mitwalli <i>et al.</i> (2006)(25)	1980-2001	age, sex, blood pressure, type of donor, and immunosuppressive medication , type of donor (living related, living unrelated, and cadaver donors), hepatitis status, hepatitis-positive or hepatitis-negative
Pereira <i>et al.</i> (1998)(27)	1987-1990	Effects of hepatitis C infection and renal transplantation on survival in end-stage renal disease. The New England Organ Bank Hepatitis C Study Group

and Mahmoud (100%) of hemodialysis patients confirmed it by HCV-RNA (PCR). In response to our request, Einollahi *et al.* replied that nearly 70% of HCV positive antibody anti HCV antibody positive RTRs were confirmed by HCV RNA (PCR).

Effect on patient and graft survival

The Q-test for heterogeneity revealed $p < 0.0001$ ($Q =$

69.81, $df = 15$) and $p < 0.0001$ ($Q = 66.15$, $df = 11$) for patient and graft survival, respectively. Further, a meta-analysis was done with a random model showed a combined hazard ratio in HCV-infected recipients that was 1.69-fold (1.33-1.97, $p < 0.0001$) (Figure 2) and 1.56 times (1.22-2.004, $p < 0.0001$) (Figure 3) greater than in HCV-negative recipients for mortality and graft loss, respectively.




Figure 2. Hazard ratio in HCV-infected recipients for patient survival




Figure 3. Hazard ratio in HCV-infected recipients for graft survival

Publication bias

For patient and graft survival rates, publication bias was examined using Bagg and Manzumdar and Egger's regression asymmetry, both of which were non-significant [($p = 0.753$, $p = 0.226$; Figure 4) and ($p = 0.304$, $p = 0.55$; Figure 5), respectively]. Similar results were observed in the funnel plots.

Sensitivity analysis

All eligible studies included in meta-analysis. Because

the elimination of each study did not have an impact on the combined hazard ratio, the overall estimation was robust (Figure 6).

Discussion

Hepatitis C infection is a risk factor for graft loss and death in renal transplant recipients (8). Although our report and recent studies have emphasized the detrimental role of hepatitis C in long-term patient and graft survival after renal transplantation (10), several studies have demonstrated that patient and graft survival on-

Table 3. Adjusted relative risk for mortality and graft loss

Author	95% confidence interval	
	aRR for mortality	aRR for graft loss
Einollahi et al. (2003)(11)	4.308 (2.88-6.4)	2.609 (2.07-3.27)
Luan et al. (2008)(1)	1.3 (1.2-1.4)	N.A. ^a
Aroldi et al. (2005)(12)	1.65 (1.13-2.42)	1.4 (1.17-1.81)
Pereira et al. (1995)(26)	1 (0.49-2.02)	0.95 (0.54-1.67)
Pereira et al. (1998)(27)	2.6 (1.15-5.9)	1.3 (0.66-2.58)
Legendre et al. (1998)(13)	2.8 (1.4-5.7)	N.A. ^a
Gentil et al. (1999)(14)	3.1 (1.2-7.8)	3 (1.8-5)
Lee et al. (2001)(15)	1.57 (0.75-1.11)	1.25 (0.75-1.32)
Breitenfeldt et al. (2002)(16)	1.93 (1.01-3.42)	N.A. ^a
Bruchfeld et al. (2004)(17)	2.23 (1.48-3.34)	1.96 (1.37-2.79)
Morales et al. (2004)(18)	1.505 (1.12-2.02)	1.58 (1.27-1.97)
Ingsathit et al. (2007)(19)	1.59 (0.28-9.02)	N.A. ^a
Batty et al. (2001)(20)	1.23 (1.01-1.49)	N.A. ^a
Mahmoud et al. (2004)(21)	0.5 (0.1-1.9)	0.5 (0.3-1.2)
lin et al. (2004)(22)	0.3 (0.13-0.65)	0.8 (0.48-1.35)
Ridruejo et al. (2007)(23)	1.66 (1.01-2.77)	1.97 (1.18-3.29)
Gentil Govantes et al. (2009)(28)	N.A. ^a	1.5 (1.1-1.9)
Mitwalli et al. (2006)(25)	N.A. ^a	4.37 (1.8-4.8)

^a N.A.: Not Accessible

HCV infection after renal transplantation are the same in the short term compared with non-infected renal transplant patients (6). Conversely, kidney transplantation is a better option for HCV-positive ESRD patients versus remaining on dialysis (1). To better examine HCV-positive RTR outcomes, we performed a meta-analysis using observational studies that used adjusted data of all-cause mortality.

Impact on patient survival

Consistent with Fabrizi's meta-analysis, the aRR for mortality rate in our study was lower than in other studies (4, 8, 13, 14, 17), likely due to the greater sample size,

early detection, improvement in management, and exact follow-up. Compared with Fabrizi's meta-analysis, which included 8 articles, our study included 18 articles that comprised more than 123,000 RTRs, indicating that greater consideration has been given to the controversy of HCV-infected RTR outcomes and kidney transplantation in the past 5 years. Several studies have demonstrated lower patient and graft survival in HCV-positive RTRs, related in part to associated complications, such as cirrhosis, hepatocellular carcinoma, cardiovascular disease, diabetes mellitus, sepsis, higher PRA, and deceased kidney donation (1, 10).

Impact on graft survival:

In our study, the aRR for graft loss was similar to that in Fabrizi's meta-analysis. Although during the first 5-10 years, graft and patient survival was apparently similar between negative and positive HCV-infected RTRs (4), HCV-associated glomerulonephritis, proteinuria, and diabetic nephropathy can progress rapidly to chronic allograft nephropathy (6).

Role of other factors in mortality

It appears that the increased mortality in anti-HCV-positive patients was partially related to mortality due to causes other than HCV infection. According to a novel risk score for mortality in RTRs (29), the risk score for HCV (1.5) was not more than age above 40 years in comparison to younger than 40 (2.2-6.7), pre-transplant diabetes mellitus (1.8), post-transplant diabetes mellitus (1.5), serum creatinine levels at the first year after transplantation (1.7), and proteinuria greater than 1g during the first year of operation (2.7). In a recent meta-analysis, mortality due to liver complications, such as cirrhosis and hepatocellular carcinoma, among HCV-infected RTRs increased in most studies that were included, with an aRR of 1.79, compared with HCV-negative recipients (6). In a systematic review, cardiovascular and infectious diseases were also important causes of death in HCV-positive RTRs (6).

Because mortality and graft loss are multifactorial, we used the aRR that had been obtained by the Cox regression model in each study to appraise the isolated influence of HCV infection on patient and graft survival. In contrast to studies that reported a negative impact, the majority of studies that demonstrated a positive impact of transplantation on HCV-infected patient and graft survival rates did not use the Cox regression model; consequently, studies that observed a positive impact or not on HCV-positive patients were excluded from this systematic review and meta-analysis. Although our study and other similar articles on the effect of HCV infection on patient and graft survival did not have any publication bias, it appears that we included only papers with a negative impact (Figure 2, 3). Consistent with previous surveys, we observed that the aRR of all-cause mortality and graft failure was significantly higher for seropositive




Figure 4. publication bias for patient survival




Figure 5. publication bias for graft survival




Figure 6. sensitivity analysis for patient's survival

HCV recipients after kidney transplantation.

Role of immunosuppression in HCV-positive kidney transplant recipients

The progression of liver failure in HCV-positive RTRs following immunosuppression is debated. While previous studies have illustrated a detrimental effect on liver function in these patients (10, 11), more recent studies have

observed relatively slow development of liver fibrosis in such patients (1). Luan (2008) performed a study using national data and Cox regression analysis to estimate hazard ratios, adjusted for donor, recipient, and transplant variables. A total of 3708 HCV-positive and 75,629 HCV-negative kidney transplant recipients were included, wherein no calcineurin inhibitors (cyclosporine A or tacrolimus) or steroids had a significant impact on patient mortality. Moreover, the use of mycophenolyte-mofetile (MMF) not only was associated with a significantly reduction in mortality rate, it also had a protective effect (1), despite its association with increased HCV viremia (1). According to another study, HCV replication increases after kidney transplantation, likely due to immunosuppression (1). In contrast, in cultured hepatocytes, cyclosporine A, but not tacrolimus, prevents HCV replication. Notably, more than 50% of HCV-positive kidney transplant recipients who are treated with cyclosporine A have stable liver function and decreased liver fibrosis (1). Nevertheless, in HCV-positive kidney transplant patients, the use of antibody induction has no correlation with viral load (1) and does not have a negative influence on patient survival in these patients (6). It appears that the anti-HCV activity of cyclosporine A differs from its immunosuppressive effects (10). Thus, based on the protective effects of new immunosuppressive drugs, such as MMF and cyclosporine, we hope for greater survival of HCV-positive renal transplant recipients. Yet, controversy still exists regarding the impact of HCV infection on the outcomes of renal transplantation.

Limitations

The majority of articles are not complete; some did not consider Cox regression, and the aRR for patient and graft survival was not reported. Some contributing factors, such as alcohol or drug consumption, were not noted. After renal transplantation, HCV-positive patients have lower patient and graft survival rates compared with HCV-negative patients. However, HCV infection is not a contraindication for renal transplantation; and HCV therapy before transplantation is important to improve the outcome of the patients after transplantation.

Financial support

None declared.

Conflict of interest

None declared.

References

1. Luan FL, Schaubel DE, Zhang H, Jia X, Pelletier SJ, Port FK, et al. Impact of immunosuppressive regimen on survival of kidney transplant recipients with hepatitis C. *Transplantation*. 2008;**85**(11):1601-6.
2. Romero E, Galindo P, Bravo JA, Osorio JM, Perez A, Baca Y, et al. Hepatitis C virus infection after renal transplantation. *Trans-*

- plant Proc. 2008;**40**(9):2933-5.
3. Lezaic V, Stosovic M, Marinkovic J, Rangelov V, Djukanovic L. Hepatitis B and hepatitis C virus infection and outcome of hemodialysis and kidney transplant patients. *Ren Fail.* 2008;**30**(1):81-7.
 4. Einollahi B, Alavian SM. Hepatitis C virus infection and kidney transplantation: a review for clinicians. *Iran J Kidney Dis.* 2010;**4**(1):1-8.
 5. Alavian SM. Patient survival after renal transplantation in HCV and HBV infected patients needs more attention than other risk factors. *Clin Nephrol.* 2009;**72**(4):326-7.
 6. Dominguez-Gil B, Morales JM. Transplantation in the patient with hepatitis C. *Transpl Int.* 2009;**22**(12):1117-31.
 7. Toz H, Nart D, Turan I, Ersoz G, Sezis M, Asci G, et al. The acquisition time of infection: a determinant of the severity of hepatitis C virus-related liver disease in renal transplant patients. *Clin Transplant.* 2009;**23**(5):723-31.
 8. Fabrizi F, Lunghi G, Dixit V, Martin P. Meta-analysis: anti-viral therapy of hepatitis C virus-related liver disease in renal transplant patients. *Aliment Pharmacol Ther.* 2006;**24**(10):1413-22.
 9. Savas N, Ocal S, Colak T, Karakayali H, Yilmaz U, Haberal M. Clinical course of hepatitis C virus infection in renal transplant recipients. *Transplant Proc.* 2007;**39**(4):984-6.
 10. Fabrizi F, Bromberg J, Elli A, Dixit V, Martin P. Review article: hepatitis C virus and calcineurin inhibition after renal transplantation. *Aliment Pharmacol Ther.* 2005;**22**(8):657-66.
 11. Einollahi B, Hajarizadeh B, Bakhtiari S, Lesanpezheshki M, Khataami MR, Nourbala MH, et al. Pretransplant hepatitis C virus infection and its effect on the post-transplant course of living renal allograft recipients. *J Gastroenterol Hepatol.* 2003;**18**(7):836-40.
 12. Aroldi A, Lampertico P, Montagnino G, Passerini P, Villa M, Campise MR, et al. Natural history of hepatitis B and C in renal allograft recipients. *Transplantation.* 2005;**79**(9):1132-6.
 13. Legendre C, Garrigue V, Le Bihan C, Mamzer-Bruneel MF, Chaix ML, Landais P, et al. Harmful long-term impact of hepatitis C virus infection in kidney transplant recipients. *Transplantation.* 1998;**65**(5):667-70.
 14. Gentil MA, Rocha JL, Rodriguez-Algarra G, Pereira P, Lopez R, Bernal G, et al. Impaired kidney transplant survival in patients with antibodies to hepatitis C virus. *Nephrol Dial Transplant.* 1999;**14**(10):2455-60.
 15. Lee WC, Shu KH, Cheng CH, Wu MJ, Chen CH, Lian JC. Long-term impact of hepatitis B, C virus infection on renal transplantation. *Am J Nephrol.* 2001;**21**(4):300-6.
 16. Breitenfeldt MK, Rasenack J, Berthold H, Olschewski M, Schroff J, Strey C, et al. Impact of hepatitis B and C on graft loss and mortality of patients after kidney transplantation. *Clin Transplant.* 2002;**16**(2):130-6.
 17. Bruchfeld A, Wilczek H, Elinder CG. Hepatitis C infection, time in renal-replacement therapy, and outcome after kidney transplantation. *Transplantation.* 2004;**78**(5):745-50.
 18. Morales JM. Hepatitis C virus infection and renal disease after renal transplantation. *Transplant Proc.* 2004;**36**(3):760-2.
 19. Ingsathit A, Thakkinstian A, Kantachavesiri S, Sumethkul V. Different impacts of hepatitis B virus and hepatitis C virus on the outcome of kidney transplantation. *Transplant Proc.* 2007;**39**(5):1424-8.
 20. Batty DS, Jr., Swanson SJ, Kirk AD, Ko CW, Agodoa LY, Abbott KC. Hepatitis C virus seropositivity at the time of renal transplantation in the United States: associated factors and patient survival. *Am J Transplant.* 2001;**1**(2):179-84.
 21. Mahmoud IM, Elhabashi AF, Elsayy E, El-Husseini AA, Sheha GE, Sobh MA. The impact of hepatitis C virus viremia on renal graft and patient survival: a 9-year prospective study. *Am J Kidney Dis.* 2004;**43**(1):131-9.
 22. Lin HH, Huang CC, Huang JY, Yang CW, Wu MS, Fang JT, et al. Impact of HCV infection on first cadaveric renal transplantation, a single center experience. *Clin Transplant.* 2004;**18**(3):261-6.
 23. Ridruejo E, Cusumano A, Diaz C, Davalos Michel M, Jost L, Jost h L, et al. Hepatitis C virus infection and outcome of renal transplantation. *Transplant Proc.* 2007;**39**(10):3127-30.
 24. Gentil Govantes MA, Rodriguez-Benot A, Sola E, Osuna A, Mazuecos A, Bedoya R, et al. Trends in kidney transplantation outcome: the Andalusian Kidney Transplant Registry, 1984-2007. *Transplant Proc.* 2009;**41**(5):1583-5.
 25. Mitwalli AH, Alam A, Al-Wakeel J, Al Suwaida K, Tarif N, Schaar TA, et al. Effect of chronic viral hepatitis on graft survival in Saudi renal transplant patients. *Nephron Clin Pract.* 2006;**102**(2):c72-80.
 26. Periera BJ, Wright TL, Schmid CH, Levey AS. The impact of pre-transplantation hepatitis C infection on the outcome of renal transplantation. *Transplantation.* 1995;**60**(8):799-805.
 27. Pereira BJ, Natov SN, Bouthot BA, Murthy BV, Ruthazer R, Schmid CH, et al. Effects of hepatitis C infection and renal transplantation on survival in end-stage renal disease. The New England Organ Bank Hepatitis C Study Group. *Kidney Int.* 1998;**53**(5):1374-81.
 28. Gentil MA, Perez-Valdivia MA, Munoz-Terol JM, Borrego J, Mazuecos A, Osuna A, et al. Are we still making progress in patient survival after kidney transplantation? Results of a regional registry. *Transplant Proc.* 2009;**41**(6):2085-8.
 29. Hernandez D, Sanchez-Fructuoso A, Gonzalez-Posada JM, Arias M, Campistol JM, Rufino M, et al. A novel risk score for mortality in renal transplant recipients beyond the first posttransplant year. *Transplantation.* 2009;**88**(6):803-9.