

Occult hepatitis B infection in different high risk patients

Amitis Ramezani ^{1, 2}, Mohammad Banifazl ³, Ali Eslamifar ¹, Masoomeh Sofian ⁴, Arezoo Aghakhani ^{1*}

¹Clinical Research Department, Pasteur Institute of Iran, Tehran, IR Iran

² Pediatric Infectious Disease Research Center, Tehran University of Medical sciences, Tehran, IR Iran

³ Iranian Society for Support of Patients With Infectious Diseases, Tehran, IR Iran

⁴ Tuberculosis and Pediatric Infectious Research Center (TPIRC), Arak University of Medical Sciences, Arak, IR Iran

ARTICLE INFO

Article type: Letter to Editor

Article history: Received: 15 Jun 2012 Revised: 25 Jun 2012 Accepted: 05 Jul 2012

Keywords: Hepatitis B Renal Dialysis HIV

Dear Editor,

We read with great interest the article "epidemiology of occult hepatitis B infection among thalassemia, hemophilia and hemodialysis patients" by Arababadi *et al.* (1). because of our common field of interest in investigation of occult hepatitis B infection (OBI) in high risk patients such as HIV infected cases and hemodialysis patients. OBI is defined as the presence of HBV-DNA in the liver tissue or serum without detectable hepatitis B surface antigen (HBsAg) (2). About 20 % of OBI cases are negative for all HBV markers except HBV-DNA, 50 % are positive for hepatitis B core antibody (anti-HBc) and 35 % are positive for hepatitis B surface antibody (anti-HBs) (3). We investigated OBI in HIV positive patients (10, 11). OBI is an important issue in these patients because HIV/HBV co-infected individuals are at increased risk of chronic hepatitis, cirrho-

DOI: 10.5812/hepatmon.7094

Copyright ©2012 Kowsar Corp. All rights reserved.

Please cite this paper as:

Ramezani A, Banifazl M, Eslamifar A, Sofian M, Aghakhani A. Occult hepatitis B infection in different high risk patients. *Hepat Mon.* 2012;**12**(7): 467-8. DOI: 10.5812/hepatmon.7094

Copyright © 2012 Kowsar Corp. All rights reserved.

sis, and hepatocellular carcinoma (4). In the published reports, the prevalence of occult HBV infection in HIV infected patients ranged between 0-10 % using standard PCR methods (5-7) and 35-89 % using more sensitive assays (8, 9). In our study, out of 106 enrolled HIV infected patients, 20.75 % had isolated anti-HBc (HBsAg negative, anti-HBs negative and anti-HBc positive). HBV-DNA was detected in 13.6 % of patients with isolated anti-HBc. We divided these patients based on their HCV status. Out Of 63 anti-HCV positive cases, 28.6 % had isolated anti-HBc and 16.7% of the latter group had OBI. But we did not find any OBI in HIV patients without HCV infection (10, 11). We also conducted the same study in hemodialysis patients (12). The prevalence of occult HBV infection in dialysis patients was reported with a range between 0 to 58 % in published studies (13-16). We found that 6.2 % of 289 enrolled patients had isolated anti-HBc and OBI was detected in 50 % of these patients. Only one of our patients with occult HBV was co-infected with HCV, so a conclusion cannot be reached regarding the association of occult HBV infection and HCV in this study (12).

Overall, the rate of OBI detection can be affected by sen-

^{*} Corresponding author: Arezoo Aghakhani, Clinical Research Department, Pasteur Institute of Iran, No 69, Pasteur Ave., Tehran, 13164, IR Iran. Tel: +98-2166968852, Fax: +98-2166465147, E-mail: araghakhani@hotmail.com

sitivity of the HBV-DNA assay, the sample size, power of the study and composition of the study populations. In conclusion we agree with Arababadi *et al.* (1). Suggestions indicating that OBI is relatively common in high risk cases such as hemodialysis and HIV infected patients. Therefore, the risk of HBV transmission is probable in these groups and screening of high risk groups is recommended.

Authors' Contribution

All authors contributed in the writing of the paper.

Financial Disclosure

None Declared.

References

- Kazemi Arababadi M, Nasiri Ahmadabadi B, Yousefi Daredor H, Kennedy D. Epidemiology of Occult Hepatitis B Infection Among Thalassemic, Hemophilia, and Hemodialysis Patients. *Hepat* Mon. 2012;12(5):315-9.
- Aghakhani A, Banifazl M, Velayati AA, Eslamifar A, Ramezani A. Occult Hepatitis B Virus Infection in Hemodialysis Patients: A Concept for Consideration. *Ther Apher Dial.* 2012.
- Hollinger FB, Sood G. Occult hepatitis B virus infection: a covert operation. J Viral Hepat. 2010;17(1):1-15.
- Alavian SM, Kabir A, Ahmadi AB, Lankarani KB, Shahbabaie MA, Ahmadzad-Asl M. Hepatitis C infection in hemodialysis patients in Iran: a systematic review. *Hemodial Int.* 2010;14(3):253-62.
- 5. Neau D, Winnock M, Jouvencel AC, Faure M, Castera L, Legrand E, et al. Occult hepatitis B virus infection in HIV-infected patients with isolated antibodies to hepatitis B core antigen: Aquitaine cohort, 2002-2003. *Clin Infect Dis.* 2005;**40**(5):750-3.
- 6. Nunez M, Rios P, Perez-Olmeda M, Soriano V. Lack of 'occult'

hepatitis B virus infection in HIV-infected patients. *AIDS*. 2002;**16**(15):2099-101.

- Shire NJ, Rouster SD, Rajicic N, Sherman KE. Occult hepatitis B in HIV-infected patients. J Acquir Immune Defic Syndr. 2004;36(3):869-75.
- Hofer M, Joller-Jemelka HI, Grob PJ, Luthy R, Opravil M. Frequent chronic hepatitis B virus infection in HIV-infected patients positive for antibody to hepatitis B core antigen only. Swiss HIV Cohort Study. Eur J Clin Microbiol Infect Dis. 1998;17(1):6-13.
- 9. Piroth L, Binquet C, Vergne M, Minello A, Livry C, Bour JB, et al. The evolution of hepatitis B virus serological patterns and the clinical relevance of isolated antibodies to hepatitis B core antigen in HIV infected patients. *J Hepatol.* 2002;**36**(5):681-6.
- Azadmanesh K, Mohraz M, Aghakhani A, Edalat R, Jam S, Eslamifar A, et al. Occult hepatitis B virus infection in HIV-infected patients with isolated hepatitis B core antibody. *Intervirology*. 2008;51(4):270-4.
- Ramezani A, Mohraz M, Aghakhani A, Banifazl M, Eslamifar A, Khadem-Sadegh A, et al. Frequency of isolated hepatitis B core antibody in HIV-hepatitis C virus co-infected individuals. *Int J STD AIDS*. 2009;**20**(5):336-8.
- 12. Aghakhani A, Banifazl M, Kalantar E, Eslamifar A, Ahmadi F, Razeghi E, et al. Occult hepatitis B virus infection in hemodialysis patients with isolated hepatitis B core antibody: a multicenter study. *Ther Apher Dial*. 2010;**14**(3):349-53.
- 13. Cabrerizo M, Bartolome J, Caramelo C, Barril G, Carreno V. Molecular analysis of hepatitis B virus DNA in serum and peripheral blood mononuclear cells from hepatitis B surface antigen-negative cases. *Hepatology*. 2000;**32**(1):116-23.
- Fabrizi F, Messa PG, Lunghi G, Aucella F, Bisegna S, Mangano S, et al. Occult hepatitis B virus infection in dialysis patients: a multicentre survey. Aliment Pharmacol Ther. 2005;21(11):1341-7.
- Minuk GY, Sun DF, Greenberg R, Zhang M, Hawkins K, Uhanova J, et al. Occult hepatitis B virus infection in a North American adult hemodialysis patient population. *Hepatology*. 2004;40(5):1072-7.
- Oesterreicher C, Hammer J, Koch U, Pfeffel F, Sunder-Plassmann G, Petermann D, et al. HBV and HCV genome in peripheral blood mononuclear cells in patients undergoing chronic hemodialysis. *Kidney Int*. 1995;48(6):1967-71.