Published online 2021 August 29.

Research Article

Social Disparities in Post-diagnosis Health Behaviors of Iranian Breast Cancer Survivors: The Mediating Role of Psychosocial Factors as Potential Buffering Agents

Mohammad Tavakol ¹, Reza Tayari Ashtiani^{1,*}, Majid Koosheshi², Mohammad Esmaeil Akbari ³ and Maryam Khayamzadeh ³

¹Department of Sociology, School of Social Sciences, University of Tehran, Tehran, Iran
²Department of Demography, School of Social Sciences, University of Tehran, Tehran, Iran
³Cancer Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Corresponding author: Department of Sociology, School of Social Sciences, University of Tehran, Tehran, Iran. Email: reza.t.ashtiani@gmail.com

Received 2020 May 10; Revised 2021 June 22; Accepted 2021 June 22.

Abstract

Background: Breast cancer (BC) is surging as a public health issue in Iran and engagement in positive health behaviors improves the odds of survival and reduces the risk of concomitant comorbidities in BC survivors.

Objectives: The primary purpose of this study was to investigate the impact of socioeconomic inequalities on leisure-time physical activity (LTPA) and fruit and vegetable (F & V) consumption among Iranian BC survivors. Another objective of this study was to examine the role of psychosocial factors, such as stress, self-efficacy, and social support as mediators between socioeconomic status (SES) and these health behaviors. Few studies have investigated social disparities in the health behaviors of cancer survivors. Likewise, the mediating role of psychosocial factors in the SES-health behavior gradient has rarely been explored in the cancer context. Psychosocial factors might have positive implications for socioeconomically disadvantaged survivors.

Methods: Cross-sectional data were obtained from 196 patients with BC by a telephone-administrated questionnaire. The Behavioral Risk Factor Surveillance System (BRFSS) 2013 was partially employed to measure exercise and F & V consumption. Cross-sectional analyses (correlation and logistic regression) were conducted to assess the relationship between SES, psychosocial factors, and health behaviors (P < 0.05).

Results: Regarding LTPA and F & V consumption, more than half of the survivors complied with the recommended levels, consecutively. Both education and family income exhibited a positive association with LTPA and F & V consumption. Furthermore, self-efficacy and social support showed a positive relationship with LTPA and F & V consumption.

Conclusions: The SES impacted health behaviors directly and via intermediary psychosocial factors. The results can inform future studies and interventions; psychosocial factors could buffer the effects of social inequalities on health behaviors. A key policy priority should, therefore, be planning and implementing psychosocial empowerment interventions to promote exercise and a healthy diet among impoverished cancer survivors.

Keywords: Breast Neoplasms, Health Behavior, Social Class, Social Support, Self Efficacy, Exercise, Diet

1. Background

The incidence and mortality of breast cancer (BC) have shown an annual increase of 3.1% and 1.8% from 1980 to 2010, respectively(1). There is a broad consensus on the role of socioeconomic disparities in mortality and morbidity of cancer (2,3), which puts a more severe strain on developing countries such as Iran.

Cancer is the 3rd cause of death in Iran (4-6). BC is the most common cancer and the 5th cause of cancer deaths among Iranian women (7). It is also responsible for 14.2% of overall cancer mortalities (8, 9). Moreover, to amelio-

rate complications arising from the increased length of survival (2, 10, 11), the significance of post-diagnosis health behaviors must be underscored. A review of the literature shows that health habits play a crucial role in increasing the odds of survival (12, 13).

Health behaviors have been instrumental in our understanding of the social gradient in health (14-17); one reason is that health behaviors potentially mediate the relationship between socioeconomic status (SES) and health (18, 19). Furthermore, health behaviors themselves are influenced by SES disparities (18, 20-22). This indicates a need to

Copyright © 2021, Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (http://creativecommons.org/licenses/by-nc/4.0/) which permits copy and redistribute the material just in noncommercial usages, provided the original work is properly cited.

understand the mechanisms by which SES impacts health behaviors (23-27). Identifying these factors might be far more advantageous to survivors than the general population. Nonetheless, there have been few empirical studies paying attention to the mediating role of psychosocial factors in the SES-health behavior gradient in the cancer context (10, 28).

2. Objectives

In this study, we attempted to explore the impact of SES on health behaviors in BC survivors as well as identifying the role of stress, self-efficacy, and social support as mediating factors. We hypothesized that the SES impact on both exercise and F & V consumption could partly be explained by psychosocial mediators. We also hypothesized that the average exercise and F & V consumption of survivors would fairly be higher than the general population. Another hypothesis was that even though the diagnosis of a life-threatening disease might be a teachable moment and lead to tremendous lifestyle changes in survivors, the impact of SES on survivors' lifestyles remains significant. In other words, the effects of SES on health behaviors cannot be completely overcome.

The present study fills a gap in the existing literature; to our knowledge, our study is the first to examine the intermediary role of psychosocial factors in the SES-health behavior gradient in BC survivors. The results of the current study can inform future studies and psychosocial empowerment interventions; enhancing psychosocial factors could potentially buffer the effects of social inequalities on health habits in socioeconomically disadvantaged survivors.

3. Methods

3.1. Study Design and Participants

The data were collected by telephone interviews from 196 BC survivors (from the initial 235) admitted to the Cancer Research Center of Shahid Beheshti University of Medical Sciences from March 2014 to February 2015. Patients' information were retrieved from medical records. The participants were selected randomly from the patients' list, and Krejcie and Morgan table was used for determining the sample size. The questionnaire comprised 97 questions and was administrated in Farsi. The inclusion criteria were as follows: (1) age > 18; (2) minimum 6 months interval between the primary treatment and the interview; (3) absence of serious disability and mobility impairments; (4) no previous mental illness history; and (5) absence of secondary or reoccurring cancer. All patients had gone under surgery and had been treated by either or both chemotherapy and radiotherapy. The oral consent was obtained by two trained interviewers, reading an approved written script to the participants, and briefly explaining the purpose of the study, how they have been selected, and the estimated time of the interview. After ensuring that the participants adequately understood the objectives of the study, they began the interview. Obtaining written consent was not feasible since participants were reachable only by phone. This study was derived from a master's degree thesis submitted to the Faculty of Social Sciences of the University of Tehran and was approved by the Sociology Department, which was approved by the Ethical Committee of University of Tehran Research.

3.2. Measures

3.2.1. Socioeconomic Status

The data were measured by family income and educational attainment level. Family income was defined as the combined income of the family from all resources classified in 4 categories: less than 10 million, between 10 and 20, between 20 and 40, and more than 40 million Iranian rials. Educational attainment was initially categorized into 7 categories and, for statistical analysis, was reclassified into three categories.

3.2.2. Leisure-Time Physical activity

Leisure-time physical activity (LTPA) was measured by the exercise section of the Behavioral Risk Factor Surveillance system (BRFSS) 2013 (29). We used recommended guidelines from the Centers for Disease Control and Prevention (CDC) and The American College of Sports Medicine (ACSM) for adequate weekly physical activity (30). The recommended level was coded as 1, while below the recommended level was coded as 0. Reliability was calculated, using Cronbach's alpha ($\alpha = 0.61$).

3.2.3. F & V consumption

F & V consumption was measured by the BRFSS. The measurement of the F & V consumption was initially based on the American Cancer Society Guidelines, recommending 5 or more F & V servings per day (29). The recommended level was coded as 1, while below the recommended level was coded as 0 ($\alpha = 0.53$).

3.2.4. Self-Efficacy

Self-efficacy was assessed by the brief version of the Self-Efficacy for Exercise scale developed by Sallis et al. (31) ($\alpha = 0.93$).

3.2.5. Stress

Stress was assessed by the validated Farsi version (32) of the Depression Anxiety Stress Scales-21 (DASS-21) introduced by Lovibond and Lovibond (33) ($\alpha = 0.82$).

3.2.6. Social Support

Social support was measured by the validated Farsi version of the MOS Social Support Survey (MOS-SSS) by Sherbourne and Stewart (34). The scale has been widely implemented in the cancer survivorship context (35, 36), and the validity of the Farsi version is well-documented (37) (α = 0.93).

3.3. Data analysis

Descriptive statistics were used to assess the sociodemographic and clinical characteristics of the sample. Eta correlation coefficients were used to assess the relationship between education, family income, stress, social support, and self-efficacy. The associations between education, income, LTPA, and F & V intake were evaluated by one-way ANOVA. After checking for the normality of the distribution and measurement levels, logistic regression was used to examine the relationship between education, income, psychosocial factors, and health behaviors. Data management and analysis were performed, using SPSS 20.0.

4. Results

The demographic and health-related characteristics of the study sample were presented in Table 1. The average age of participants was 48.78 (SD = 10.86). Regarding the stage of diagnosis, 35%, 40%, and 23% of the survivors were diagnosed with stage 1, 2, and 3 BC, respectively, and only 2% were diagnosed with stage 4 BC. Regarding the exercise and F & V consumption, 52% and 53% of the survivors complied with the recommended levels. Psychosocial variables' mean and SD for stress, self-efficacy, and social support were 16.69 (6.36), 24.15 (11.8), and 69.73 (15.68), respectively. Education was negatively associated with stress (η = -0.281) and positivity associated with self-efficacy (η = 0. 251) and social support (η = 0.178). Similarly, there existed a negative association between family income and stress $(\eta = -0.255)$ and a positive relationship between family income and social support (η = 0.223), but no significant relationship between family income and self-efficacy was observed. Also, self-efficacy was positively related to exercise, whereas a negative relationship existed between stress and exercise; social support exhibited a positive relationship with F & V consumption.

As for the logistic regression analysis for education and LTPA (Table 2), respondents in the middle and highesteducation tertile were physically active. It is apparent in

Table 1. Demographics and Health Care Characteristics		
Characteristic	Total, No. (%)	
Marital status		
Married	155 (79.1)	
Bachelor	18 (9.2)	
Widowed	16 (8.2)	
Divorced	7 (3.6)	
Place of living		
Tehran	81 (41.3)	
Other cities-rural areas	115 (58.7)	
Family breadwinner		
Yes	35 (17.9)	
No	160 (81.6)	
Education		
Under high school	64 (32.7)	
High school diploma	67 (34.2)	
College or higher	65 (33.2)	
Monthly family income, million (IRR)		
< 10	39 (20.5)	
10 - 20	79 (41.6)	
20 - 40	49 (25.8)	
> 40	23 (12.1)	
Body mass index		
Normal	50 (25.9)	
Overweight	94 (48.7)	
Obese	49 (25.4)	
Health insurance coverage		
Covered	194 (99)	
Not covered	2 (1)	
General health status		
Excellent	14 (7.1)	
Very good	32 (16.3)	
Good	89 (45.4)	
Fair	56 (28.6)	
Poor	5 (2.6)	

the second model that the relationship between middleeducation tertile and physical activity became insignificant. In the third model, the association between the highest-education group and physical activity remained significant. The same regression analysis was employed to examine the relationship between family income and exercise (Table 3).

The logistics regression for education and F & V intake

Table 2. Logistic Regression Analysis for Education and LTPA ^a			
	Model 1, OR (CI 95 %)	Model 2, OR (CI 95%)	Model 3, OR (CI 95 %)
Education			
Lowest	1		
Middle	2.47 (1.17 - 5.20)*	2.19 (0.99 - 4.84)	1.82 (0.81 - 4.11)
Highest	5.21 (2.35 - 11.54)***	4.01 (1.73 - 9.29)***	3.40 (1.44 - 7.99)***
PA self-efficacy		1.09 (1.046 - 1.14)***	1.09 (1.04 - 1.13)***
Stress			0.93 (0.99 - 0.86)*

^aModels are adjusted for age and general health status.

	Model 1, OR (CI 95 %)	Model 2, OR (CI 95 %)
Education		
Lowest	1	
Middle	2.42 (1.17 - 4.98)*	2.33 (1.11 - 4.87)*
Highest	2.72 (1.28 -5.77)**	2.28 (1.05 - 4.94)*
Social support		1.03 (1.01 - 1.05)**

^aModels are adjusted for age

comprised two models (Table 4). In the first model, education was initially entered into the model and both middle and highest tertiles were positively associated with consumption of F & V. Inserting social support to the second model, both education tertiles retained their significant relationship with the F & V intake. Social support was also associated with F & V intake. A similar regression analysis was employed to assess the relationship between family income and F & V intake (Table 5).

5. Discussion

The present study aimed at investigating the nature of the relationship between SES and LTPA and F & V consumption among BC survivors. To do so, we argued that BC survivors are more active than the general population and adhere more to the recommended F & V consumption than individuals without cancer. We also argued that SES effects on health behaviors are partially explained by mediating psychosocial factors, namely stress, self-efficacy, and social support.

5.1. Health-Related Behaviors

In this study, roughly half of the survivors reported engaging in some type of physical activity. The level of participation in LTPA varies from 20% to 72.5% in different studies (20, 38, 39). Also, more than half of the survivors complied with the recommended F & V consumption level. Regarding both health habits, the reported levels in survivors were remarkably higher than the levels reported by Iran's general population. This confirms our hypothesis on significant differences in the level of activity and healthy diet between survivors and the general population (40, 41).

5.2. Socioeconomic and Psychosocial Factors Related to Health Behaviors

Both education and family income were predictors of exercise in survivors. However, education was a stronger predictor. Our findings were consistent with previous research (39, 42, 43). As for F & V consumption, those with higher education and family income were more likely to consume F & V. Similar to exercise, education was a stronger predictor. Other studies have also reached a similar conclusion in the general population (44-46). These findings validate our hypothesis that even after a lifechanging event such as cancer, the social class remains a strong predictor of health behaviors in survivors; in our study, only participants in high and very high family income categories were physically active and consumed the recommended level of F & V, whereas middle-income survivors appeared to exhibit the same patterns as lowincome families. This denotes that Iranian BC survivors in the middle-income cluster are, to some extent, as vulnerable to economic hardship as low-income groups, and lack the necessary resources to keep up with a healthy lifestyle.

Furthermore, socioeconomic inequalities were observed in all examined psychosocial factors. Education affected both stress and self-efficacy levels, but family income did not show such an impact on self-efficacy. This might be explained by the fact that the majority of participants were homemakers, not the primary breadwinners of the family; this means that the potential benefits of employment and earning income on self-efficacy did not apply to them.

Our findings support our premise that the relationship between both education and family income with LTPA was

f able 4. Logistic Regression Analysis ^a for Family Income and LTPA ^b			
	Model 1, OR (CI 95 %)	Model 2, OR (CI 95%)	Model 3, OR (CI 95 %)
Family income			
Low	1		
Middle	1.92 (0.83 - 4.45)	1.64 (0.68 - 3.95)	1.49 (0.61 - 3.66)
High	2.98 (1.2 - 7.42)*	1.95 (0.76 - 5.40)	1.75 (0.66 - 4.68)
Very high	6.17 (1.88 - 20.23)***	5.10 (1.49 - 17.48)**	4.12 (1.18 - 14.42)*
PA self-efficacy		1.09 (1.05 - 1.14)***	1.09 (1.04 - 1.13)***
Stress			0.92 (0.99 - 086)*
^a Models are adjusted for age and general health status.			

^bLTPA is leisure-time physical activity

Table - Logistic Degression Apal	ucic for Family Incom	and E.S. VConcumption ^a
able 5. Logistic Regression Anal	ysis ior raining meon	e and r & v consumption

	Model 1, OR (CI 95 %)	Model 2, OR (CI 95 %)
Family income		
Low	1	
Middle	1.66 (0.74 - 3.72)	1.54 (0.68 - 3.53)
High	3.25 (1.32 - 7.99)*	2.73 (1.09 - 6.87)*
Very high	5.13 (1.62 - 16.22)**	4.26 (1.32 - 13.76)*
Social support		1.03 (1.01 - 1.05)**

^aModels are adjusted for age.

partially mediated by stress and self-efficacy. Similar findings were reported by other studies (23, 24). We also determined that social support, to some extent, was responsible for education and income disparities in F & V consumption.

Psychosocial factors could be used to mitigate health behavior disparities caused by educational deficits and economic disadvantages. Self-efficacy and social support have been shown to have a direct effect on health behaviors. The authors believe that one feasible solution lies within psychosocial interventions to influence positive health behaviors in cancer survivors. Regarding selfefficacy, less-educated patients should be the primary targets for intervention. Since self-efficacy is one of the key factors in raising exercise levels, interventions to directly promote self-efficacy could be of benefit; boosting selfefficacy appears to be one of the few practical measures capable of enhancing exercise levels in lower-education clusters. However, to date, the majority of research on this subject has solely concentrated on and is limited to providing a descriptive account of existing inequalities without any practical and feasible guidelines for intervention.

Acknowledgments

We are grateful to Somayeh Zoughipor for assistance in data collection and to Samira Sharifi for help with data analysis. This work also benefited from insightful comments from Abou Ali Vedadhir.

Footnotes

Authors' Contribution: Study concept and design: AR. Analysis and interpretation of data: AR, KM, and TM. Drafting of the manuscript: AR. Critical revision of the manuscript for important intellectual content: AR, TM, AM, and KM. Statistical analysis: KM, and AR. Administrative, technical, and material support: KM. Study supervision: TM.

Conflict of Interests: There was no conflict of interests.

Ethical Approval: This study was derived from a master's degree thesis submitted to the Faculty of Social Sciences of the University of Tehran and was approved by the Sociology Department, which was approved by the Ethical Committee of University of Tehran Research.

Funding/Support: The authors received no financial support for the research, authorship, and/or publication of this article.

Informed Consent: Verbal consent was obtained since the interviews were conducted by telephone.

References

- Ferlay J, Forman D, Mathers CD, Bray F. Breast and cervical cancer in 187 countries between 1980 and 2010. *Lancet*. 2012;**379**(9824):1390–1. doi: 10.1016/s0140-6736(12)60595-9.
- Taheri M, Abbasi M, Akbari ME, Almasi-Hashiani A, Tavakol M. Decomposition of Socioeconomic Inequalities in Metastasis, Recurrence, Stage, Grade, and Self-Rated Health of Gastrointestinal Cancer Patients. Int J Cancer Manag. 2019;12(3). doi: 10.5812/ijcm.85802.

- Woods LM, Rachet B, Coleman MP. Origins of socio-economic inequalities in cancer survival: a review. Ann Oncol. 2006;17(1):5-19. doi: 10.1093/annonc/mdj007. [PubMed: 16143594].
- Mousavi SM, Gouya MM, Ramazani R, Davanlou M, Hajsadeghi N, Seddighi Z. Cancer incidence and mortality in Iran. *Ann Oncol.* 2009;**20**(3):556–63. doi: 10.1093/annonc/mdn642. [PubMed: 19073863].
- Mohamad T, Mohsen NR. Health And Social Capital Inequalities Among People Affected/Not Affected By Cancer, Referring To Cancer Institute Of Tehran. *Payavard Salamat.* 2012;6(1).
- Taheri M, Tavakol M, Esmaeil Akbari M, Almasi-Hashiani A, Abbasi M. Associations of demographic, socioeconomic, self-rated health, and metastasis in colorectal cancer in Iran. *Med J Islam Repub Iran*. 2019;**33**:17. doi: 10.34171/mjiri.33.17. [PubMed: 31380307]. [PubMed Central: PMC6662537].
- Rahimzadeh M, Pourhoseingholi MA, Kavehie B. Survival Rates for Breast Cancer in Iranian Patients: a Meta- Analysis. *Asian Pac J Cancer Prev.* 2016;17(4):2223–7. doi: 10.7314/apjcp.2016.17.4.2223. [PubMed: 27221922].
- Akbari A, Razzaghi Z, Homaee F, Khayamzadeh M, Movahedi M, Akbari ME. Parity and breastfeeding are preventive measures against breast cancer in Iranian women. *Breast Cancer*. 2011;18(1):51–5. doi: 10.1007/s12282-010-0203-z. [PubMed: 20217489].
- Tavakol M, Naserirad M. Causal Aspects of Social Capital of Iranian Patients with Cancer: Evidence of Predictive, Modifying and Descriptive Effects in Health Inequality. *Iran J Public Health*. 2013;42(10):1117–25. [PubMed: 26060619]. [PubMed Central: PMC4436539].
- Tabrizi FM. Health promoting behavior and influencing factors in Iranian breast cancer survivors. *Asian Pac J Cancer Prev.* 2015;16(5):1729– 36. doi: 10.7314/apjcp.2015.16.5.1729. [PubMed: 25773817].
- Bishop MM, Lee SJ, Beaumont JL, Andrykowski MA, Rizzo JD, Sobocinski KA, et al. The preventive health behaviors of long-term survivors of cancer and hematopoietic stem cell transplantation compared with matched controls. *Biol Blood Marrow Transplant*. 2010;**16**(2):207– 14. doi: 10.1016/j.bbmt.2009.09.015. [PubMed: 19781657]. [PubMed Central: PMC2819641].
- Bellizzi KM, Rowland JH, Jeffery DD, McNeel T. Health behaviors of cancer survivors: examining opportunities for cancer control intervention. *J Clin Oncol.* 2005;23(34):8884–93. doi: 10.1200/JCO.2005.02.2343. [PubMed: 16314649].
- McTiernan A, Friedenreich CM, Katzmarzyk PT, Powell KE, Macko R, Buchner D, et al. Physical Activity in Cancer Prevention and Survival: A Systematic Review. *Med Sci Sports Exerc*. 2019;**51**(6):1252-61. doi:10.1249/MSS.000000000001937. [PubMed: 31095082]. [PubMed Central: PMC6527123].
- Oberguggenberger A, Meraner V, Sztankay M, Hilbert A, Hubalek M, Holzner B, et al. Health Behavior and Quality of Life Outcome in Breast Cancer Survivors: Prevalence Rates and Predictors. *Clin Breast Cancer*. 2018;**18**(1):38–44. doi: 10.1016/j.clbc.2017.09.008. [PubMed: 29017754].
- Amireault S, Fong AJ, Sabiston CM. Promoting Healthy Eating and Physical Activity Behaviors: A Systematic Review of Multiple Health Behavior Change Interventions Among Cancer Survivors. *Am J Lifestyle Med.* 2018;**12**(3):184–99. doi: 10.1177/1559827616661490. [PubMed: 30202391]. [PubMed Central: PMC6124968].
- Clifford BK, Mizrahi D, Sandler CX, Barry BK, Simar D, Wakefield CE, et al. Barriers and facilitators of exercise experienced by cancer survivors: a mixed methods systematic review. *Support Care Cancer*. 2018;26(3):685–700. doi: 10.1007/s00520-017-3964-5. [PubMed: 29185105].
- Stalsberg R, Eikemo TA, Lundgren S, Reidunsdatter RJ. Physical activity in long-term breast cancer survivors - A mixed-methods approach. *Breast.* 2019;46:126–35. doi: 10.1016/j.breast.2019.05.014. [PubMed: 31158651].
- Pampel FC, Krueger PM, Denney JT. Socioeconomic Disparities in Health Behaviors. Annu Rev Sociol. 2010;36:349–70. doi: 10.1146/annurev.soc.012809.102529. [PubMed: 21909182]. [PubMed Central:

PMC3169799].

- Contoyannis P, Jones AM. Socio-economic status, health and lifestyle. *J Health Econ.* 2004;23(5):965–95. doi: 10.1016/j.jhealeco.2004.02.001. [PubMed: 15353189].
- Park B, Kong SY, Kim J, Kim Y, Park IH, Jung SY, et al. Health Behaviors of Cancer Survivors in Nationwide Cross-Sectional Survey in Korea: Higher Alcohol Drinking, Lower Smoking, and Physical Inactivity Pattern in Survivors with Higher Household Income. *Medicine (Baltimore)*. 2015;94(31). e1214. doi: 10.1097/MD.000000000001214. [PubMed: 26252280]. [PubMed Central: PMC4616611].
- Harper S, Lynch J. Trends in socioeconomic inequalities in adult health behaviors among U.S. states, 1990-2004. *Public Health Rep.* 2007;**122**(2):177-89. doi: 10.1177/003335490712200207. [PubMed: 17357360]. [PubMed Central: PMC1820442].
- Karimi SE, Rafiey H, Sajjadi H, Nosrati Nejad F. Identifying the Social Determinants of Breast Health Behavior: a Qualitative Content Analysis. *Asian Pac J Cancer Prev.* 2018;**19**(7):1867–77. doi: 10.22034/APJCP.2018.19.7.1867. [PubMed: 30049199]. [PubMed Central: PMC6165651].
- Lehto E, Konttinen H, Jousilahti P, Haukkala A. The role of psychosocial factors in socioeconomic differences in physical activity: a population-based study. *Scand J Public Health*. 2013;**41**(6):553–9. doi: 10.1177/1403494813481642. [PubMed: 23508949].
- Cerin E, Leslie E. How socio-economic status contributes to participation in leisure-time physical activity. Soc Sci Med. 2008;66(12):2596–609. doi: 10.1016/j.socscimed.2008.02.012. [PubMed: 18359137].
- Droomers M, Schrijvers CT, van de Mheen H, Mackenbach JP. Educational differences in leisure-time physical inactivity: a descriptive and explanatory study. *Soc Sci Med.* 1998;47(11):1665–76. doi: 10.1016/s0277-9536(98)00272-x. [PubMed: 9877337].
- Krueger PM, Chang VW. Being poor and coping with stress: health behaviors and the risk of death. *Am J Public Health*. 2008;**98**(5):889–96. doi: 10.2105/AJPH.2007.114454. [PubMed: 18382003]. [PubMed Central: PMC2374822].
- Conti G, Hansman C. Personality and the education-health gradient: a note on "understanding differences in health behaviors by education". *J Health Econ.* 2013;**32**(2):480–5. doi: 10.1016/j.jhealeco.2012.07.005. [PubMed: 23245875]. [PubMed Central: PMC3767122].
- Frank-Stromborg M, Pender NJ, Walker SN, Sechrist KR. Determinants of health-promoting lifestyle in ambulatory cancer patients. Soc Sci Med. 1990;31(10):1159–68. doi: 10.1016/0277-9536(90)90237-m. [PubMed: 2274804].
- CDC. BRFSS Data Quality, Validity, and Reliability. 2013. Available from: http://www.cdc.gov/brfss/publications/data_qvr.htm.
- Pescatello LS. ACSM's guidelines for exercise testing and prescription. Lippincott Williams & Wilkins; 2013.
- Sallis JF, Pinski RB, Grossman RM, Patterson TL, Nader PR. The development of self-efficacy scales for healthrelated diet and exercise behaviors. *Health Educ Res.* 1988;3(3):283–92. doi: 10.1093/her/3.3.283.
- 32. Moradi MF, Ahadi H, Jomehri F, Rahgozar M. Relationship between psychological distress and quality of life in women with breast cancer. J Zabol Univ Med Sci Health Service. 2012;4(2):51-9.
- Lovibond PF, Lovibond SH. The structure of negative emotional states: comparison of the Depression Anxiety Stress Scales (DASS) with the Beck Depression and Anxiety Inventories. *Behav Res Ther.* 1995;**33**(3):335–43. doi: 10.1016/0005-7967(94)00075-u. [PubMed: 7726811].
- Sherbourne CD, Stewart AL. The MOS social support survey. Soc Sci Med. 1991;32(6):705–14. doi: 10.1016/0277-9536(91)90150-b. [PubMed: 2035047].
- 35. Lee MK, Park S, Lee ES, Ro J, Kang HS, Shin KH, et al. Social support and depressive mood 1 year after diagnosis of breast cancer compared with the general female population: a prospective cohort study. Support Care Cancer. 2011;19(9):1379–92. doi: 10.1007/s00520-010-0960-4.

[PubMed: 20676695].

- Nausheen B, Gidron Y, Peveler R, Moss-Morris R. Social support and cancer progression: a systematic review. J Psychosom Res. 2009;67(5):403-15. doi: 10.1016/j.jpsychores.2008.12.012. [PubMed: 19837203].
- Tamannaeifar MR, Mansourinik A. The relationship between personality characteristics, social support and life satisfaction with university students' academic performance. *Q J Res Plan High Educ*. 2014;**20**(1):149–66.
- Pudkasam S, Polman R, Pitcher M, Fisher M, Chinlumprasert N, Stojanovska L, et al. Physical activity and breast cancer survivors: Importance of adherence, motivational interviewing and psychological health. *Maturitas*. 2018;**116**:66–72. doi: 10.1016/j.maturitas.2018.07.010. [PubMed: 30244781].
- Neil-Sztramko SE, Winters-Stone KM, Bland KA, Campbell KL. Updated systematic review of exercise studies in breast cancer survivors: attention to the principles of exercise training. Br J Sports Med. 2019;53(8):504–12. doi: 10.1136/bjsports-2017-098389. [PubMed: 29162619].
- Esteghamati A, Noshad S, Nazeri A, Khalilzadeh O, Khalili M, Nakhjavani M. Patterns of fruit and vegetable consumption among Iranian adults: a SuRFNCD-2007 study. *Br J Nutr.* 2012;**108**(1):177-81. doi: 10.1017/S0007114511005393. [PubMed: 22017813].
- 41. Fakhrzadeh H, Djalalinia S, Mirarefin M, Arefirad T, Asayesh H, Safiri S, et al. Prevalence of physical inactivity in Iran: a systematic re-

view. J Cardiovasc Thorac Res. 2016;8(3):92-7. doi: 10.15171/jcvtr.2016.20. [PubMed: 27777692]. [PubMed Central: PMC5075364].

- Gidlow C, Johnston LH, Crone D, Ellis N, James D. A systematic review of the relationship between socio-economic position and physical activity. *Health Educ J*. 2016;65(4):338–67. doi: 10.1177/0017896906069378.
- Naik H, Qiu X, Brown MC, Eng L, Pringle D, Mahler M, et al. Socioeconomic status and lifestyle behaviours in cancer survivors: smoking and physical activity. *Curr Oncol.* 2016;23(6):e546-55. doi: 10.3747/co.23.3166. [PubMed: 28050143]. [PubMed Central: PMC5176380].
- Estaquio C, Druesne-Pecollo N, Latino-Martel P, Dauchet L, Hercberg S, Bertrais S. Socioeconomic differences in fruit and vegetable consumption among middle-aged French adults: adherence to the 5 A Day recommendation. J Am Diet Assoc. 2008;108(12):2021–30. doi: 10.1016/j.jada.2008.09.011. [PubMed: 19027405].
- 45. Giskes K, Turrell G, Patterson C, Newman B. Socioeconomic differences among Australian adults in consumption of fruit and vegetables and intakes of vitamins A, C and folate. J Hum Nutr Diet. 2002;15(5):375-85. discussion 387-90. doi: 10.1046/j.1365-277x.2002.00387.x. [PubMed: 12270018].
- Roos E, Talala K, Laaksonen M, Helakorpi S, Rahkonen O, Uutela A, et al. Trends of socioeconomic differences in daily vegetable consumption, 1979-2002. *Eur J Clin Nutr.* 2008;62(7):823–33. doi: 10.1038/sj.ejcn.1602798. [PubMed: 17522606].