Published online 2022 January 8.

Research Article

Effect of 5'-fluoro-2'-deoxycytidine, 5-azacytidine, and 5-aza-2'-deoxycytidine on DNA Methyltransferase 1, CIP/KIP Family, and INK4a/ARF in Colon Cancer HCT-116 Cell Line

Masumeh Sanaei¹ and Fraidoon Kavoosi^{1,*}

¹Research Center for Non-communicable Diseases, Jahrom University of Medical Sciences, Jahrom, Iran

corresponding author: Research Center for Non-communicable Diseases, Jahrom University of Medical Sciences, Jahrom, Iran. Email: kavoosifraidoon@gmail.com

Received 2020 October 19; Revised 2021 December 05; Accepted 2021 December 06.

Abstract

Background: Cyclin-dependent kinase inhibitors (CKIs) are the negative regulator of cell cycle progression, which inhibits cyclincdk complexes, resulting in cell cycle arrest. Recently, we evaluated the effect of 5-Aza-CdR on DNMT1 gene expression in the WCH-17 hepatocellular carcinoma (HCC) cell line.

Objectives: The current study was designed to analyze the effects of 5-aza-2'-deoxycytidine (5-Aza-CdR, decitabine), 5-azacytidine (5-AzaC, vidaza), and 5'-fluoro-2'-deoxycytidine (FdCyd) on INK4a/ARF, CIP/KIP, and DNA methyltransferase 1 gene expression, apoptosis induction, and cell growth inhibition in colon cancer HCT-116 cell line.

Methods: The colon cancer HCT-116 cell line was treated with 5-azaC, 5-Aza-CdR, and FdCyd at 24 and 48h. To determine colon cancer HCT-116 cell viability, cell apoptosis, and the relative expression level of the INK4a/ARF, CIP/KIP, and DNA methyltransferase 1 genes, MTT assay, flow cytometry, and qRT-PCR were done, respectively.

Results: 5-azaC, 5-Aza-CdR, and FdCyd significantly inhibited colon cancer HCT-116 cell growth and induced apoptosis. Besides, they significantly increased CIP/KIP (p21CIP1, p27KIP1, and p57KIP2) and INK4 (p14^{ARF}, p15^{INK4b}, and p16^{INK4a}) and decreased DNMT1 gene expression. Besides, minimal and maximal apoptosis were seen in the groups treated with FdCyd and 5-Aza-CdR, respectively. The IC50 for CAF for FdCyd was 1.72 \pm 0.23 and 1.63 \pm 0.21 μ M at 24 and 48h, respectively. The IC50 for CAF for 5-AzaC was 2.18 \pm 0.33 and 1.98 \pm 0.29 μ M at 24 and 48h, respectively. The IC50 for CAF for 5-AzaC was 4.08 \pm 0.61 and 3.18 \pm 0.50 μ M at 24 and 48h, respectively.

Conclusions: The 5-azac, 5-Aza-CdR, and FdCyd can reactivate the INK4a/ARF and CIP/KIP families through inhibition of DNMT1 activity.

Keywords: Cyclin-dependent Kinases, DNA (Cytosine-5-)-Methyltransferases, Colonic Neoplasms

1. Background

The mammalian cell cycle is a highly ordered process and is divided into discrete phases including G1 the phase (gap 1), DNA synthesis phase (S phase), G2 phase, and mitosis phase (M phase). In this cycle, genetic information is transmitted from one cell to the next. The cycle is derived by the cyclin-dependent kinases (Cdks) and their regulatory subunit, cyclin, together, which propel the cell cycle through the various phases of the cell cycle (1). The transition between each phase is controlled by the kinase activity composed of cyclins and their partner, Cdk. Cyclins have fluctuated levels during the cell cycle. In contrast, the level of Cdk proteins typically remains unchanged throughout the cycle (2). Cyclin-dependent kinase inhibitors (CKIs) are the negative regulator of cell cycle progression, which inhibit cyclin-cdk complexes, resulting in cell cycle arrest (3). Two families of CKIs have been identified: CIP/KIP and INK4 families. The first family members comprise p21CIP1 (p21), p27KIP1 (p27), and p57KIP2 (p57) (4). The INK4a/ARF includes 3 important tumor suppressor genes (TSGs): p14^{ARF} (p14), p15^{INK4b} (p15), and p16^{INK4a} (p16) (5). The posttranslational modifications of cell cycle include phosphorylation, histone acetylation, DNA methylation, ADPribosylation, and ubiquitination (6).

Aberrant DNA methylation is known as an important epigenetic change occurring in cancer. The silencing of the INK4a/ARF (7) and CIP/KIP (8) families by methylation

Copyright © 2021, Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (http://creativecommons.org/licenses/by-nc/4.0/) which permits copy and redistribute the material just in noncommercial usages, provided the original work is properly cited.

plays an important role in several cancers. The hypermethylation of INK4a/ARF has been reported in breast cancer (9), Mantle cell lymphoma (MCL)(10), colorectal cancer (11), colon cancer (12), and gastric cancer (13). Similarly, the hypermethylation of the CIP/KIP family has been shown in the lung, gastric, and colon cancer (14). In mammals, cytosine methylation is achieved by 3 DNA methyltransferases (DNMT's) comprising DNMT1, DNMT3A, and DNMT3B (15). Several studies have indicated the role of DNMT1 in the regulation of the expression of TSGs in colon cancer cells (16).

DNA methyltransferase inhibitors (DNMTIs) can inhibit DNMTs activity resulting in the re-activation of silenced TSGs. These compounds are divided into two groups including the nucleoside inhibitors and the nonnucleoside inhibitors. First group includes 5'-fluoro-2'deoxycytidine (FdCyd), 5-azacytidine (5-AzaC, vidaza), and 5-aza-2'-deoxycytidine (5-Aza-CdR, decitabine). The second group includes epigallocatechin3-gallate (EGCG), procaine, and RG108 (17). The inhibitory effect of these compounds has been demonstrated in lymphoid cancer, ovarian cancer, cervical cancer, lung cancer (18), breast cancer (19), and human colon cancer cells (20, 21). Recently, we evaluated the effect of 5-Aza-CdR on DNMT1 gene expression in the WCH-17 hepatocellular carcinoma (HCC) cell line (22).

2. Objectives

The current study was designed to analyze the effects of 5-azac, 5-Aza-CdR, and FdCyd on the INK4a/ARF family (p15INK4a, p14, and p15), CIP/KIP family (p21, p27, and p57), and DNA methyltransferase 1 gene expression, cell growth inhibition, and cell apoptosis induction in colon cancer HCT-116 cell line.

3. Methods

Human colon cancer HCT-116 cell line was provided from the National Cell Bank of Iran-Pasteur Institute and maintained in DMEM supplemented with fetal bovine serum 10% and antibiotics (0.1 mg/mL streptomycin and 100 U/mL penicillin). 5-azac, 5-Aza-CdR, and FdCyd were purchased from Sigma (St. Louis, MO, USA) and dissolved in DMSOto make a master stock solution (23). Further concentration was obtained by diluting the provided solution. Other compounds including materials and kits were purchased as provided for our previous works (22, 24, 25), including FBS (fetal bovine serum), MTT, Real-time

3.1. Cell Culture and Cell Viability

The HCT-116 cells were cultured in DMEM supplemented with 10% FBS and antibiotics (0.1 mg/ml streptomycin and 100 U/mL penicillin) at 37°C in 5% CO2 for 24 h. Subsequently, the cells were seeded into 96-well plates $(4 \times 10^5$ cells per well). After 1 day, culture medium was removed and the experimental medium containing various doses of 5-azac (0, 0.5, 1, 2.5, 5, and 10 μ M), 5-Aza-CdR $(0, 0.5, 1, 2.5, 5, and 10 \mu M)$, and FdCyd (0, 0.5, 1, 2.5, 5, and10 μ M). The concentrations were selected based on and in the range of our previous works and other researchers' reports. The control groups were treated with the solvent (DMSO) only at a concentration of 0.05 %. After 24 and 48 h of treatment, the treated and untreated HCT-116 cells were evaluated by MTT assay to obtain cell viability, the MTT solution (5 mg/mL) was added to each well and allowed incubation for 4 h at 37°C. To dissolve all of the crystals, the solution was replaced by DMSO for 10 min. Subsequently, the absorbance spectrum was determined by a microplate reader at a wavelength of 570 nM.

3.2. Cell Apoptosis Assay

To determine cell apoptosis, the HCT-116 cells were cultured at a density of 4×10^5 cells/well and incubated overnight and, then, the cells were treated with 5-azac, 5-Aza-CdR, and FdCyd, based on IC50 values indicated in Table 1 for different periods (24 and 48 h). Subsequently, the treated and untreated HCT-116 cells were harvested and prepared for flow cytometry by trypsinization, washing twice with cold PBS, and then stained with annexin-V-(FITC) and propidium iodide (PI). The apoptotic cells were determined by FACScan flow cytometry.

3.3. Real-time Quantitative Reverse Transcription Polymerase Chain Reaction (qRT-PCR)

To determine the relative expression level of the INK4a/ARF family (p15INK4a, p14, and p15), CIP/KIP family (p21, p27, and p57), and DNA methyltransferase 1 gene, qRT-PCR was done. The HCT-116 cells were treated with 5-azaC, 5-Aza-CdR, and FdCyd, based on IC50 values indicated in Table 1 for different periods (24 and 48 h). Total RNA from the HCT-116 cells was extracted, using the RNeasy kit according

Fable 1. IC50 Values and Apoptosis. The Cells Were Treated with Compounds, to Determine IC50 Values and Apoptosis							
Cell Line	Duration/Hour	IC50 Value/µM	LogIC50	R Squared	Apoptosis (%)	P-Value ^a	
FdCyd	24	1.72	0.2357	0.7815	7.64	< 0.0003	
FdCyd	48	1.63	0.2121	0.9729	11.15	< 0.0001	
5-AzaC	24	2.18	0.3391	0.8771	9.45	< 0.0001	
5-AzaC	48	1.98	0.2981	0.8066	50.1	< 0.0001	
5-Aza-CdR	24	4.08	0.6116	0.7221	13.16	< 0.0001	
5-Aza-CdR	48	3.18	0.5034	0.7001	83.66	< 0.0001	

^aThe P-value indicates the difference between the control group (untreated cell line) with each experimental group treated with the compound.

to the protocol and treated by RNase-free DNase to eliminate the genomic DNA before cDNA synthesis (26). Realtime PCR reactions were performed, using the Steponeplus. Thermal cycling conditions were initial denaturation at 95°C for 10 minutes, followed by 40 cycles of denaturation at 95°C for 20 seconds, annealing at 58°C for 15 seconds, and extension at 72°C for 15 seconds. Thermal cycling conditions for DNMT1 was initial denaturation at 95°C for 10 minutes, followed by 40 cycles of denaturation at 95°C for 15 seconds, annealing at 60°C for 20 seconds, and extension at 72°C for 20 seconds.

The data were analyzed, using the comparative Ct $(\Delta\Delta ct)$ method. A melting curve was used to determine the melting temperature of specific amplification products and primer dimmers. GAPDH was used as a reference gene for internal control. The primer sequences of the genes used in the current article are shown in Table 2 (27-32). The relative RT-PCR determines the expression level in comparison with a reference sample. It is based on the expression levels of a target gene versus a housekeeping gene (Pfaffl 2004) (33).

3.4. Statistical Analysis

Data from three independent experiments were analyzed with one-way analysis of variance (ANOVA) using Graphpad Prism Software version 8.0. A significant difference is expressed as P < 0.05.

4. Results

4.1. Cell Viability

The viability of colon cancer HCT-116 cells treated with 5-azac, 5-Aza-CdR, and FdCyd was measured by MTT assay. As indicated in Figure 1, 5-azac, 5-Aza-CdR, and FdCyd induced significant cell growth inhibition. The half-maximal inhibitory concentration (IC50) values are demonstrated in Table 1. As indicated in Table 1, The IC50 for CAF for Fd-Cyd was 1.72 \pm 0.23 and 1.63 \pm 0.21 μ M at 24 and 48h, respectively. The IC50 for CAF for 5-AzaC was 2.18 \pm 0.33 and 1.98 \pm 0.29 μ M at 24 and 48h, respectively. The IC50 for CAF for 5-Aza-CdR was 4.08 \pm 0.61 and 3.18 \pm 0.50 μ M at 24 and 48h, respectively.

4.2. Cell Apoptosis

To determine cell apoptosis, the HCT-116 cells were treated with FdCyd (1.72 \pm 0.23 and 1.63 \pm 0.21 μ M), 5-AzaC (2.18 \pm 0.33 and 1.98 \pm 0.29 μ M), and 5-Aza-CdR (4.08 \pm 0.61 and 3.18 \pm 0.50 μ M) for 24 and 48 h and stained using annexin-V-(FITC) and PI as mentioned in the method section. As depicted in Figure 2, all of the compounds significantly induced apoptosis (Figure 3). Besides, minimal and maximal apoptosis were seen in the groups treated with FdCyd and 5-Aza-CdR, respectively. The percentage of HCT-116 apoptotic cells is shown in Table 1.

4.3. Gene Expression

The effect of 5-azac, 5-Aza-CdR, and FdCyd on INK4a/ARF family (p15INK4a, p14, and p15), CIP/KIP family (p21, p27, and p57), and DNA methyltransferase 1 gene expression was assessed by qRT-PCR. To determine the relative gene expression, the HCT-116 cells were treated with FdCyd (1.72 \pm 0.23 and 1.63 \pm 0.21 μ M), 5-AzaC (2.18 \pm 0.33 and 1.98 \pm 0.29 μ M), and 5-Aza-CdR (4.08 \pm 0.61 and 3.18 \pm 0.50 μ M) for 24 and 48h, respectively. The result indicated that treatment with 5-azac, 5-Aza-CdR, and FdCyd (24 and 48h) upregulated INK4a/ARF (p15INK4a, p14, and p15) and CIP/KIP (p21, p27, and p57) families, and down-regulated DNMT1 gene expression significantly (Figure 4).

5. Discussion

Cellular gene transcription is directly under the influence of the genomic structure and organization. DNA





100

0

0

Viability % 50



0.5

1

C1

Concentration (µM)



FdCyd, 48h

C2

Concentration (µM)





10

Primer	Primer Sequences (5' to 3')	Product Length/bp	Reference
P14	Forward: TACTGAGGAGCCAGCGTCTA	146	(27)
	Reverse: TGCACGGGTCGGGTGAGAGT	140	
P15	Forward: AAGCTGAGCCCAGGTCTCCTA	02	(28)
	Reverse: CCACCGTTGGCCGTAAACT	93	
P15	Forward: CTTCCTGGACACGCTGGT	160	(29)
	Reverse: GCATGGTTACTGCCTCTGGT	102	
P21	Forward: CGATGGAACTTCGACTTTGTCA	220	(30)
	Reverse: GCACAAGGGTACAAGACAGTG	220	
P 27	Forward: GGTTAGCGGAGCAATGCG	127	(30)
	Reverse: TCCACAGAACCGGCATTTG	127	
P 57	Forward: GCGGCGATCAAGAAGCTGT	52	(31)
	Reverse: GCTTGGCGAAGAAATCGGAGA		
DNMT	Forward: GCACAAACTGACCTGCTTCA	313	(32)
DINMIT	Reverse: GCCTTTTCACCTCCATCAAA	213	
GAPDH	Forward: TGTGGGCATCAATGGATTTGG	116	(31)
	Reverse: ACACCATGTATTCCGGGTCAAT	116	

Table 2. The Primer Sequences of INK4a/ARF, CIP/KIP, and DNMT1 Genes^a

^a The relative expression level of the INK4a/ARF family (p15INK4a, p14, and p15), CIP/KIP family (p21, p27, and p57), and DNA methyltransferase 1 gene were determined through qRT-PCR.

methylation is an important epigenetic modification that has widespread influences on gene transcription and expression (34). It is profoundly altered in human cancers. The silencing of TSGs by this epigenetic change is known as a key mechanism in tumorigenesis (35). The potential anticancer activity of DNMT inhibitors has been extensively evaluated in recent years. These compounds are widely studied because DNA demethylation induces the reactivation of TSGs that are silenced by promoter hypermethylation (36).

In the present study, we demonstrated that DNA demethylating agents 5-azac, 5-Aza-CdR, and FdCyd inhibited HCT-116 cell growth and induced apoptosis in colon cancer HCT-116 cell lines. Furthermore, we did a further evaluation to find the molecular mechanism of the compounds. Thus, we found that these agents could re-activate INK4a/ARF family (p15INK4a, p14, and p15) and the CIP/KIP family (p21, p27, and p57) by inhibition of DNA methyltransferase 1 gene expression.

Similar apoptotic pathways have been shown by other researchers. It has been indicated that 5-Aza-CdR increases the expression of both p15^{INK4a} and p19^{INK4d} in the human lung cancer cell line (37). In vitro studies have shown such molecular mechanisms for the member of INK4 in col-

orectal cancer (38), leukemia (39), ovarian cancer (40), gastric cancer (41), and HCC (42). As mentioned above DNA methyltransferase inhibitors play their role through the re-activation of the CIP/KIP family. A similar pathway has been indicated in HCC (43), gastric cancer (44), breast, and lung cancer (45). We reported that DNA demethylating agents 5-azac, 5-Aza-CdR, and FdCyd play their role through inhibition of DNMT1 activity.

In addition to this function, several in vitro studies have been demonstrated that these agents can inhibit DNMT3a and DNMT3b in the colon cancer HCT-116 (46), DNMT3B in human endometrial cancer (47), DNMT1, and/or DNAT 3b mediates in lung cancer, esophageal cancer, and malignant pleural mesothelioma cells (48), and DNMT3b in testicular germ cell tumors (TGCT) (49).

Additionally, other members of DNA methyltransferase inhibitors can induce apoptosis with similar molecular mechanisms. Our previous work indicated that zebularine (a membership of HDACIs) can induce apoptosis through down-regulation of DNMT1, DNMT3a, and DNMT3b and up-regulation of p21Cip1/Waf1/Sdi1, p27Kip1, and p57Kip2 in colon cancer LS 174T cell line (50). Besides, we reported that zebularine induces apoptosis by down-regulation of DNMT1, 3a, and 3b and up-regulation



Figure 2. The apoptotic effect of 5-azac, 5-Aza-CdR, and FdCyd on HCT-116 cell line versus control groups at different periods (24 and 48h). The cells were treated with FdCyd (1.72 and 1.63 μM), 5-AzaC (2.18 and 1.98 μM), and 5-Aza-CdR (4.08 and 3.18 μM) for 24 and 48h, respectively. Quadrant (Q) 2 and 3, late and primary apoptosis, respectively, were calculated in this graph. A: FdCyd treated groups (AI: Control; A2: 24h; A3: 48h); B: 5-azac treated groups (BI: Control; B2: 24h; B3: 48h); C: 5-Aza-CdR treated groups (CI: Control; C2: 24h; C3: 48h).

of *p21Cip1/Waf1/Sdi1*, *p27Kip1*, *p57Kip2* in colon cancer LS 180 cell line (51). Meanwhile, DNMT1 inhibition is not the only apoptotic pathway of DNMT1s. It has been reported that 5 aza 2' deoxycytidine treatment resulted in significant FAS

gene up-regulation in the HT 29 cell line and it plays its role through the extrinsic apoptotic pathway (52). Inconsistent with our report, it has been reported that 5 AZA treatments cannot induce a significant change in cell proliferation,



Figure 3. The comparative apoptotic effects of 5-azac, 5-Aza-CdR, and FdCyd on HCT-116 cell line. The cells were treated with FdCyd (1.72 and 1.63 μM), 5-AzaC (2.18 and 1.98 μM), and 5-Aza-CdR (4.08 and 3.18 μM) for 24 and 48h, respectively. Asterisks indicate significant differences between the HCT-116 treated and untreated control groups. All compounds induced significant apoptosis in HCT-116 cell line (part A). Minimal and maximal apoptosis were seen in the groups treated with FdCyd and 5-Aza-CdR, respectively (parts B, and C). It should be noted that ^{***} and ^{****} indicate P < 0.0003 and P < 0.0001, respectively.

cell cycle arrest, cell apoptosis, and mtDNA copy number in HCT116, SW480, LS 174T, and HT 29 cell lines (53). Finally, DN-MTIs can play their apoptotic roles through various molecular mechanisms.

We indicated that inhibition of DNMT1 activity by 5-azac, 5-Aza-CdR, and FdCyd induces re-activation of INK4a/ARF family (p15INK4a, p14, and p15) and CIP/KIP family (p21, p27, and p57), resulting in apoptosis induction in colon cancer HCT-116 cell line.

We did not assess the other DNMTs gene expression such as DNMT3a and DNMT3b in this work. Thus, this evaluation is recommended.

5.1. Conclusions

In summary, our findings indicated that 5-azac, 5-Aza-CdR, and FdCyd inhibited colon cancer HCT-116 cell line and induced apoptosis in this cell line. The most likely molecular mechanism underlying these compounds inhibited HCT-116 cell growth and induced apoptosis involves downregulation of DNMT1 and up-regulation of CIP/KIP (p21, p27, and p57) and INK4 (p14, p15, and p15INK4a) genes expression. This result suggests that 5-azac, 5-Aza-CdR, and FdCyd may have wide therapeutic applications in colon cancer.

Footnotes

Authors' Contribution: M.S., K.F.: Study concept and design, acquisition of data; analysis and interpretation of data; critical revision of the manuscript for important intellectual content; statistical analysis; M.S.: Drafting of the manuscript; administrative, technical, or material support; K.F.: Study supervision. All authors approved the final version of the manuscript for publication.

Conflict of Interests: The authors report no conflict of interest.

Ethical Approval: This work was approved by the Ethics Committee of Jahrom University of Medical science with a code number of IR.JUMS.REC.1398.099.

Funding/Support: This article was supported by the adjutancy of research of Jahrom University of Medical Sciences, Iran.

A2

A1



Figure 4. The relative expression level of the INK4a/ARF family (p15INK4a, p14, and p15), CIP/KIP family (p21, p27, and p57), and DNA methyltransferase 1 gene in the colon cancer HCT-116 cell line. The cells were treated with FdCyd (1.72 and 1.63 μ M), 5-AzaC (2.18 and 1.98 μ M), and 5-Aza-CdR (4.08 and 3.18 μ M) for 24 and 48h, respectively. A significant difference was seen between treated and untreated control groups. Asterisks indicate significant differences between the treated and untreated control groups. A: FdCyd treated groups; B: 5-azac treated groups; C: 5-Aza-CdR treated groups. It should be noted that *, ***, and **** indicate P < 0.0195, P < 0.0011, and P < 0.0001, respectively.

A

References

- 1. Israels ED, Israels LG. The cell cycle. *Oncologist*. 2000;**5**(6):510–3. doi: 10.1634/theoncologist.5-6-510. [PubMed: 11110604].
- Lim S, Kaldis P. Cdks, cyclins and CKIs: roles beyond cell cycle regulation. *Development*. 2013;**140**(15):3079–93. doi: 10.1242/dev.091744. [PubMed: 23861057].
- Shankland SJ, Wolf G. Cell cycle regulatory proteins in renal disease: role in hypertrophy, proliferation, and apoptosis. *Am J Physiol Renal Physiol*. 2000;**278**(4):F515–29. doi: 10.1152/ajprenal.2000.278.4.F515. [PubMed: 10751212].
- 4. Funk JO. Cell cycle checkpoint genes and cancer. 5. Wiley; 2001. p. 1-6.
- Zhang Z, Rosen DG, Yao JL, Huang J, Liu J. Expression of p14ARF, p15INK4b, p16INK4a, and DCR2 increases during prostate cancer progression. *Mod Pathol*. 2006;19(10):1339–43. doi: 10.1038/modpathol.3800655. [PubMed: 16799475].
- Shanmugam MK, Arfuso F, Arumugam S, Chinnathambi A, Jinsong B, Warrier S, et al. Role of novel histone modifications in cancer. *Oncotarget*. 2018;9(13):11414–26. doi: 10.18632/oncotarget.23356. [PubMed: 29541423]. [PubMed Central: PMC5834259].
- Park MT, Lee SJ. Cell cycle and cancer. J Biochem Mol Biol. 2003;36(1):60– 5. doi: 10.5483/bmbrep.2003.36.1.060. [PubMed: 12542976].
- Chim CS, Fung TK, Wong KF, Lau JS, Law M, Liang R. Methylation of INK4 and CIP/KIP families of cyclin-dependent kinase inhibitor in chronic lymphocytic leukaemia in Chinese patients. *J Clin Pathol.* 2006;**59**(9):921–6. doi: 10.1136/jcp.2005.035089. [PubMed: 16565223]. [PubMed Central: PMC1860467].
- Rivandi M, Khorrami MS, Fiuji H, Shahidsales S, Hasanzadeh M, Jazayeri MH, et al. The 9p21 locus: A potential therapeutic target and prognostic marker in breast cancer. *J Cell Physiol*. 2018;233(7):5170–9. doi: 10.1002/jcp.26332. [PubMed: 29240242].
- Hutter G, Scheubner M, Zimmermann Y, Kalla J, Katzenberger T, Hubler K, et al. Differential effect of epigenetic alterations and genomic deletions of CDK inhibitors [p16(INK4a), p15(INK4b), p14(ARF)] in mantle cell lymphoma. *Genes Chromosomes Cancer*. 2006;45(2):203-10. doi: 10.1002/gcc.20277. [PubMed: 16258956].
- Ishiguro A, Takahata T, Saito M, Yoshiya G, Tamura Y, Sasaki M, et al. Influence of methylated p15 and p16 genes on clinicopathological features in colorectal cancer. *J Gastroenterol Hepatol*. 2006;21(8):1334–9. doi: 10.1111/j.1440-1746.2006.04137.x. [PubMed: 16872319].
- Burri N, Shaw P, Bouzourene H, Sordat I, Sordat B, Gillet M, et al. Methylation silencing and mutations of the p14ARF and p16INK4a genes in colon cancer. *Lab Invest*. 2001;81(2):217–29. doi: 10.1038/labinvest.3780230. [PubMed: 11232644].
- Zhang F, Wang L, Wu PP, Yan ZW, Zheng L, Yu YY, et al. In situ analysis of p16/INK4 promoter hypermethylation in esophageal carcinoma and gastric carcinoma. *Chin J Dig Dis.* 2004;5(4):149–55. doi: 10.1111/j.1443-9573.2004.00172.x. [PubMed: 15612883].
- Fang JY, Lu YY. Effects of histone acetylation and DNA methylation on p21(WAF1) regulation. World J Gastroenterol. 2002;8(3):400– 5. doi: 10.3748/wjg.v8.i3.400. [PubMed: 12046058]. [PubMed Central: PMC4656409].
- Robertson KD. DNA methylation, methyltransferases, and cancer. Oncogene. 2001;20(24):3139–55. doi: 10.1038/sj.onc.1204341. [PubMed: 11420731].
- Fang JY, Lu R, Mikovits JA, Cheng ZH, Zhu HY, Chen YX. Regulation of hMSH2 and hMLH1 expression in the human colon cancer cell line SWI116 by DNA methyltransferase 1. *Cancer Lett.* 2006;**233**(1):124–30. doi: 10.1016/j.canlet.2005.03.005. [PubMed: 16473673].
- 17. Lyko F, Brown R. DNA methyltransferase inhibitors and the development of epigenetic cancer therapies. J Natl Cancer Inst.

2005;97(20):1498-506. doi: 10.1093/jnci/dji311. [PubMed: 16234563].

- Gnyszka A, Jastrzebski Z, Flis S. DNA methyltransferase inhibitors and their emerging role in epigenetic therapy of cancer. *Anticancer Res.* 2013;33(8):2989–96. [PubMed: 23898051].
- Billam M, Sobolewski MD, Davidson NE. Effects of a novel DNA methyltransferase inhibitor zebularine on human breast cancer cells. *Breast Cancer Res Treat*. 2010;**120**(3):581-92. doi: 10.1007/s10549-009-0420-3. [PubMed: 19459041]. [PubMed Central: PMC3901992].
- Robert MF, Morin S, Beaulieu N, Gauthier F, Chute IC, Barsalou A, et al. DNMT1 is required to maintain CpG methylation and aberrant gene silencing in human cancer cells. *Nat Genet.* 2003;**33**(1):61–5. doi: 10.1038/ng1068. [PubMed: 12496760].
- Thakur S, Feng X, Qiao Shi Z, Ganapathy A, Kumar Mishra M, Atadja P, et al. ING1 and 5-azacytidine act synergistically to block breast cancer cell growth. *PLoS One*. 2012;7(8). e43671. doi: 10.1371/journal.pone.0043671. [PubMed: 22916295]. [PubMed Central: PMC3423394].
- Sanaei M, Kavoosi F. Effects of 5-aza-2'-deoxycytidine and Valproic Acid on Epigenetic-modifying DNMT1 Gene Expression, Apoptosis Induction and Cell Viability in Hepatocellular Carcinoma WCH-17 cell line. *Iran J Pediatr Hematol Oncol.* 2019;9(2):83–90. doi: 10.18502/ijpho.v9i2.607.
- Sanaei M, Kavoosi F. Effect of 5-Aza-2'-Deoxycytidine in Comparison to Valproic Acid and Trichostatin A on Histone Deacetylase 1, DNA Methyltransferase 1, and CIP/KIP Family (p21, p27, and p57) Genes Expression, Cell Growth Inhibition, and Apoptosis Induction in Colon Cancer SW480 Cell Line. Adv Biomed Res. 2019;8:52. doi: 10.4103/abr.abr_91_19. [PubMed: 31516890]. [PubMed Central: PMC6712896].
- Sanaei M, Kavoosi F. Effect of DNA Methyltransferase in Comparison to and in Combination with Histone Deacetylase Inhibitors on Hepatocellular Carcinoma HepG2 Cell Line. Asian Pac J Cancer Prev. 2019;20(4):1119-25. doi: 10.31557/APJCP.2019.20.4.1119. [PubMed: 31030484]. [PubMed Central: PMC6948907].
- Kavoosi F, Sanaei M. Comparative Analysis of the Effects of Valproic Acid and Tamoxifen on Proliferation, and Apoptosis of Human Hepatocellular Carcinoma WCH 17 CellLin. *Iran J Pediatr Hematol Oncol.* 2018;8(1):12–20.
- Sanaei M, Kavoosi F. Effect of Curcumin and Trichostatin A on the Expression of DNA Methyltransfrase 1 in Hepatocellular Carcinoma Cell Line Hepa 1-6. Iran J Pediatr Hematol Oncol. 2018;8(4):193–201.
- Maruo S, Zhao B, Johannsen E, Kieff E, Zou J, Takada K. Epstein-Barr virus nuclear antigens 3C and 3A maintain lymphoblastoid cell growth by repressing p16INK4A and p14ARF expression. *Proc Natl Acad Sci U S A*. 2011;108(5):1919–24. doi: 10.1073/pnas.1019599108. [PubMed: 21245331]. [PubMed Central: PMC3033265].
- Li G, Ji Y, Liu C, Li J, Zhou Y. Reduced levels of p15INK4b, p16INK4a, p21cip1 and p27kip1 in pancreatic carcinoma. *Mol Med Rep.* 2012;5(4):1106-10. doi: 10.3892/mmr.2012.771. [PubMed: 22293850]. [PubMed Central: PMC3493078].
- Wang Y, Zang X, Wang Y, Chen P. High expression of p16INK4a and low expression of Bmit are associated with endothelial cellular senescence in the human cornea. *Mol Vis.* 2012;18:803-15. [PubMed: 22509111]. [PubMed Central: PMC3324359].
- Wang L, Wang G, Yang D, Guo X, Xu Y, Feng B, et al. Euphol arrests breast cancer cells at the G1 phase through the modulation of cyclin D1, p21 and p27 expression. *Mol Med Rep.* 2013;8(4):1279–85. doi: 10.3892/mmr.2013.1650. [PubMed: 23969579].
- 31. Xu Y, Zhong C, Ding S, Huang H, Shen Z. MicroRNA-221 promotes human non-small cell lung cancer cell H460 growth. Int J Clin

Exp Med. 2015;**8**(2):2024–30. [PubMed: 25932132]. [PubMed Central: PMC4402779].

- Xu Y, Chao L, Wang J, Sun Y. miRNA-148a regulates the expression of the estrogen receptor through DNMT1-mediated DNA methylation in breast cancer cells. *Oncol Lett.* 2017;14(4):4736–40. doi: 10.3892/ol.2017.6803. [PubMed: 29085474]. [PubMed Central: PMC5649610].
- 33. Pfaffl MW. Quantification strategies in real-time PCR. *AZ of quantitative PCR*. 2004;1:89–113.
- Metivier R, Gallais R, Tiffoche C, Le Peron C, Jurkowska RZ, Carmouche RP, et al. Cyclical DNA methylation of a transcriptionally active promoter. *Nature*. 2008;452(7183):45–50. doi: 10.1038/nature06544. [PubMed: 18322525].
- Kondo Y. Epigenetic cross-talk between DNA methylation and histone modifications in human cancers. *Yonsei Med J.* 2009;**50**(4):455– 63. doi: 10.3349/ymj.2009.50.4.455. [PubMed: 19718392]. [PubMed Central: PMC2730606].
- Zhu WG, Otterson GA. The interaction of histone deacetylase inhibitors and DNA methyltransferase inhibitors in the treatment of human cancer cells. *Curr Med Chem Anticancer Agents*. 2003;3(3):187– 99. doi: 10.2174/1568011033482440. [PubMed: 12769777].
- Zhu WG, Dai Z, Ding H, Srinivasan K, Hall J, Duan W, et al. Increased expression of unmethylated CDKN2D by 5-aza-2'-deoxycytidine in human lung cancer cells. *Oncogene*. 2001;20(53):7787–96. doi: 10.1038/sj.onc.1204970. [PubMed: 11753657].
- 38. Zheng S, Chen P, McMillan A, Lafuente A, Lafuente MJ, Ballesta A, et al. Correlations of partial and extensive methylation at the p14(ARF) locus with reduced mRNA expression in colorectal cancer cell lines and clinicopathological features in primary tumors. *Carcinogenesis*. 2000;**21**(11):2057-64. doi: 10.1093/carcin/21.11.2057. [PubMed: 11062168].
- Farinha NJ, Shaker S, Lemaire M, Momparler L, Bernstein M, Momparler RL. Activation of expression of p15, p73 and E-cadherin in leukemic cells by different concentrations of 5-aza-2'-deoxycytidine (Decitabine). *Anticancer Res.* 2004;24(1):75-8.
- Meng CF, Su B, Li W. DNA demethylation is superior to histone acetylation for reactivating cancer-associated genes in ovarian cancer cells. *Mol Med Rep.* 2011;4(6):1273–8. doi: 10.3892/mmr.2011.557. [PubMed: 21850374].
- Meng CF, Zhu XJ, Peng G, Dai DQ. Promoter histone H3 lysine 9 dimethylation is associated with DNA methylation and aberrant expression of p16 in gastric cancer cells. *Oncol Rep.* 2009;**22**(5):1221–7. doi:10.3892/or_00000558. [PubMed: 19787243].
- Liu LH, Xiao WH, Liu WW. Effect of 5-Aza-2'-deoxycytidine on the P16 tumor suppressor gene in hepatocellular carcinoma cell line HepG2. World J Gastroenterol. 2001;7(1):131–5. doi: 10.3748/wjg.v7.i1.131. [PubMed: 11819749]. [PubMed Central: PMC4688690].
- Matsuda Y. Molecular mechanism underlying the functional loss of cyclindependent kinase inhibitors p16 and p27 in hepatocellular carcinoma. *World J Gastroenterol.* 2008;14(11):1734–40. doi: 10.3748/wjg.14.1734. [PubMed: 18350604]. [PubMed Central: PMC2695913].
- 44. Sun D, Toan X, Zhang Y, Chen Y, Lu R, Wang X, et al. Mammalian target of rapamycin pathway inhibition enhances the effects of 5-aza-dC

on suppressing cell proliferation in human gastric cancer cell lines. *Sci China C Life Sci.* 2008;**51**(7):640–7. doi: 10.1007/s11427-008-0080-2. [PubMed: 18622747].

- Kobatake T, Yano M, Toyooka S, Tsukuda K, Dote H, Kikuchi T, et al. Aberrant methylation of p57KIP2 gene in lung and breast cancers and malignant mesotheliomas. *Oncol Rep.* 2004;12(5):1087–92. [PubMed: 15492797].
- 46. Schneider-Stock R, Diab-Assef M, Rohrbeck A, Foltzer-Jourdainne C, Boltze C, Hartig R, et al. 5-Aza-cytidine is a potent inhibitor of DNA methyltransferase 3a and induces apoptosis in HCT-116 colon cancer cells via Gadd45- and p53-dependent mechanisms. *J Pharmacol Exp Ther.* 2005;**312**(2):525–36. doi: 10.1124/jpet.104.074195. [PubMed: 15547111].
- Cui M, Wen Z, Chen J, Yang Z, Zhang H. 5-Aza-2'-deoxycytidine is a potent inhibitor of DNA methyltransferase 3B and induces apoptosis in human endometrial cancer cell lines with the up-regulation of hMLH1. *Med Oncol*. 2010;**27**(2):278–85. doi: 10.1007/s12032-009-9204-1. [PubMed: 19306077].
- Kassis ES, Zhao M, Hong JA, Chen GA, Nguyen DM, Schrump DS. Depletion of DNA methyltransferase 1 and/or DNA methyltransferase 3b mediates growth arrest and apoptosis in lung and esophageal cancer and malignant pleural mesothelioma cells. J Thorac Cardiovasc Surg. 2006;131(2):298–306. doi: 10.1016/j.jtcvs.2005.05.022. [PubMed: 16434257].
- Beyrouthy MJ, Garner KM, Hever MP, Freemantle SJ, Eastman A, Dmitrovsky E, et al. High DNA methyltransferase 3B expression mediates 5-aza-deoxycytidine hypersensitivity in testicular germ cell tumors. *Cancer Res*. 2009;69(24):9360–6. doi: 10.1158/0008-5472.CAN-09-1490. [PubMed: 19951990]. [PubMed Central: PMC2795063].
- 50. Sanaei M, Kavoosi F. Effect of Zebularine in Comparison to and in Combination with Trichostatin A on CIP/KIP Family (p21Cip1/Waf1/Sdi1, p27Kip1, and p57Kip2), DNMTs (DNMT1, DNMT3a, and DNMT3b), Class I HDACs (HDACs 1, 2, 3) and Class II HDACs (HDACs 4, 5, 6) Gene Expression, Cell Growth Inhibition and Apoptosis Induction in Colon Cancer LS 174T Cell Line. Asian Pac J Cancer Prev. 2020;21(7):2131–9. doi: 10.31557/APJCP.2020.21.7.2131. [PubMed: 32711442]. [PubMed Central: PMC7573409].
- Sanaei M, Kavoosi F. Investigation of the Effect of Zebularine in Comparison to and in Combination with Trichostatin A on p21Cip1/Wafi/Sdi1, p27Kip1, p57Kip2, DNA Methyltransferases and Histone Deacetylases in Colon Cancer LS 180 Cell Line. *Asian Pac J Cancer Prev.* 2020;**21**(6):1819–28. doi: 10.31557/APJCP.2020.21.6.1819. [PubMed: 32592383]. [PubMed Central: PMC7568903].
- Manoochehri M, Borhani N, Karbasi A, Koochaki A, Kazemi B. Promoter hypermethylation and downregulation of the FAS gene may be involved in colorectal carcinogenesis. *Oncol Lett.* 2016;**12**(1):285– 90. doi: 10.3892/ol.2016.4578. [PubMed: 27347139]. [PubMed Central: PMC4906593].
- Tong H, Zhang L, Gao J, Wen S, Zhou H, Feng S. Methylation of mitochondrial DNA displacement loop region regulates mitochondrial copy number in colorectal cancer. *Mol Med Rep.* 2017;**16**(4):5347–53. doi: 10.3892/mmr.2017.7264. [PubMed: 28849075]. [PubMed Central: PMC5647067].