

Family History as a Risk for Upper Gastrointestinal Tract Cancer: A Case Control Study

Safae A¹, Moghimi Dehkordi B¹, Fatemi SR¹, Maserat E¹, Ghafarnejad F¹, Zali MR¹

Abstract

Background: Although, family history of cancer is an important risk factor for upper gastrointestinal cancers development, but limited information is available on the upper gastrointestinal cancers associated with family history in Iran. The purpose of this study was to define upper gastrointestinal cancers risk associated with family history of cancer.

Methods: This study was conducted as a case control study. A total number of 1,010 cases of upper gastrointestinal cancer and 1,010 healthy controls were recruited. For family history of cancer, questions were asked about any malignant tumor in first and second degree relatives. Adjusted odds ratio estimates for the association family history and upper gastrointestinal cancers risk and corresponding 95% confidence intervals were obtained.

Results: A family history of any malignant tumor in relatives was associated with 1.3 fold increased risks of upper gastrointestinal cancers. A first-degree family history of esophageal and gastric cancer was significantly associated with upper gastrointestinal cancers development, with an adjusted OR of 4.7(CI 95%: 2.6-8.4).

Conclusion: Our findings suggested that risk for upper gastrointestinal cancers increases among individuals with family history of cancer. Therefore, appropriate screening strategies especially in relatives of patients should be considered to prevent and control of disease.

Key Words: Upper gastrointestinal tract; Gastrointestinal neoplasm; Esophageal neoplasm, Case-Control Study

Please cite this article as: Safae A, Moghimi-Dehkordi B, Fatemi SR, Maserat E, Ghafarnejad F, Zali MR. Family History as a Risk for Upper Gastrointestinal Tract Cancer: A Case Control Study. *Iran J Cancer Prev.* 2011;4(3):114-8.

1. Research Center of Gastroenterology and Liver Diseases, Shahid Beheshti University of Medical Science, Tehran, Iran

Corresponding Author:

Bijan Moghimi Dehkordi
MSc of Epidemiology
Tel: (+98) 21 22 43 25 15
Email: b_moghimi_de@yahoo.com
Received: 3 Jan. 2011
Accepted: 20 Jun. 2011
Iran J Cancer Prev. 2011; 3:114-8

Introduction

Malignancies of the upper gastrointestinal (UGI) tract form a heterogeneous group of cancers characterized by unique epidemiology and biology[1]. They are much more common in Asia than in western countries.[2-4]. Upper gastrointestinal cancers cause 55% of all cancer-related deaths in Iran, with gastric cancer being the most common[5]. Although the incidence of gastric cancer (GC) has declined in the past several decades, it remains the second most common cause of cancer-related deaths in the world[6, 7]. Gastric cancer is an important cause of mortality due to cancer and is predicted to be the eighth leading cause of all death worldwide in the year 2010[8]. GC is the most common malignancy in Iran[9-11]. In recent years, cancer morbidity and mortality increased in Iran, and GC is ordered second among all cancers in this country[12-14].

Also, esophageal cancer (EC) is among the 10 most common tumors and is the 6th leading cause of malignancy deaths worldwide[2, 15]. The incidence rate appears to have decreased during the last 3 decades but is still 6th common cancer in Iran.

A number of environmental factors are now known to be related to the development of the GC[16]. Smoking, alcohol consumption, Helicobacter pylori infection, dietary habits and obesity may be risk factors of gastric cancer[6]. However, neither the relative risks associated with these factors nor their prevalence in the Iranian population is of appropriate magnitude to justify the extremely high incidence. It is therefore important that familial factors also be considered as possible risk factors contributing to the development of GC[17]. Most epidemiologic studies reported a risk of gastric cancer between 1.5 and 3.5 for subjects with relatives with gastric cancer[17].

The principle risk factors for esophageal cancer are thought to be environmental. Smoking, alcohol consumption, a dietary deficiency of fruits and vegetables, the consumption of very hot beverages and exposure to carcinogens due to opium consumption as have been stressed important risk factors, but recently the genetic nature of esophageal cancer has received much attention[4]. One study from Iran showed that subjects with one or more first degree relatives with esophageal cancer have a odds ratio of developing esophageal cancer about 3.5[4].

It has been suggested that familial esophageal cancer may develop earlier and have a poorer prognosis than sporadic esophageal cancer. Therefore, it is reasonable to hypothesize that FH of cancer might also predict survival of upper gastrointestinal cancer[18].

Family history (FH) by different relative types and risk of UGI cancers has been only rarely reported in Iran[4, 19]. Therefore, this study was under taken to confirm the impact of familial susceptibility on UGI cancer development.

Materials and Methods

This research was conducted in Research Institute for Gastroenterology and Liver Disease (RIGLD), Shahid Beheshti University of Medical Science, Tehran capital of Iran, as a case control study. A total number of 1,010 cases of UGI cancer and 1,010 healthy controls were recruited. We included cases in this study who: had their diagnosis histologically confirmed without previous history of cancer. Controls were randomly selected among the healthy participants in a health survey conducted by the Department of Health System Research of RIGLD in that a total of 5,500 subjects aged ≥ 15 years were invited to participate in an interview about the occurrence of cancer in their first- or second-degree relatives[20, 21].

After obtaining informed consent, both cases and controls were interviewed to obtain information on demographic characteristics, and FH. For FH of cancer, questions were asked about any malignant tumor in first degree relatives, including father, mother, siblings, and offspring. In addition, information on cancer history was also collected on second degree relatives as the cases and controls (i.e., grand father or mother, aunt and uncle).

The FH was extracted from a standard history form completed by the patient or from the record created by a health care provider. The ethical

committee at Shahid Beheshti University of Medical Sciences approved the study.

Adjusted odd ratio(OR) estimates for the association FH and UGI cancers risk and corresponding 95% confidence intervals were obtained. To account for potential confounders, multivariate logistic regression analyses were conducted. All the regression models included terms for age and sex. Statistical analysis was performed with SPSS (version 13.0). All analyses were two-sided, and statistical significance was defined as a P-value less than 0.05.

Results

A total of 2,020 cases and controls were enrolled in this study. The mean (\pm SD) age of cases was 53.8 ± 18.6 years. 68.1% of patients were male. Of the 1,010 controls, 529 (52.4%) were female and 481(47.6%) were male. By design, the proportion of men was higher in cases than in controls, and the age distribution was higher in cases (P-value is $P < 0.0001$).

Two hundred twenty two (22%) of cases and 237 (23.5%) of controls reported the occurrence different types of malignant tumors in their family members ($P > 0.05$). A FH of any malignant tumor in relatives was associated with 1.3 fold increased risks of UGI cancers (Table 1).

Table 2 presents the distribution of UGI cancers in the first-degree relatives of cases and controls. A first-degree family history of esophageal and gastric cancer was significantly associated with UGI cancers development, with an adjusted OR of 4.7 (CI 95%: 2.6-8.4). Overall, 3.5% of cases versus 2.5% of controls reported a family history of UGI cancer in second degree relatives. Results indicating that having a positive family history of UGI cancer in second degree relatives increase one's risk of UGI cancer about 3 fold.

None of the other cancer sites in relatives showed a significant association with risk of UGI cancers (Table 3).

Discussion

In present case-control study focusing on FH of cancer and risk of UGI cancers, we observed increased risk for UGI cancers among individuals with FH of any malignant tumor and UGI cancers. The excess risk was most apparent for first degree relatives compared to second degree relatives. This result is in agreement with another studies[4, 18].

Table1. Associations between family history of any malignant tumor and risk of UGI cancers

	Cases n (%)	Controls n (%)	Crude OR (CI: 95%)	Adjusted OR ⁺ (CI: 95%)
Family history of Cancer				
FDR[*]/SDR^{**}				
No	788(78.0)	773(76.5)		
Yes	222(22.0)	237(23.5)	0.94(0.78-1.12)	1.3(1.01-1.66)
FDR				
No	864(85.5)	896(88.7)		
Yes	146(14.5)	114(11.3)	1.28(1.01-1.64)	1.24(0.92-1.68)
SDR				
No	918(90.9)	872(86.3)		
Yes	92(9.1)	138(13.7)	0.63(0.48-0.84)	1.27(0.90-1.78)

* Family history of cancer in First Degree Relatives

** Family history of cancer in Second Degree Relatives

+ Adjusted according to age and sex

Table2. Associations between family history of UGI malignant tumor and risk of UGI cancers

	Cases n (%)	Controls n (%)	Crude OR (CI: 95%)	Adjusted OR ⁺ (CI: 95%)
Family history of UGI malignancy				
FDR/SDR				
No	910(90.1)	970(96.0)		
Yes	100(9.9)	40(4.0)	2.5(1.73-3.61)	3.67(2.34-5.76)
FDR[*]				
No	935(92.6)	993(98.3)		
Yes	75(7.4)	17(1.7)	4.68(2.75-7.99)	4.67(2.60-8.41)
SDR^{**}				
No	974(96.4)	985(97.5)		
Yes	36(3.6)	25(2.5)	1.44(0.86-2.39)	2.81(1.49-5.32)

* Family history of cancer in First Degree Relatives

** Family history of cancer in Second Degree Relatives

+ Adjusted according to age and sex

Table3. Associations between family history of any malignant tumor except esophageal and gastric cancer and risk of UGI cancers

	Cases n (%)	Controls n (%)	Crude OR (CI: 95%)	Adjusted OR ⁺ (CI: 95%)
Family history of other Cancer				
FDR/SDR				
No	880(87.1)	801(79.3)		
Yes	130(12.9)	209(20.7)	0.62(0.50-0.77)	0.80(0.61-1.06)
FDR[*]				
No	928(91.9)	912(90.3)		
Yes	82(8.1)	98(9.7)	0.84(0.62-1.12)	0.74(0.53-1.66)
SDR^{**}				
No	947(93.8)	889(88.0)		
Yes	63(6.2)	121(12.0)	0.52(0.38-0.71)	1.03(0.70-1.51)

* Family history of cancer in First Degree Relatives

** Family history of cancer in Second Degree Relatives

+ Adjusted according to age and sex

Some studies suggest that FH increases the risk of cancer at many sites and is not site-specific [22]. The association between FH and UGI cancer risk in present survey strengthened as FH became more specific: OR=1.3 for FH of any cancer, OR=3.67 for FH of UGI cancer. Our estimated OR is higher than

one Chinese study[18]. Differences from previous reports could be due to variation in the prevalence of UGI cancers in the study populations. In contrast, no association was observed between FH of non-UGI cancer and risk of UGI cancer. This suggests that the shared susceptibility is UGI specific, and FH of non-

UGI cancer is not important for UGI cancer risk prediction[17, 18].

A number of previous studies reported increased risk of gastric cancer among individuals with a positive family history of cancer [17, 23]. Only a few studies have been examined the risk of gastric cancer by anatomic sub sites (cardia and non-cardia)[24]. FH of gastric cancer was associated with increased risk of non-cardia cancer only, and FH of any cancer was associated with increased risk of cardia cancer only[24]. Ying Gao et al. [18] found that FH of gastric cardia cancer was associated with increased risk for both gastric cardia and gastric non-cardia adenocarcinoma, but no association was observed between FH of gastric non-cardia cancer and risk of either gastric non-cardia adenocarcinoma or gastric cardia adenocarcinoma.

It is suggest that younger age at onset is often related with genetic role in disease. Wen et al.[25] With a case-control study showed that, FH correlated with early age of onset for esophageal cancer. Garavello et al. [16,26] found also that younger individuals with a FH of cancer have higher risks for development of cancer. In this study we don't have access to the age of onset of disease.

Although, inherited susceptibility is a very important risk factor in UGI cancers, but co-occurrence of UGI cancers among family members does not necessarily reflect shared genetic susceptibility; it could also be due to shared environmental exposures[17, 18, 27]. Therefore, studying different types of relatives (blood and non-blood relatives sharing household) might provide information to help differentiate genetic and environmental components in esophageal and gastric pathogenesis. While several studies have reported that environmental factors like may play a more important role than genetic factors in UGI development[4] some studies haven't found a strong correlation between FH of UGI and environmental factors, such as lifestyles[28]. This contrast may be due to variation in the frequency of esophageal and gastric susceptibility alleles, or due to variation in environmental risk factors, or due to a combination of the two[1, 4]. In the west countries, risk factors include male sex, smoking, and alcohol use, and these are not risk factors in Iran. Further studies will provide information about the relative contribution of genetic and environmental factors in the development of UGI cancers in Iran.

Limitations of this study are: potential recall bias due to the nature of the case-control study design. Information on family history was self-reported, and it is possible that cases of GC and EC may tend to

recall a FH of UGI cancer or other cancers more accurately than controls. Aggregation of cancer among non-blood relatives in the same household, like wives and husbands, supports environmental compared to genetic factors in cancer etiology. But we don't access to information about non-blood relatives and also about number of first and second degree relatives affected. Also, unfortunately, it was not possible to obtain pathological confirmation of the diagnosis of cancer in the relatives, and some diagnosis may have been missed. Another limitation of this study is potential misclassification bias. Different diagnostic criteria and classification methods may have led to incorrect designation of tumor location and histology for a proportion of patient.

So the interpretation and generalization results of this study should be done with caution.

A strong point of this study is the large and representative sample. Another strong point of the study is that we conducted multivariate analyses, which allowed us to control the effects of confounder factors in analysis.

Conclusion

Our findings suggested that risk for UGI cancers increases among individuals with family history of cancer. Future studies will provide information about the relative contribution of genetic and environmental factors in the development of upper gastrointestinal cancers in Iran. We support the establishment of a national program to investigate potential risk factors for UGI cancers by tumor site and possible gene-environment interactions in patients from different racial groups. Such analysis may help us to better understand the epidemiology of UGI cancers and provide a basic foundation for developing prevention strategies among individuals at high risk.

Acknowledgement

This study was funded by Research Center for Gastroenterology and Liver Diseases of Shahid Beheshti University of Medical Sciences, Tehran, Iran. The authors also acknowledge the valuable collaboration of local Health Center staffs for data gathering.

Conflicts of Interest

None to declare.

Authors' Contribution

AS conceived and designed this study and interpreted the results and drafted the manuscript. BMD designed and carried out the analyses. SRF

revised the manuscript. FGH and EM contributed to data gathering and data entering to the software. MRZ supervised the project. All authors read and improved the final manuscript.

References

1. Shah MA, Kurtz RC. Upper gastrointestinal cancer predisposition syndromes. *Hematol Oncol Clin North Am*. 2010;24(5):815-35.
2. Marjani HA, Biramijamal F, Hossein-Nezhad A, Islami F, Pourshmas A, Semnani S. Prevalence of esophageal cancer risk factors among Turkmen and non-Turkmen ethnic groups in a high incidence area in Iran. *Arch Iran Med*. 2010;13(2):111-5.
3. Somi MH, Mousavi SM, Rezaeifar P, Naghashi SH. Cancer incidence among the elderly population in the Northwest of Iran: A population based study. *Iranian Journal of Cancer Prevention*. 2009;2(3):117-26.
4. Akbari MR, Malekzadeh R, Nasrollahzadeh D, Amanian D, Sun P, Islami F, et al. Familial risks of esophageal cancer among the Turkmen population of the Caspian littoral of Iran. *Int J Cancer*. 2006;119(5):1047-51.
5. Taghavi N, Nasrollahzadeh D, Merat S, Yazdanbod A, Hormazdi M, Sotoudeh M, et al. Epidemiology of upper gastrointestinal cancers in Iran: a sub site analysis of 761 cases. *World J Gastroenterol*. 2007;13(40):5367-70.
6. Hemminki K, Sundquist J, Ji J. Familial risk for gastric carcinoma: an updated study from Sweden. *British journal of cancer*. 2007;96(8):1272-7.
7. Moghimi-Dehkordi B, Safaei A, Zali MR. Survival rates and prognosis of gastric cancer using an actuarial life-table method. *Asian Pacific Journal of Cancer Prevention*. 2008;9(2):317-22.
8. Pourhoseingholi MA, Pourhoseingholi A, Vahedi M, Moghimi-Dehkordi B, Safaei A, Ashtari S, et al. Alternative for Cox Regression: Parametric Model to Analysis the Survival of Cancer Patients. *Iranian Journal of Cancer Prevention*. 2010;4(1):1-9.
9. Moghimi-Dehkordi B, Safaei A, Pourhoseingholi MA, Zali MR. Effect of demographic and clinicopathologic factors on prognosis of early gastric cancer in Iran. *Asian Pacific Journal of Cancer Prevention*. 2008;9(4):585-8.
10. Mehrabian AA, Esna-Ashari F, Zham H, Hadizadeh M, Bohlooli M, Khayamzadeh M, et al. Gastric cancer prevalence, according to survival data in Iran (National Study-2007). *Iranian Journal of Public Health*. 2010;39(3):27-31.
11. Movahedi M, Afsharfard A, Moradi A, Nasermoaddeli A, Khoshnevis J, Fattahi F, et al. Survival rate of gastric cancer in Iran. *Journal of Research in Medical Sciences*. 2009;14(6):367-73.
12. Safaei A, Moghimi-Dehkordi B, Fatemi SR, Ghiasi S, Pourhoseingholi MA, Zali MR. Clinicopathological features of gastric cancer: A study based on cancer registry data. *Iranian Journal of Cancer Prevention*. 2009;2(2):67-70.
13. Moghimi-Dehkordi B, Safaei A, Tabei SZ. A comparison between Cox proportional hazard models and logistic regression on prognostic factors in gastric cancer. *East African journal of public health*. 2009; 6 Suppl(1):20-2.
14. Moghimi-Dehkordi B, Safaei A, Fatemi R, Ghiasi S, Zali MR. Impact of age on prognosis in Iranian patients with gastric carcinoma: review of 742 cases. *Asian Pac J Cancer Prev*. 2010;11(2):335-8.
15. Tabatabaee SA, Hashemi SM, Eidy M, Davarpanah Jazi AH. Predicting factors for anastomotic leakage after esophageal cancer resection. *Iranian Journal of Cancer Prevention*. 2009;2(2):103-6.
16. Brenner H, Rothenbacher D, Arndt V. Epidemiology of stomach cancer. *Methods Mol Biol*. 2009;472:467-77.
17. Foschi R, Lucenteforte E, Bosetti C, Bertuccio P, Tavani A, La Vecchia C, et al. Family history of cancer and stomach cancer risk. *Int J Cancer*. 2008;123(6):1429-32.
18. Gao Y, Hu N, Han X, Giffen C, Ding T, Goldstein A, et al. Family history of cancer and risk for esophageal and gastric cancer in Shanxi, China. *BMC cancer*. 2009;9:269.
19. Yaghoobi M, Rakhshani N, Sadr F, Bijarchi R, Joshaghani Y, Mohammadkhani A, et al. Hereditary risk factors for the development of gastric cancer in younger patients. *BMC Gastroenterol*. 2004;4:28.
20. Safaei A, Moghimi-Dehkordi B, Pourhoseingholi MA, Vahedi M, Maserat E, Ghiasi S, et al. Risk of colorectal cancer in relatives: A case control study. *Indian Journal of Cancer*. 2010;47(1):27-30.
21. Barzkar M, Pourhoseingholi MA, Habibi M, Moghimi-Dehkordi B, Safaei A, Pourhoseingholi A, et al. Uninvestigated dyspepsia and its related factors in an Iranian community. *Saudi Med J*. 2009;30(3):397-402.
22. Ji J, Hemminki K. Familial risk for esophageal cancer: an updated epidemiologic study from Sweden. *Clin Gastroenterol Hepatol*. 2006;4(7):840-5.
23. Chen MJ, Wu DC, KO YC, Chiou YY. Personal history and family history as a predictor of gastric cardiac adenocarcinoma risk: a case-control study in Taiwan. *The American journal of gastroenterology*. 2004;99(7):1250-7.
24. Cai L, Zheng ZL, Zhang ZF. Risk factors for the gastric cardia cancer: a case-control study in Fujian Province. *World J Gastroenterol*. 2003;9(2):214-8.
25. Wen D, Wang S, Zhang L, Zhang J, Wei L, Zhao X. Differences of onset age and survival rates in esophageal squamous cell carcinoma cases with and without family history of upper gastrointestinal cancer from a high-incidence area in North China. *Familial cancer*. 2006;5(4):343-52.
26. Garavello W, Negri E, Talamini R, Levi F, Zambon P, Dal Maso L, et al. Family history of cancer, its combination with smoking and drinking, and risk of squamous cell carcinoma of the esophagus. *Cancer Epidemiol Biomarkers Prev*. 2005;14(6):1390-3.
27. Al-Dabbagh TQ. Risk factors associated with esophageal cancer in North of Iran. *Saudi Med J*. 2008;29(1):153.
28. Chang-Claude J, Becher H, Blettner M, Qiu S, Yang G, Wahrendorf J. Familial aggregation of oesophageal cancer in a high incidence area in China. *International journal of epidemiology*. 1997;26(6):1159-65.