



The Effect of Ethanol Baneh Skin Extract on the Expressions of Bcl - 2, Bax, and Caspase - 3 Concentration in Human Prostate Cancer pc3 Cells

Maryam Amiri,¹ Foad Nasrollahi,² Siamak Barghi,³ Nazanin Ebrahimi,⁴ Afsaneh Rajizadeh,⁵ Nasrin Dehghan Nayyeri,³ Faranak Kazerouni,³ Ali Rahimpour,^{3,*} Saeed Namaki,² and Hadis Ahmadi⁶

¹Lecturer, Department of Parasitology, School of Medicine, Jiroft University of Medical Sciences, Jiroft, Iran

²Department of Immunology, Faculty of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

³Department of Laboratory Medicine, Faculty of Paramedical Sciences, Shahid Beheshti University of Medical Sciences, Tehran, Iran

⁴Department of Biochemistry, Medical School, Kerman University of Medical Sciences, Kerman, Iran

⁵MSc of Nutrition in Public Health, Health Services Management Research Center, Institute for Futures Studies in Health, Kerman University of Medical Sciences, Kerman, Iran

⁶Department of Biochemistry, Medical School, Rafsanjan University of Medical Sciences, Rafsanjan, Iran

*Corresponding author: Ali Rahimpour, Department of Laboratory Medicine, Faculty of Paramedical Sciences, Shahid Beheshti University of Medical Sciences, Tehran, Iran. Tel: +98-9126904482, Fax: +98-2122721150, E-mail: a.rahimpour@sbmu.ac.ir

Received 2016 November 30; Revised 2017 February 11; Accepted 2018 February 25.

Abstract

Background: Traditional herbs are effective in the treatment of many diseases. It is proven that traditional herbs can decrease the risk of cancer and grow the survivals of patients.

Objectives: Based on the results of our previous research, ethanol Baneh skin extract has cytotoxic effect on pc3 cells; thus, we decide to determine the effect of Baneh extracts on Bcl2 and Bax expression and amount of concentration of caspase3.

Methods: The expressions of apoptotic - related genes were determined by real time PCR and the concentration of caspase3 was determined by ELISA method.

Results: Ethanol Baneh skin extract increased the mRNA expression of Bax and decreased the mRNA expression of Bcl2 in pc3 cells and also increased the amount of caspase3.

Conclusions: The result of this study indicates the Ethanol Baneh skin extract including photochemical that might be a candidate for the herbal anti - cancer drugs.

Keywords: Expression of Bax and Bcl - 2, Concentration of Caspase3, Prostate Cancer, Ethanol Baneh Skin Extract

1. Background

Prostate cancer is known as one of the most common causes of deaths in men worldwide (1). The incidence rate of prostate cancer in Tehran is more than other provinces in Iran (2). Prostate cancer progresses slowly and metastasis to the bone and lymph nodes. If detected early, prostate cancer can be effectively cured; however, patients mostly refer to the doctors when the disease has already gone to androgen - independent prostate growth; so, androgen - ablative therapies seem to be ineffective in those stages. Some studies are performed to search the novel treatment modalities as well as new targets and anti - cancer agents following the discovery of the clinical impact of advanced prostate cancer (3). Traditional medicine is known as an effective treatment for many diseases; some phytochemicals have been proven to grow the survival of patients and decrease the risk of cancer (4, 5). Most of these medicines have significant anti - cancer as well as apoptosis effects through cellular mechanisms and targeting various

molecular towards cancer (6). Apoptosis is a very important physiological procedure essential for normal growth and preservation of tissue homeostasis (7). Induction of apoptosis in tumor cells by chemotherapeutic agents is controlled by proportion of Bcl2/Bax proteins in the mitochondria (8). Natural products can affect as apoptosis inducer; so, they can reduce cancer incidence by deleting pre-cancerous cells. Induction of cell death in tumor cells is the final target of cancer chemotherapy (9). Iran is the largest manufacturer of wild pistachio in the world. Wild Pistachio is named Baneh in Iran. Baneh trees, *Pistacia atlantica* subsp. *mutica*, from Anacardiaceae family have been growing throughout the central, eastern, and western, parts of Iran (10). Baneh fruit has been used for a diversity of medicinal and nutritional target (11).

2. Objectives

According to the results of our pervious study, Ethanol Baneh skin extract has cytotoxic effect on pc3 cells and also

induces apoptosis in prostate cancer cells (12). In this study, we decided to examine the effect of Baneh skin extract on gene expression of bax and bcl-2 and also caspase3 concentration.

3. Methods

3.1. Materials

We purchased RPMI 1640, Penicillin - Streptomycin, Fetal Bovine Serum (FBS), and Trypsin - EDTA solution from Sigma Aldrich (St. Louis, MO, USA). PC3 prostate cancer cell lines were taken from national cell bank of Iran (Pasteur institute, Iran). Total RNA extraction kit was obtained from GeneALL (Biotech, Korea) and RevertAid First Strand cDNA Synthesis Kit were purchased from Thermo scientific (Thermo Scientific, Schwerte, Germany), respectively. SYBR Green Master Mix was purchased from Takara (Japan). Caspase3 assay kit was obtained from Cusabio Biotech (Wuhan, China).

3.2. Cell Culture

Human prostate cancer cell lines were grown in RPMI 1640 supplemented with 100 mg/mL streptomycin, 100 U/mL penicillin, and 10% fetal bovine serum and in a humidified incubator containing 5% CO₂.

3.3. Preparation of Baneh Extract

Fresh unripe fruit from *P. atlantica kurdica* was gathered from Ilam province, Iran. The fruit skins were dried and, then, powdered, using a mechanical grinder. A total of 10 g of powder was mixed with 50 mL H₂O and 50 mL pure Ethanol for 48 hours and, then, filtrated. The filtrate was evaporated under the reduced pressure in vacuum evaporator and stored at 4 °C. The Baneh extract was dried and dissolved in PBS (1 g in 2 mL) and the filter was sterilized. To prepare the working concentrations, the stock solution of the extract was freshly diluted with RPMI 1640 complete culture medium.

3.4. RNA Extraction and Real Time PCR

Pc-3 cells (5×10^5 cells/well) were seeded in 6-well plate. After 24 hours, the cells were treated with ethanol baneh skin extract (0.78, 1.5, 3.13, 6.25, 12.5 mg/mL) for 48 hours. Then, RNA extraction was performed by kit GeneALL (Biotech, Korea). The concentration and purity was determined, using Nanodrop. For checking the integrity of RNA, extracted RNA was applied on 1% agarose gel to observe 18 s and 28 s bands. Then, using the Revert Aid First Strand cDNA Synthesis kit, 1 µg of total RNA was reverse transcribed to cDNA and Revert Aid Reverse Transcriptase (RT), a recombinant M - MuLV RT. Real time quantitative PCR

(Applied Biosystems, USA) was cycled 40 times between 95°C/15 s and 60 °C/1 min, using SYBR Green PCR Master Mix. Expression of Bcl2 and Bax genes was normalized to the internal control (GAPDH), a housekeeping gene. For analysis of real time PCR, Pfaffl method was used. The relative expression ratio (R) of a target gene was calculated based on E and the CP (CT) deviation of an unknown sample versus a control, and expressed and compared to a reference gene.

Table 1. Real Time PCR Primers Sequences

Name	Sequences (5' -> 3')
bcl - F	CATGTGTGGAGAGCGTCAA
bcl - R	GCCGGTTCAGGTAICTAGTCA
bax - F	GCTGGACATTGGACTTCCTC
BAX - R	TCAGCCCATCTTCTCCAGA
GAPDH - F	TGCACCACCAACTGCTTAG
GAPDH - R	GATGCAGGGATGATGTTTC

3.5. The Measurement of Caspase - 3

Measuring the caspase3 was used by ELISA method according to protocol. (Cusabio Biotech, Wuhan, China).

3.6. Statistical Analysis

Results were expressed as means ± SD and P < 0.05 was considered statistically significant. Linear regression was used for relationship between increasing baneh extract on caspase3 in pc3 cells. Student's t - test was used to analyze the effect of different concentrations of Ethanol baneh extract on gene expression. Statistical analyses were performed, using SPSS version 18.

4. Results

4.1. Ethanol Baneh Skin Extract Increased the Expression of Bax Genes and Decreased the Expression of Bcl2 Gene

The mRNA levels of Bax and Bcl2 were analyzed for determining the apoptotic effects of Ethanol baneh skin extract treatment (0.78, 1.5, 3.13, 6.25, 12.5 mg/mL) on pc3 cells after 48 hours. As shown in [Figure 1](#), baneh extract treatment (48 hours) increased bax gene expression and significant reduction was seen in the expression of Bcl2 as compared with the control. To analyze the effect of different concentrations of Ethanol baneh skin extract on gene expression, the paired t - test was used. A significant difference was seen among gene expression of bax and bcl2 to different concentrations of baneh skin extract (0.78, 1.5, 3.13, 6.25 mg/mL), but there was not any significant difference among gene expression bax and bcl2 to bane skin extract in 6.25, 12.5 mg/mL concentrations.




Figure 1. Real Time PCR Analysis of the Gene Expression of Bcl-2 and Bax on Pc3 Cells After 48 Hours Treatment with Baneh Skin Extract (0.78, 1.5, 3.13, 6.25, 12.5 mg/mL)

4.2. Ethanol Baneh Skin Extract Increased the Concentration of Caspase3

We exposed pc3 cells to different concentrations of the baneh extract (0.78, 1.5, 3.13, 6.25, 12.5 mg/mL) for 24 hours. According to Linear regression, a significant relationship (P -value < 0.05) was seen, as each unit (mg/mL) increased in baneh skin extract concentration, caspase3 concentration 0.03 ng/mL was increased (Table 2 and Figure 2).

Table 2. The Effect of Ethanol Baneh Skin Extract on Caspase3 Concentration. As Each Unit (mg/mL) Increased in Baneh Skin Extract Concentration, Caspase3 Concentration 0.7 ng/mL Was Increased.

Variable	Regression Coefficient	P Value
Baneh concentration	0.03	0.01

5. Discussion

Many treatments such as radiotherapy and chemotherapy induced apoptosis in cells (13, 14). Side effects of many common cancer treatments made the researchers work on the field of herbal medicine (8). According to epidemiological research, strong evidence shows that environmental factors are affected in the occurrence of some types of human cancer (15). Natural anti-oxidants decrease the




Figure 2. Effect of Ethanol Baneh Skin Extract on Caspase3 Concentration in Pc3 Cells

risk of chronic diseases, cancer, diabetes, and cardiovascular disease; so, they are important for human health (16). Deletion of tumors by the infusion of apoptosis is the most common mechanism of cancer therapeutics (17). One of the first regulators of apoptosis is bcl-2 that is known as anti-apoptotic member of bcl-2 family (18). It is encoded as a human proto-oncogene (19) and represses apoptosis by blocking the escape of cytochrome c from the mitochondria to prevent the activation of caspases (18). Damage or stress signals induce Bax monomers oligomerization, and conductance to the release the proapoptotic proteins and activation of some caspases (20). Researchers have recently considered the role of family of caspase proteases (21). Apoptosis is the programmed cell death, in which caspase3 plays upstream “initiator” roles (22, 23). Polyphenolic compounds of plants and flavonoids are well known as dietary anti-oxidants Pistacia species contain source of natural anti-oxidants, flavonoids, and phenolics such as quercetin and α -tocopherol (24). Phenols have an effect on cell cycle arrest and they can activate apoptotic signal transduction pathways in cancerous cells (25). There are a number of medical species in Anacardiaceae family, which contain special compounds with anti-bacterial, fungicidal, and cytotoxic effects. In addition, some studies have approved the cytotoxic activity of the extract of *Lithraea molleoides* (Anacardiaceae) on HepG2 cells. Some examples are the effects of *Semecarpus anacardium* nut oil on growth inhibition in leukemia cell lines and the cytotoxic effects of its nuts on breast cancer cell line (26). Gum mastic (GM) of *P. lentiscus* var *chia* has anti-proliferative effect of on prostate and colon cancer cell lines (3). The flavonoid amount of Baneh (*Pistacia atlantica* sub *kurdica*) extract was more than the extract of the *P. atlantica* Desf

leaves and *P. terebinthus* fruits (27). The amount of DPPH IC50 in the Baneh extract was more than in the DPPH IC50 of *P. atlantica* leaves and the *P. lentiscus* fruits (28). Analyzing the phenolic combination in different part of baneh was found in hull and shell extracts of all genotypes, Baneh from Ilam province has the highest phenolic content and antioxidant activity, for this reason, in this study the skin of bane fruit from Ilam was used (29). Natural compounds have toxic effects and interfere via apoptotic pathways and compounds with pro apoptotic effects intercept cancer incidence by enhancing the deletion of original precancerous cells. As it is known for polar compounds to have anti-cancer effects, generally alcoholic extracts were used for anti-cancer screening (30). Baneh extract indicated cytotoxic effect and apoptosis effect in human breast cancer T47D cells (29) and pericarp polyphenol-rich extract of Baneh induction of apoptosis and cell cycle arrest in human colon carcinoma HT29 cells (27). In previous study, we confirmed that Ethanol baneh skin extract has anti-cancer and inhibitory effect on prostate cancer cells in a time- and dose-dependent manner (11). The molecular analysis of apoptosis proved the activation of caspase3 more efficiently in Baneh than in Doxorubicin treated cancer cells (29). With respect to these results, we decided to examine the effect of baneh skin extract on bcl2 and bax expression and caspase3 concentration for the first time.

In this study, the mRNA expression levels of Bax and Bcl2 genes, which are involved in apoptosis, were investigated. All gene expressions were normalized to GAPDH (reference gene). The results showed that the mRNA levels of bax were increased and the mRNA levels of bcl2 were decreased in pc-3 cells after treatment with Ethanol baneh skin extract for 48 hours; so, the ratio of Bcl-2/Bax was increased. Caspases are important agents in apoptosis. Caspase3 is an executive caspase in both inner and outer pathway. Based on the results of this study, caspase3 concentration in pc3 cells was increased by baneh extract. In summary, Ethanol baneh skin extract could induce apoptosis in pc-3 cells through the modulation of expression of Bax and bcl-2 genes. Baneh extract increased Bax gene expression and decreased the expression of Bcl2, and also increased caspase3 concentration. However, further studies are necessary to understand the exact mechanisms of the activities of baneh skin extract.

Acknowledgments

The authors would like to thank Dr. Ali Hashemi, Faculty of Medical Sciences, Shahid Beheshti University of Medical Sciences and Dr. Ali Rahimpour, Faculty of Paramedical Sciences, Shahid Beheshti University of Medical Sciences for their useful recommendation.

Footnotes

Authors' Contribution: All authors participated in study design, data analysis, interpretation, and manuscript writing.

Conflict of interests: There is no conflict of interest in this article.

Financial Disclosure: This research was funded by a grant from the Shahid Beheshti University of Medical Sciences.

References

1. Siegel RL, Miller KD, Jemal A. Cancer statistics, 2016. *CA Cancer J Clin*. 2016;**66**(1):7-30. doi: [10.3322/caac.21332](https://doi.org/10.3322/caac.21332). [PubMed: [26742998](https://pubmed.ncbi.nlm.nih.gov/26742998/)].
2. Cancer Department. *Iran cancer report*. Center for non-infectious disease control, ministry of health and medical education; 2008.
3. Wang G, Reed E, Li QQ. Apoptosis in prostate cancer: progressive and therapeutic implications (Review). *Int J Mol Med*. 2004;**14**(1):23-34. [PubMed: [15202013](https://pubmed.ncbi.nlm.nih.gov/15202013/)].
4. Issa AY, Volate SR, Wargovich MJ. The role of phytochemicals in inhibition of cancer and inflammation: New directions and perspectives. *J Food Compos Anal*. 2006;**19**(5):405-19. doi: [10.1016/j.jfca.2006.02.009](https://doi.org/10.1016/j.jfca.2006.02.009).
5. Yoon J, Yoo HS, Lee YW, Cho CK. An overview of current oriental medicine herbal cancer research in Korea. *Chin J Integr Med*. 2011;**17**(4):251-6. doi: [10.1007/s11655-011-0710-6](https://doi.org/10.1007/s11655-011-0710-6). [PubMed: [21509666](https://pubmed.ncbi.nlm.nih.gov/21509666/)].
6. Awad AB, Chinnam M, Fink CS, Bradford PG. beta-Sitosterol activates Fas signaling in human breast cancer cells. *Phytomedicine*. 2007;**14**(11):747-54. doi: [10.1016/j.phymed.2007.01.003](https://doi.org/10.1016/j.phymed.2007.01.003). [PubMed: [17350814](https://pubmed.ncbi.nlm.nih.gov/17350814/)].
7. Satyanarayana A, Kaldis P. Mammalian cell-cycle regulation: several Cdk, numerous cyclins and diverse compensatory mechanisms. *Oncogene*. 2009;**28**(33):2925-39. doi: [10.1038/onc.2009.170](https://doi.org/10.1038/onc.2009.170). [PubMed: [19561645](https://pubmed.ncbi.nlm.nih.gov/19561645/)].
8. Kaufmann SH, Earnshaw WC. Induction of apoptosis by cancer chemotherapy. *Exp Cell Res*. 2000;**256**(1):42-9. doi: [10.1006/excr.2000.4838](https://doi.org/10.1006/excr.2000.4838). [PubMed: [10739650](https://pubmed.ncbi.nlm.nih.gov/10739650/)].
9. Reddy L, Odhav B, Bhoola KD. Natural products for cancer prevention: a global perspective. *Pharmacol Ther*. 2003;**99**(1):1-13. [PubMed: [12804695](https://pubmed.ncbi.nlm.nih.gov/12804695/)].
10. Karimi HR, Zamani Z, Ebadi A, Fatahi MR. Morphological diversity of *Pistacia* species in Iran. *Genet Resour Crop Evol*. 2008;**56**(4):561-71. doi: [10.1007/s10722-008-9386-y](https://doi.org/10.1007/s10722-008-9386-y).
11. Hu FB, Willett WC. Optimal diets for prevention of coronary heart disease. *JAMA*. 2002;**288**(20):2569-78. [PubMed: [12444864](https://pubmed.ncbi.nlm.nih.gov/12444864/)].
12. Amiri M, Kazerouni F, Namaki S, Darbandi Tamijani H, Rahimpour H, Boroumand N, et al. Cytotoxic Effects of the Ethanol Bane Skin Extract in Human Prostate Cancer Pc3 Cells. *Iran J Cancer Prev*. 2016;**9**(2):4755. doi: [10.17795/ijcp-4755](https://doi.org/10.17795/ijcp-4755). [PubMed: [27482333](https://pubmed.ncbi.nlm.nih.gov/27482333/)].
13. Pfaffl MW. A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Res*. 2001;**29**(9):45. [PubMed: [11328886](https://pubmed.ncbi.nlm.nih.gov/11328886/)].
14. Galluzzi L, Buque A, Kepp O, Zitvogel L, Kroemer G. Immunogenic cell death in cancer and infectious disease. *Nat Rev Immunol*. 2017;**17**(2):97-111. doi: [10.1038/nri.2016.107](https://doi.org/10.1038/nri.2016.107). [PubMed: [27748397](https://pubmed.ncbi.nlm.nih.gov/27748397/)].
15. Nobili S, Lippi D, Witort E, Donnini M, Bausi L, Mini E, et al. Natural compounds for cancer treatment and prevention. *Pharmacol Res*. 2009;**59**(6):365-78. doi: [10.1016/j.phrs.2009.01.017](https://doi.org/10.1016/j.phrs.2009.01.017). [PubMed: [19429468](https://pubmed.ncbi.nlm.nih.gov/19429468/)].
16. Doll R, Peto R. The causes of cancer: quantitative estimates of avoidable risks of cancer in the United States today. *J Natl Cancer Inst*. 1981;**66**(6):1191-308. [PubMed: [7017215](https://pubmed.ncbi.nlm.nih.gov/7017215/)].

17. Knekt P, Kumpulainen J, Jarvinen R, Rissanen H, Heliovaara M, Reunanen A, et al. Flavonoid intake and risk of chronic diseases. *Am J Clin Nutr*. 2002;**76**(3):560-8. [PubMed: [12198000](#)].
18. Peirce SK, Chen WY. Human prolactin and its antagonist, hPRL-G129R, regulate bax and bcl-2 gene expression in human breast cancer cells and transgenic mice. *Oncogene*. 2004;**23**(6):1248-55. doi: [10.1038/sj.onc.1207245](#). [PubMed: [14647416](#)].
19. Cory S, Adams JM. The Bcl2 family: regulators of the cellular life-or-death switch. *Nat Rev Cancer*. 2002;**2**(9):647-56. doi: [10.1038/nrc883](#). [PubMed: [12209154](#)].
20. Kumar R, Vadlamudi RK, Adam L. Apoptosis in mammary gland and cancer. *Endocr Relat Cancer*. 2000;**7**(4):257-69. [PubMed: [11174847](#)].
21. Debatin KM, Poncet D, Kroemer G. Chemotherapy: targeting the mitochondrial cell death pathway. *Oncogene*. 2002;**21**(57):8786-803. doi: [10.1038/sj.onc.1206039](#). [PubMed: [12483532](#)].
22. Alnemri ES, Livingston DJ, Nicholson DW, Salvesen GS, Thornberry NA, Wong WW, et al. Human ICE/CED-3 protease nomenclature. *Cell*. 1996;**87**(2):171.
23. Stennicke HR, Jurgensmeier JM, Shin H, Deveraux Q, Wolf BB, Yang X, et al. Pro-caspase-3 is a major physiologic target of caspase-8. *J Biol Chem*. 1998;**273**(42):27084-90. [PubMed: [9765224](#)].
24. Slee EA, Harte MT, Kluck RM, Wolf BB, Casiano CA, Newmeyer DD, et al. Ordering the cytochrome c-initiated caspase cascade: hierarchical activation of caspases-2, -3, -6, -7, -8, and -10 in a caspase-9-dependent manner. *J Cell Biol*. 1999;**144**(2):281-92. [PubMed: [9922454](#)].
25. Benhammou N, Bekkara FA, Panovska TK. Antioxidant and antimicrobial activities of the Pistacia lentiscus and Pistacia atlantica extracts. *African Journal of Pharmacy and Pharmacology*. 2008;**2**(2):22-8.
26. Dai J, Mumper RJ. Plant phenolics: extraction, analysis and their antioxidant and anticancer properties. *Molecules*. 2010;**15**(10):7313-52. doi: [10.3390/molecules15107313](#). [PubMed: [20966876](#)].
27. Chakraborty S, Roy M, Taraphdar AK, Bhattacharya RK. Cytotoxic effect of root extract of Tiliacora racemosa and oil of Semecarpus anacardium nut in human tumour cells. *Phytother Res*. 2004;**18**(8):595-600.
28. Gamet-Payraastre L, Li P, Lumeau S, Cassar G, Dupont MA, Chevolleau S, et al. Sulforaphane, a naturally occurring isothiocyanate, induces cell cycle arrest and apoptosis in HT29 human colon cancer cells. *Cancer Res*. 2000;**60**(5):1426-33. [PubMed: [10728709](#)].
29. Hatamnia AA, Abbaspour N, Darvishzadeh R. Antioxidant activity and phenolic profile of different parts of Bene (Pistacia atlantica subsp. kurdica) fruits. *Food Chem*. 2014;**145**:306-11. doi: [10.1016/j.foodchem.2013.08.031](#). [PubMed: [24128482](#)].
30. Mathivadhani P, Shanthi P, Sachdanandam P. Apoptotic effect of Semecarpus anacardium nut extract on T47D breast cancer cell line. *Cell Biol Int*. 2007;**31**(10):198-206. doi: [10.1016/j.cellbi.2007.04.004](#). [PubMed: [17572113](#)].