Published online: 2024 January 25.

Research Article



The Correlation Between GRACE Risk Score and SYNTAX Angiographic Score in Acute Coronary Syndrome: A Cross-Sectional Study

Mohammad Hasan Namazi (1)¹, Isa Khaheshi (1)¹, Maryam Alaei (1)², Yasaman Tavakoli (1)³, Amir Moradi (1)⁴, Omid Amali⁵, Morteza Safi (1)¹, Saeed Alipour Parsa (1)¹, Vahid Eslami (1)¹, Zohre Zamiri (1)^{1,*}

¹ Cardiovascular Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran

² School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

³ Student Research Committee, School of Medicine, Mazandaran University of Medical Sciences, Sari, Iran

⁴ Atherosclerosis Research Center, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

⁵ Student Research Committee, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

corresponding author: Cardiovascular Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran. Email: zohre_zamiri@yahoo.com

Received 2023 November 20; Revised 2024 January 2; Accepted 2024 January 8.

Abstract

Background: The Global Registry of Acute Coronary Events (GRACE) is used in patients with acute coronary syndrome (ACS) to stratify the risk of mortality. The Synergy Between Percutaneous Coronary Intervention (SYNTAX) score explains the extent of coronary artery disease (CAD) and guides to an appropriate treatment strategy.

Objectives: This study aimed to determine the correlation between GRACE and SYNTAX scores.

Methods: A total of 101 ACS patients were recruited in this case-control study. Coronary angiography (CA) was performed for all of the participants. Correlation analysis was performed to investigate the relationship between GRACE risk and SYNTAX angiographic scores.

Results: A total of 83 men and 18 women who had ACS with an average age of 57.2 ± 11.6 years (minimum of 33 and maximum of 89 years) were investigated. The SYNTAX angiographic score and the GRACE risk score for participants of this study were 15.09 \pm 5.87 and 114.47 \pm 26.2, respectively. A strong positive correlation, which was statistically significant, was demonstrated between the GRACE risk score and the SYNTAX angiographic score (r = 0.867, P < 0.001)

Conclusions: Our findings point out a significant strong positive correlation exists between GRACE risk score and SYNTAX angiographic score in patients with unstable angina (UA), ST-elevation myocardial infarction (STEMI), or non-ST elevation myocardial infarction (NSTEMI).

Keywords: Acute Coronary Syndrome, Coronary Angiography, GRACE Score, SYNTAX Score

1. Background

Coronary artery disease (CAD) constitutes the leading cause of death worldwide (1-3). In patients with acute coronary syndrome (ACS), which includes ST elevation myocardial infarction (STEMI), non-ST elevation myocardial infarction (NSTEMI), and unstable angina (UA), early risk stratification is the cornerstone of therapeutic strategies (4, 5). The Global Registry of Acute Coronary Events (GRACE) is recommended by American and European current clinical guidelines to stratify the risk of mortality in-hospital and within 6 months (6, 7). It is measured by clinical variables, including age, systolic blood pressure, heart rate, cardiac arrest, and Killip classification; electrocardiographic parameters, including ST-segment deviation; and laboratory values, including creatinine level and elevated cardiac enzymes (8). Although the GRACE score calculates the risk of recurrent events and mortality, it is not intended for prediction of the severity and complexity of CAD.

The Synergy Between Percutaneous Coronary Intervention (SYNTAX) score, a comprehensive grading system based on coronary angiography (CA) findings, aims to classify the extent and complexity of CAD (9).This score guides in making the decision between

Copyright © 2024, International Journal of Cardiovascular Practice. This open-access article is available under the Creative Commons Attribution-NonCommercial 4.0 (CC BY-NC 4.0) International License (https://creativecommons.org/licenses/by-nc/4.0/), which allows for the copying and redistribution of the material only for noncommercial purposes, provided that the original work is properly cited. performing coronary artery bypass grafting (CABG) surgery or percutaneous coronary intervention (PCI) (10). However, if a non-invasive assessment method of coronary anatomy prior to CA could be developed, the therapeutic decision, intensity, and timing of intervention could be modified.

To date, only a number of studies have investigated the usefulness of GRACE scoring in predicting severe CAD. Most of these studies did not enroll patients with STEMI (11-14). However, the GRACE score can also guide cardiologists to assess the need for an early PCI after fibrinolytic therapy in STEMI patients (15).

2. Objectives

In the present study, we aimed to determine the correlation between the GRACE and SYNTAX scores among patients admitted with ACS (UA, NSTEMI, and STEMI). Therefore, we sought to assess the value of the GRACE score for predicting angiographic features.

3. Methods

This cross-sectional study prospectively enrolled a total of 101 consecutive patients diagnosed with ACS who underwent CA, presenting to the Cardiac Emergency Unit of Shahid Modarres Hospital, Shahid Beheshti University of Medical Sciences, Tehran, Iran, between 2021 and 2023. In our study, we included clinically diagnosed ACS patients who were indicated for CA because they met 1 of the following criteria: (a) patients presenting to the emergency department with chest pain less than 2 months since the onset of symptoms; (b) patients with electrocardiogram (ECG) changes indicative of acute ischemia; and (c) patients with elevated cardiac enzyme levels. There were no exclusion criteria in our study. The study was performed in accordance with the principles of the Declaration of Helsinki (16) and was reviewed and approved by the Ethics Committee of Shahid Beheshti University of Medical Sciences (ID: IR.SBMU.MSP.REC.1400.390). Written informed consent was provided by all ACS patients participating in the study.

Baseline clinical features were documented for each study participant at the time of admission. A checklist was used to assess demographics and cardiovascular risk factors (i.e., smoking, alcohol consumption, dyslipidemia, diabetes mellitus (DM), hypertension, family history, history of previous CAD, history of previous myocardial infarction (MI)). In addition, the parameters used in the GRACE scoring system (i.e., age, systolic blood pressure, heart rate, Killip classification, cardiac arrest in-hospital, elevated cardiac enzymes, creatinine level, ST-segment deviation) were recorded and assessed by a cardiologist (8). The GRACE ACS Risk and Mortality Calculator (available online at www.mdcalc.com) was used to calculate the GRACE risk.

Coronary angiography was performed for all participants in the Shahid Modarres Hospital Catheterization Laboratory. An experienced interventional cardiologist assessed all coronary angiograms visually, blinded to patients' other clinical data. To assess the complexity and extent of coronary artery lesions, the interventionist calculated the SYNTAX score for all participants using the algorithm and calculator as previously described in detail on the website (https://syntaxscore.org).

3.1. Statistical Analysis

Data are reported as mean \pm standard deviation (SD) for continuous variables and as frequency and percentage for categorical variables. Mann-Whitney U test and Wilcoxon test were used for comparison of GRACE and SYNTAX scores between various subgroups of the study (i.e., diabetic and non-diabetic, with and without heart failure, negative and positive family history of CAD, with or without dyslipidemia, smokers, and non-smokers). The chi-square test was used for the comparison of categorical variables between the groups. The Pearson correlation analysis was performed to determine the association between SYNTAX and GRACE scores. We categorized patients based on their GRACE/SYNTAX scores to assess the difference between mean values of systolic blood pressure, heart rate, and creatinine. The GRACE score < 109 was considered lowrisk, $109 \leq \text{GRACE}$ score < 140 was considered intermediate-risk, and the GRACE score \geq 140 was considered high-risk. SYNTAX score < 23 was considered low-risk, $23 \leq$ SYNTAX score < 32 was considered intermediate-risk, and SYNTAX score \geq 32 was considered high-risk. Two-tailed statistical tests were applied, and statistical significance was defined as P values < 0.05. Statistical analysis was performed using SPSS v. 22 (IBM SPSS Statistics for Windows, Armonk, NY: IBM Corp.).

4. Results

A total of 101 consecutive patients (18 (17.8%) women and 83 (82.2%) men) who were diagnosed with ACS (UA, NSTEMI, and STEMI) and underwent CA were recruited for the study. The mean age of the enrolled patients was 57.2 ± 11.6 years. Moreover, 48.5% of patients had a history of MI. In addition, 31.7% and 44.4% of the participants suffered from DM and dyslipidemia, respectively.

Parameters	SYNTAX Score < 23	SYNTAX Score > 23	P-Value
Systolic blood pressure (mmHg)	119.01 ± 18.89	114.78 ± 6.23	0.45
Heart rate (bpm)	81.53 ± 11.12	93.15 ± 3.68	0.04
Creatinine (mg/dL)	1.12 ± 0.2	1.04 ± 0.06	0.28

Besides, 60.4% of these patients were smokers. Of the total number of patients, 49 (48.5%) were classified as low-risk, 38 (37.6%) as intermediate-risk, and 14 (13.9%) as high-risk, based on the GRACE score that was calculated on admission. The mean GRACE score in the study was 114.5 \pm 26.2. According to the determined SYNTAX score, low-complexity CAD (SYNTAX score < 23) was observed in the majority of the participants (85.1%), whilst only 14.9% of participants were classified as intermediate-complexity CAD (23 \leq SYNTAX score < 32), and none of the study participants obtained a SYNTAX score \geq 32. The mean SYNTAX score in our study population was 15.1 \pm 5.9. Baseline clinical and laboratory parameters are reported in Table 1.

This study demonstrated a significant positive correlation between the GRACE score and SYNTAX score (based on angiography) (r = 0.867, P < 0.001) (Figure 1).



Figure 1. Correlation between the GRACE score and the Synergy Between Percutaneous Coronary Intervention (SYNTAX) angiographic score

Amongst various subgroups of the study (i.e., diabetic and non-diabetic, with and without heart failure, negative and positive family history of CAD, with or without dyslipidemia, smokers, and non-smokers), only heart failure seemed to cause a statistically significant difference in GRACE and SYNTAX scores between patients with heart failure and those without heart failure. The median GRACE and SYNTAX scores were both higher in patients with heart failure than those without heart failure (P = 0.002 and P < 0.001, respectively).

5. Discussion

In the present work, we tried to evaluate the correlation between the GRACE and SYNTAX scores among patients admitted with ACS (UA, NSTEMI, and STEMI) in order to assess the value of the GRACE score for predicting angiographic features. Based on our data, the mean GRACE score was 114.5 \pm 26.2. The lowcomplexity CAD (SYNTAX score < 23) was observed in 85.1% of participants, whilst only 14.9% of participants were classified as intermediate-complexity CAD (23 \leq SYNTAX score < 32), and none of the study participants obtained a SYNTAX score \geq 32. Only heart failure seemed to cause a statistically significant difference in GRACE and SYNTAX scores between patients with or without heart failure. Our study demonstrated a significant positive correlation between the GRACE score and the SYNTAX score (based on angiography) in patients with ACS.

A proper treatment strategy can be selected by early identification of CAD complexity since CABG is proven to be beneficial to patients with high and intermediate SYNTAX scores (SYNTAX score \geq 23) (17). Therefore, the GRACE score may serve as a non-invasive method for predicting CAD severity prior to angiography.

To date, a number of studies have investigated the association of GRACE score with the severity of CAD involvements (12-14, 17-25). Consistent with our findings, Rahmani et al. (13) also reported that there is a weak positive correlation between GRACE and SYNTAX scores in UA and NSTEMI patients (r = 0.34, P < 0.001). A similar patient population was investigated by Avci et al. (12), and a positive significant correlation (r = 0.338, P < 0.001) was demonstrated between the GRACE and SYNTAX scores. In a similar study, Hammami et al. (19) found a correlation between these 2 scores, which was weak but significant (r = 0.23, P < 0.001).

To the best of our knowledge, in this regard, only 2 studies included STEMI patients in addition to NSTEMI

and UA patients. In line with our findings, Sofidis et al. (20) found that GRACE and SYNTAX scores in patients with ACS correlated positively. Based on their study, the correlation was weak, although statistically significant (r = 0.32, P < 0.001). In another study with the same population, including STEMI, Bekler et al. (18) demonstrated a positive correlation between these scores as well (r = 0.427, P < 0.001).

Our data provides further evidence of the correlation between GRACE and SYNTAX angiographic score, which is a statistically significant, strong positive correlation (r = 0.867, P < 0.001) and also includes STEMI patients.

Although the GRACE risk score is useful in estimating the risk of mortality in-hospital and within 6 months, our study also indicates that it could represent the severity of coronary artery involvement prior to angiography.

Certain limitations need to be considered in interpreting our findings. Our investigation was conducted at a single center, and a relatively small number of participants were involved. None of our participants obtained a severe SYNTAX score (SYNTAX \geq 32).

5.1. Conclusions

Our findings point out a significant strong positive correlation exists between GRACE score and SYNTAX score patients with UA/NSTEMI/STEMI who were indicated to undergo CA. Our results need to be accredited prospectively by large-scale, multi-center investigations.

Footnotes

Authors' Contribution: Study concept and design: MHN, and ZZ; acquisition of data: IKH, and MHN; analysis and interpretation of data: MA; drafting of the manuscript: YT, and AM; critical revision of the manuscript: OA; statistical analysis: OA; administrative, technical, and material support: MS, and SAP; study supervision: VE.

Conflict of Interests: The authors had no conflict of interest.

Data Availability: The dataset presented in the study is available on request from the corresponding author during submission or after publication.

Ethical Approval: The study was performed in accordance with the principles of the Declaration of Helsinki (16) and was reviewed and approved by the

Ethics Committee of Shahid Beheshti University of Medical Sciences (ID: IR.SBMU.MSP.REC.1400.390).

Funding/Support: There was no funding to be declared. **Informed Consent:** Written informed consent was provided by all ACS patients participating in the study.

References

- Roth GA, Johnson C, Abajobir A, Abd-Allah F, Abera SF, Abyu G, et al. Global, regional, and national burden of cardiovascular diseases for 10 causes, 1990 to 2015. *J Am Coll Cardiol.* 2017;**70**(1):1-25. [PubMed ID: 28527533]. [PubMed Central ID: PMC5491406]. https://doi.org/10.1016/j.jacc.2017.04.052.
- Townsend N, Wilson L, Bhatnagar P, Wickramasinghe K, Rayner M, Nichols M. Cardiovascular disease in Europe: Epidemiological update 2016. Eur Heart J. 2016;37(42):3232-45. [PubMed ID: 27523477]. https://doi.org/10.1093/eurheartj/ehw334.
- No Authors Listed. Corrigendum to: Cardiovascular disease in Europe: Epidemiological update 2016. Eur Heart J. 2019;40(2):189. [PubMed ID: 29846568]. https://doi.org/10.1093/eurheartj/ehy342.
- Yan AT, Yan RT, Tan M, Casanova A, Labinaz M, Sridhar K, et al. Risk scores for risk stratification in acute coronary syndromes: Useful but simpler is not necessarily better. *Eur Heart J*. 2007;28(9):1072-8. [PubMed ID: 17437970]. https://doi.org/10.1093/eurheartj/ehm004.
- Chan Pin Yin D, Azzahhafi J, James S. Risk assessment using risk scores in patients with acute coronary syndrome. *J Clin Med.* 2020;9(9). [PubMed ID: 32967247]. [PubMed Central ID: PMC7565031]. https://doi.org/10.3390/jcm9093039.
- Ibanez B, James S, Agewall S, Antunes MJ, Bucciarelli-Ducci C, Bueno H, et al. 2017 ESC Guidelines for the management of acute myocardial infarction in patients presenting with ST-segment elevation: The task force for the management of acute myocardial infarction in patients presenting with ST-segment elevation of the european society of cardiology (ESC). *Eur Heart J.* 2018;**39**(2):119-77. [PubMed ID: 28886621]. https://doi.org/10.1093/eurheartj/ehx393.
- Anderson JL, Adams CD, Antman EM, Bridges CR, Califf RM, Casey DJ, et al. 2012 ACCF/AHA focused update incorporated into the ACCF/AHA 2007 guidelines for the management of patients with unstable angina/non-ST-elevation myocardial infarction: A report of the American College of Cardiology Foundation/American heart association task force on practice guidelines. J Am Coll Cardiol. 2013;61(23):e179-347. [PubMed ID: 23639841]. https://doi.org/10.1016/j.jacc.2013.01.014.
- Fox KA, Fitzgerald G, Puymirat E, Huang W, Carruthers K, Simon T, et al. Should patients with acute coronary disease be stratified for management according to their risk? Derivation, external validation and outcomes using the updated GRACE risk score. *BMJ Open*. 2014;4(2). e004425. [PubMed ID: 24561498]. [PubMed Central ID: PMC3931985]. https://doi.org/10.1136/bmjopen-2013-004425.
- Sianos G, Morel MA, Kappetein AP, Morice MC, Colombo A, Dawkins K, et al. The SYNTAX Score: An angiographic tool grading the complexity of coronary artery disease. *EuroIntervention*. 2005;1(2):219-27. [PubMed ID: 19758907].
- Serruys PW, Morice MC, Kappetein AP, Colombo A, Holmes DR, Mack MJ, et al. Percutaneous coronary intervention versus coronary-artery bypass grafting for severe coronary artery disease. N Engl J Med. 2009;360(10):961-72. [PubMed ID: 19228612]. https://doi.org/10.1056/NEJMoa0804626.
- 11. Cakar MA, Sahinkus S, Aydin E, Vatan MB, Keser N, Akdemir R, et al. Relation between the GRACE score and severity of atherosclerosis in

acute coronary syndrome. *J Cardiol*. 2014;**63**(1):24-8. [PubMed ID: 24012333]. https://doi.org/10.1016/j.jjcc.2013.06.017.

- Avci BK, Ikitimur B, Tok OO, Cimci M, Erturk E, Omar TB, et al. The role of GRACE score in the prediction of high-risk coronary anatomy in patients with non-ST elevation acute coronary syndrome. *Kardiol Pol.* 2015;73(8):592-7. [PubMed ID: 25733174]. https://doi.org/10.5603/KP.a2015.0030.
- Rahmani R, Majidi B, Ariannejad H, Shafiee A. The Value of the GRACE Score for Predicting the SYNTAX Score in Patients with Unstable Angina/Non-ST Elevation Myocardial Infarction. *Cardiovasc Revasc Med.* 2020;21(4):514-7. [PubMed ID: 31495747]. https://doi.org/10.1016/j.carrev.2019.07.023.
- Beigel R, Matetzky S, Gavrielov-Yusim N, Fefer P, Gottlieb S, Zahger D, et al. Predictors of high-risk angiographic findings in patients with non-ST-segment elevation acute coronary syndrome. *Catheter Cardiovasc Interv.* 2014;83(5):677-83. [PubMed ID: 23784997]. https://doi.org/10.1002/ccd.25081.
- Chotechuang Y, Phrommintikul A, Kuanprasert S, Muenpa R, Ruengorn C, Patumanond J, et al. GRACE score and cardiovascular outcomes prediction among the delayed coronary intervention after post-fibrinolytic STEMI patients in a limited PCI-capable hospital. *Open Heart*. 2020;7(1). e001133. [PubMed ID: 32231784]. [PubMed Central ID: PMC7101037]. https://doi.org/10.1136/openhrt-2019-001133.
- World Medical A. World Medical Association Declaration of Helsinki: Ethical principles for medical research involving human subjects. *JAMA*. 2013;310(20):2191-4. [PubMed ID: 24141714]. https://doi.org/10.1001/jama.2013.281053.
- Mohr FW, Morice MC, Kappetein AP, Feldman TE, Stahle E, Colombo A, et al. Coronary artery bypass graft surgery versus percutaneous coronary intervention in patients with three-vessel disease and left main coronary disease: 5-year follow-up of the randomised, clinical SYNTAX trial. *Lancet.* 2013;**381**(9867):629-38. [PubMed ID: 23439102]. https://doi.org/10.1016/S0140-6736(13)60141-5.
- Bekler A, Altun B, Gazi E, Temiz A, Barutcu A, Gungor O, et al. Comparison of the GRACE risk score and the TIMI risk index in predicting the extent and severity of coronary artery disease in patients with acute coronary syndrome. *Anatol J Cardiol.* 2015;15(10):801-6. [PubMed ID: 25592101]. [PubMed Central ID: PMC5336965]. https://doi.org/10.5152/akd.2014.5802.

- Hammami R, Jdidi J, Mroua F, Kallel R, Hentati M, Abid L, et al. Accuracy of the TIMI and GRACE scores in predicting coronary disease in patients with non-ST-elevation acute coronary syndrome. *Rev Port Cardiol (Engl Ed)*. 2018;**37**(1):41-9. [PubMed ID: 29361323]. https://doi.org/10.1016/j.repc.2017.05.012.
- Sofidis G, Otountzidis N, Stalikas N, Karagiannidis E, Papazoglou AS, Moysidis DV, et al. Association of grace risk score with coronary artery disease complexity in patients with acute coronary syndrome. *J Clin Med.* 2021;**10**(10). [PubMed ID: 34065227]. [PubMed Central ID: PMC8160761]. https://doi.org/10.3390/jcm10102210.
- Akturk E, Askin L, Tasolar H, Turkmen S, Kaya H. Comparison of the predictive roles of risk scores of in-hospital major adverse cardiovascular events in patients with non-ST elevation myocardial infarction undergoing percutaneous coronary intervention. *Med Princ Pract.* 2018;27(5):459-65. [PubMed ID: 29672288]. [PubMed Central ID: PMC6244032]. https://doi.org/10.1159/000489399.
- 22. Barbosa CE, Viana M, Brito M, Sabino M, Garcia G, Maraux M, et al. Accuracy of the GRACE and TIMI scores in predicting the angiographic severity of acute coronary syndrome. *Arq Bras Cardiol.* 2012;**99**(3):818-24. [PubMed ID: 22936027]. https://doi.org/10.1590/s0066-782x2012005000080.
- Santos ES, Aguiar Filho Lde F, Fonseca DM, Londero HJ, Xavier RM, Pereira MP, et al. Correlation of risk scores with coronary anatomy in non-ST-elevation acute coronary syndrome. Arq Bras Cardiol. 2013;100(6):511-7. [PubMed ID: 23598458]. https://doi.org/10.5935/abc.20130088.
- 24. Tasolar H, Cetin M, Balli M, Bayramoglu A, Otlu YO, Turkmen S, et al. CHA2DS2-VASC-HS score in non-ST elevation acute coronary syndrome patients: assessment of coronary artery disease severity and complexity and comparison to other scoring systems in the prediction of in-hospital major adverse cardiovascular events. *Anatol J Cardiol*. 2016;**16**(10):742-8. [PubMed ID: 27025198]. [PubMed Central ID: PMC5324933]. https://doi.org/10.14744/Anatol]Cardiol.2015.6593.
- Ullah M, Saha GK, Ahmed R, Tushar AZ, Saha S, Akanda MAK, et al. Association of GRACE risk score with angiographic severity of coronary artery disease in patients with ST elevation myocardial infarction. *Cardiovascular Journal*. 2015;8(1):30-4. https://doi.org/10.3329/cardio.v8i1.24765.