Published online 2022 June 11.

Gender Differences in the Effect of Resilience Training on Emotional Intelligence in At-Risk Students in Shiraz, Iran

Hassan Joulaei ¹, Faranak Fathi ², Tayebeh Rakhshani ³, Mahin Nazari ⁴, Zahra Hosseinkhani ⁵, Maryam Fatemi ⁶ and Zohre Foroozanfar ⁶

¹Health Policy Research Center, Institute of Health, Shiraz University of Medical Sciences, Shiraz, Iran

²Human Resource Management, Shiraz University of Medical Sciences, Shiraz, Iran

³Department of Public Health, Nutrition Research Center, Shiraz University of Medical Sciences, Shiraz, Iran

⁴Department of Health Promotion, Shiraz University of Medical Sciences, Shiraz, Iran

⁵Metabolic Diseases Research Center, Research Institute for Prevention of Non-communicable Diseases, Qazvin University of Medical Sciences, Qazvin, Iran ⁶HIV/AIDS Research Center, Institute of Health, Shiraz University of Medical Sciences, Shiraz, Iran

*Corresponding author: HIV/AIDS Research Center, Institute of Health, Shiraz University of Medical Sciences, Shiraz, Iran. Email: foroozanfar.zohre@gmail.com

Received 2021 December 14; Revised 2022 April 27; Accepted 2022 April 28.

Abstract

Background: Adolescence is one of the most important and challenging stages in life. psychological, cognitive and emotional changes during this period can expose adolescents to high risk behaviors.

Objectives: The current study examined the effectiveness of resilience training on emotional intelligence (EI) and assessed gender differences in this regard among adolescents living in the outskirts of Shiraz, Iran.

Methods: This pre-post study included 191 students and used an intervention consisting of nine resilience-training sessions. The evaluated outcomes were EI and its 15 components. The paired-samples and independent-samples *t*-tests were used to analyze the data.

Results: Out of 191 students, 88 (46.1%) were boys. Before and after the intervention, the mean EI score for boys was 312.52 ± 37.79 and 327.31 ± 37.75 , while for girls, it was 310.74 ± 30.05 and 312.20 ± 29.51 , respectively. Following the intervention, the scores of boys in problem-solving (P = 0.007), happiness (P = 0.001), emotional self-awareness (P = 0.044), optimism (P = 0.029), self-regard (P = 0.046), impulse control (P = 0.013), and social responsibility (P = 0.042), as well as the total score of EI (P = 0.005), increased significantly. However, only the optimism score (P = 0.004) rose significantly in girls post-intervention. In addition, there were significant differences in the mean of problem-solving (P = 0.006), happiness (P = 0.001), impulse control (P = 0.042), and the total score (P = 0.035) between boys and girls, before and after the intervention.

Conclusions: Resilience training may help moderate high-risk behaviors among adolescents living on the outskirts of Iran's major cities, particularly boys. Based on the gender differences in EI components, it was suggested that female adolescents required significantly more attention. Nonetheless, gender differences in EI components were contentious, and it was concluded that a variety of factors, including the socio-cultural context, may be involved.

Keywords: Emotional Intelligence, Gender Differences, Iran, Resilience, Students

1. Background

As one of the most formative stages of life, adolescence is critical in shaping one's future personality (1). This stage of life can be full of threats, and most high-risk behaviors originate from this age (2). When compared to other stages of life, characteristics such as risk-taking and excitement can lead adolescents to engage in a multitude of high-risk behaviors (3). Marginalized or slum areas are suspected of being a source of such risks (3, 4). Therefore, adolescents living in these areas are more prone to high-risk behaviors (4,5). Numerous studies have sought to identify the factors that can aid humans during traumatic events. Resilience is a critical human ability that enables individuals to adapt to risk factors effectively (6, 7).

In general, resilience is a promising concept studied in social harms, such as violence (8) and high-risk behaviors (9), and encompasses the physical, intellectual, social, emotional, and spiritual domains. The term "resilience" is used in this context to refer to children's positive development in facing adversity and the processes that assist individuals in avoiding harmful, self-destructive, or antisocial behaviors, as well as mental disorders and even threats to their physical well-being (10). In other words, resilience is

Copyright © 2022, Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (http://creativecommons.org/licenses/by-nc/4.0/) which permits copy and redistribute the material just in noncommercial usages, provided the original work is properly cited.

a characteristic of those exposed to specific risk factors and capable of coping with risks and avoiding negative consequences, including delinquency and behavioral problems (11).

In addition, the role of emotion in the development of such behaviors should not be overlooked (12). Emotional intelligence (EI) is the capacity to manage one's emotions and develop the necessary skills to influence interpersonal relationships and day-to-day problem-solving situations (13). In this regard, prior research has focused on the associations between EI and delinquent behaviors in students, such as crime, substance abuse, vandalism, pornography, and sexual behaviors (12). Managing emotions is associated with coping strategies, and deficits in those abilities are accompanied by risk-taking behaviors (14).

Furthermore, several studies have examined the relationship between EI and resilience, concluding that resilience training may boost EI (15, 16). Subsequently, higher levels of EI have been linked to increased resilience (15). Resilience-promoting interventions have been introduced as one of the successful programs for vulnerable students in identifying the factors that contribute to some adolescents resisting or adapting to risks (10, 17).

A study in Iran discovered that the outskirts of major cities, including Shiraz, created an environment conducive to crime and social harm due to poverty, partial deprivation, and low quality of life (18). Moreover, two studies conducted on marginalized populations and slum dwellers have highlighted the significant risks associated with living in such areas, the role of poverty in aggravating social harms, and a higher prevalence of high-risk behaviors among residents (18, 19). However, harm does not merely affect everyone in the same way. Some individuals may be resilient, while others are more vulnerable to potential harm (20).

2. Objectives

To our knowledge, little research has been conducted on the relationship between EI and resilience training among adolescents, and gender differences have not been considered in the marginalized areas of Iran (21, 22). Therefore, the purpose of this study was to evaluate the impact of resilience training on improving EI in adolescents and its effectiveness in both genders in the marginalized areas of Shiraz, Iran.

3. Materials and Methods

3.1. Study Setting and Design

This pre-experimental or before-and-after study was conducted in 2019 in the middle schools located in Shiraz's

slum areas. Shiraz is the capital city of Fars province in southwestern Iran, with a population of nearly 1,869,001 and over 200.000 inhabitants living in its urban slums (23). Before-after studies are applied in small-scale interventional investigations in low-resource settings (24, 25). The study intervention consisted of nine 90-min resiliencetraining sessions, focusing on the 15 components of EI (26). All interactive training sessions were supervised and coordinated by knowledgeable psychologists. Almost all students were engaged in educational subjects through problem- and story-based learning. Table 1 lists the training subjects for each session.

3.2. Sample Size and Sampling

The participants were selected using a multistage cluster sampling procedure. We began by randomly selecting one slum area in Shiraz from a total of thirteen. Next, we considered three middle schools in that area, one for girls and two for boys. Finally, the study included all first-grade middle school students. Students were free to leave at any point during the study, and those who were not willing to participate in the study or were absent in the post-test were excluded from the study. The educational sessions drew a total of 191 male and female students (out of 221). They were also asked to sign an informed consent form prior to the intervention if they wished to participate. All students voluntarily participated in the study based on announcements by schools' managers.

3.3. Ethical Approval

The Ethics Committee of Shiraz University of Medical Sciences approved this study under the code IR.SUMS.REC.1398.1084.

3.4. Measurement

The modified Bar-On Emotional Quotient Inventory (EQ-i) was used in this study to assess the outcomes of the resilience-training intervention. The questionnaire has been validated and translated into at least 30 different languages (26). The questionnaire consists of 90 items based on a five-point Likert scale and 15 subscales, including intrapersonal skills (i.e., self-regard, emotional self-awareness, assertiveness, independence, and self-actualization), interpersonal skills (i.e., empathy, social responsibility, and interpersonal relationships), stress management skills (i.e., stress tolerance and impulse control), adaptability (i.e., reality testing, flexibility, and problem-solving), and self-actualization (i.e., optimism and happiness) (26). Samouei evaluated and confirmed the validity and reliability of the questionnaire in 2002 (27). Cronbach's alpha coefficient (α = 0.93) was used to assess

| Session | Target | Training Topic | Training Method | Educator | | | |
|---------|--|--|--|--|--|--|--|
| 1 | Interpersonal relationships, self-esteem, and self-expression | Learning about communication and social and family ties | Educational videos, lectures, question, and answer | Psychological and research expert group | | | |
| 2 | Endure psychological stress | Learning about enduring stress, shaking control | Lectures, group discussions, and question and answer | Psychiatrist | | | |
| 3 | Problem-solving skills and challenging beliefs | Learning to solve problems | Educational videos and posters | Consultant | | | |
| 4 | Self-actualization, empathy, optimism, and happiness | Learning about self-actualization, empathy, optimism, and happiness | Lectures, group discussions, and question and answer | Mental health expert | | | |
| 5 and 6 | Emotion adjustment and resilience | Learning about excitement and excitement and training on stress management and anger management | Lectures, group discussions, and question and answer | Research fellow | | | |
| 7 and 8 | Flexibility, self-esteem, and confidence | Learning about momentary resilience | Lectures, group discussions, and question and answer | Research fellow | | | |
| 9 | Perspective and goal setting for life | Learning about social and individual adequacy | Lectures, group discussions, and question and answer | Research fellow | | | |
| | | | | | | | |

Variables

Gender

Male

Female

Number of children

One

Two

the reliability of the questionnaire (27). The EQ-i was completed by study participants prior to and two months after the intervention.

3.5. Data Analysis

The quantitative variables were described as mean \pm standard deviation, and the qualitative ones were reported as numbers and percentages. First, the Kolmogorov-Smirnov test was used to determine the normality of the data. To compare the mean differences of the EI components before and after the intervention between genders, we used independent-samples *t*-tests. Moreover, to compare the pre- and post-intervention mean scores of the EI components, a paired-samples t-test was utilized. Data were analyzed by the SPSS software version 20 and Graph-Pad Prism software version 9. The significance level for all tests was considered P-value < 0.05.

4. Results

This study surveyed a total of 191 students, of whom 88 (46.1%) were boys. Nearly half of the students (49.2%) were only child, and 86.9% lived with both parents. Moreover, most mothers (30.4%) held a high school diploma, and fathers (38.7%) held a middle school education. Table 2 summarizes the socio-demographic characteristics of the participants.

Overall, the intervention increased the mean resilience scores by 8.11 points (14.19 points in boys and 2.92 points in girls). Table 3 compares the EI components before and after the intervention in both genders. Scores on all subscales had a range of 6 - 30. Following the intervention, the scores

Three or more 48 (25.1) Living with parents Living with both 166 (86.9) Living with mother 20 (10.5) Living with father 5 (2.6) Mother's education Illiterate 19 (9.9) Elementary 48 (25.1) Intermediate 56 (29.3) Diploma 58 (30.4) College education 10 (5.2) Father's education Illiterate 13 (6.8) Elementary 46 (24.1) Intermediate 74 (38.7) Diploma 42 (22) College education 16 (8.4)

Table 1. Training Intervention Program

 Table 2. Description of the Frequency of Demographic Variables of the Studied Individuals

No. (%)

88 (46.1)

103 (53.9)

94 (49.2)

49 (25.7)

of boy participants in problem-solving (P = 0.007), happiness (P = 0.001), emotional self-awareness (P = 0.044), optimism (P = 0.029), self-regard (P = 0.046), impulse control (P = 0.013), and social responsibility (P = 0.042), as well as the total EI score (P = 0.005), rose significantly. However, only the optimism score (P = 0.004) augmented significantly in girls post-intervention, while the other components and the total EI score did not show a significant difference.

Table 4 compares the differences in mean EI components before and after the intervention between the two genders. According to the findings of the current study, there was a statistically significant difference between boys and girls in the mean values of problem-solving (P = 0.006), happiness (P = 0.001), impulse control (P = 0.042), and the total score (P = 0.035). Furthermore, the optimism score had the most significant increase following the intervention in both genders.

Figure 1 illustrates the comparison of the EI components between boy and girl participants pre-intervention. It shows significant differences between boys and girls in terms of independence (P=0.01), reality testing (P=0.021), flexibility (P=0.028), and empathy (P=0.045) at this time. However, prior to the intervention, the total EI score did not differ significantly between genders (P=0.718). Figure 2 compares the EI components between boy and girl participants post-intervention. Significant differences were observed at this time between boys and girls in problem-solving (P = 0.024), happiness (P = 0.001), independence (P=0.006), and flexibility (P=0.05). Both genders had a statistically significant difference in their total EI score following the intervention (P-value = 0.004).

5. Discussion

The present research aimed to evaluate gender differences in the effectiveness of resilience training on EI among adolescents living in the marginalized areas of Shiraz in 2019. Our findings indicated that the changes in EI components, including problem-solving, happiness, impulse control, self-awareness, optimism, and self-regard, as well as the total score, were statistically significant in male adolescents following the intervention.

However, no statistically significant difference was observed in girls when the mean difference of pre- and post-test EI components was compared, except for optimism. Furthermore, there was a significant difference in problem-solving, happiness, impulse control, and total score between the genders (27). The total score of EI in male adolescents was higher than in females. The possible interpretation for such difference would be the higher rate of risk-taking behaviors in male students in Iran than the females (28), which increased male students' demands for developing resilience skills.

In addition, the results of the present study established the efficacy of resilience training on EI, consistent with previous research (15, 16, 21). In fact, resilience may help to alleviate emotional exhaustion and promote overall wellbeing (29). In line with previous research, this study established the efficacy of resilience-promoting interventions in vulnerable adolescents living in marginalized areas in both genders (10, 17). Furthermore, our findings indicated that resilience-training interventions only influenced adolescent social responsibility, self-regard, and emotional self-awareness as intrapersonal and interpersonal skills. Unlike the current study. Ghahramani et al. discovered no significant difference in emotional self-awareness between male and female adolescents (30). Molero Jurado et al. also reported that girls performed better than boys, but boys had a greater capacity for engagement (31).

Evidence indicates that resilience-building interventions can reduce the stress levels of students by improving their coping and stress management skills (32). Resilience significantly improved impulse control and stress management skills in male adolescents in the current study. In other words, numerous EI components were genderspecific (33-35). Molero Jurado et al. also noted a significant interaction between gender and stress management skills, indicating that no interaction, which is necessary for overcoming obstacles, was observed in male adolescents (31). The general mood and its components were classified as EI components associated with resilience. The mean EI score increased significantly before and after the intervention, particularly among boys.

In general, happy and optimistic people have lower stress levels and respond more positively and adaptively to specific situations (36), which reduces the likelihood of being engaged in high-risk behaviors (37). Although gender differences in mental health problems have been debated in previous research (30, 38-40), girls may experience depression at a higher rate than boys in puberty conditions (23).

In this study, male adolescents achieved a higher mean score in several EI components both pre- and postintervention, and their overall EI score was higher than female. Based on these findings, it was concluded that resilience and EI might develop differently depending on gender. Individuals with a high EI are unquestionably capable of correctly comprehending and evaluating their emotional states, adjusting their moods, and knowing how and when to express their emotions. These abilities facilitate successful coping and frequently result in increased reliance on social support, advancement motivation, and life satisfaction (16).

| Table 3. Comparison of the Pre- and Post-intervention Emotional Intelligence Components in Both Genders ^{a, b} | | | | | | | | | |
|---|------------------|--------------------|--------------------|------------------|-------------------|---------|--|--|--|
| | | Male | | Female | | | | | |
| Emotional Intelligence Components | Pre-intervention | Post-intervention | P-Value | Pre-intervention | Post-intervention | P-Value | | | |
| Problem solving | 24.01 ± 4.05 | 25.34 ± 3.50 | 0.007 ^c | 24.50 ± 3.89 | 24.00 ± 4.48 | 0.253 | | | |
| Happiness | 18.39 ± 4.94 | 19.62 ± 17.61 | 0.001 ^c | 17.94 ± 20.46 | 16.61 ± 2.70 | 0.196 | | | |
| Independence | 18.37 ± 4.94 | 19.55 ± 4.61 | 0.054 | 16.64 ± 4.26 | 17.41 ± 4.05 | 0.063 | | | |
| Stress tolerance | 19.76 ± 4.34 | 20.30 ± 4.22 | 0.386 | 19.72 ± 4.31 | 19.31 ± 4.61 | 0.397 | | | |
| Self-actualization | 18.50 ± 6.67 | 19.87 ± 5.91 | 0.074 | 18.00 ± 6.69 | 18.04 ± 6.97 | 0.931 | | | |
| Emotional self-awareness | 20.60 ± 3.50 | 21.56 ± 3.89 | 0.044 ^c | 21.0 ± 3.65 | 20.89 ± 3.21 | 0.778 | | | |
| Reality testing | 18.46 ± 4.51 | 18.97 ± 4.31 | 0.378 | 17.16 ± 3.14 | 17.49 ± 2.96 | 0.374 | | | |
| Interpersonal relationship | 19.61 ± 4.86 | 20.00 ± 4.25 | 0.551 | 18.92 ± 4.35 | 19.69 ± 4.10 | 0.104 | | | |
| Optimism | 23.19 ± 4.86 | 24.79 ± 7.09 | 0.029 ^c | 23.99 ± 3.66 | 24.84 ± 4.02 | 0.004* | | | |
| Self-regard | 22.05 ± 3.44 | 22.94 ± 3.39 | 0.046 ^c | 22.18 ± 3.33 | 22.26 ± 3.71 | 0.839 | | | |
| Impulse control | 23.54 ± 3.54 | 24.59 ± 3.34 | 0.013 ^c | 24.04 ± 3.24 | 23.99 ± 2.99 | 0.869 | | | |
| Flexibility | 20.01 ± 3.85 | 20.65 ± 4.39 | 0.273 | 18.80 ± 3.64 | 19.49 ± 3.89 | 0.103 | | | |
| Social-responsibility | 23.50 ± 3.83 | 24.44 ± 3.66 | 0.042 ^c | 23.93 ± 3.70 | 23.88 ± 3.78 | 0.868 | | | |
| Empathy | 23.31 ± 4.46 | 24.25 ± 4.27 | 0.071 | 24.60 ± 4.30 | 24.67 ± 4.25 | 0.883 | | | |
| Assertiveness | 19.17 ± 4.02 | 19.56 ± 4.48 | 0.420 | 19.28 ± 3.22 | 19.64 ± 3.11 | 0.384 | | | |
| Total score | 312.52 ± 37.79 | 327.31 ± 37.75 | 0.005 ^c | 310.74 ± 30.05 | 312.20 ± 29.51 | 0.246 | | | |

^a Data reported as mean \pm SD. ^b Paired samples *t*-test

^c Significant at 0.05



Figure 1. Emotional intelligence comparison between males and females before intervention



Figure 2. Emotional intelligence comparison between males and females after intervention

| | Male | Female | P-Value |
|----------------------------|-------------------|------------------------------------|--------------------|
| Problem solving | 1.27 ± 4.33 | $\textbf{-0.51} \pm \textbf{4.45}$ | 0.006 ^c |
| Happiness | 1.22 ± 3.41 | $\textbf{-0.33} \pm \textbf{2.57}$ | 0.001 ^c |
| Independence | 1.18 ± 5.66 | 0.77 ± 4.11 | 0.572 |
| Stress tolerance | 0.54 ± 5.65 | $\textbf{-0.41} \pm \textbf{4.89}$ | 0.212 |
| Self-actualization | 1.37 ± 7.13 | 0.04 ± 5.70 | 0.155 |
| Emotional self-awareness | 0.96 ± 4.43 | $\textbf{-0.11} \pm \textbf{3.82}$ | 0.074 |
| Reality testing | 0.51 ± 5.41 | 0.33 ± 3.75 | 0.786 |
| Interpersonal relationship | 0.38 ± 6.04 | 0.77 ± 4.80 | 0.620 |
| Optimism | 1.60 ± 6.77 | 0.85 ± 2.97 | 0.313 |
| Self-regard | 0.88 ± 4.11 | $\textbf{-0.07} \pm \textbf{3.85}$ | 0.164 |
| Impulse control | 1.04 ± 3.85 | $\textbf{-0.05} \pm \textbf{3.57}$ | 0.042 ^c |
| Flexibility | 0.64 ± 5.50 | 0.68 ± 4.24 | 0.953 |
| Social-responsibility | 0.94 ± 4.28 | $\textbf{-0.04} \pm \textbf{2.95}$ | 0.061 |
| Empathy | 0.93 ± 4.4776 | 0.07 ± 3.72 | 0.167 |
| Assertiveness | 0.48 ± 5.65 | 0.35 ± 3.86 | 0.852 |
| Total score | 14.19 ± 45.98 | 2.92 ± 25.29 | 0.035 ^c |

Table 4. Comparison of the Difference in Mean Emotional Intelligence Components Before and After the Intervention Between Genders ^{ab}

^a Data reported as mean \pm SD.

^b Independent samples *t*-test

^c Significant at 0.05

As a result, gender differences in EI components may indicate that female adolescents required significantly more attention. Nonetheless, gender differences in EI subscales were contentious, and it could be concluded that a variety of factors, including the socio-cultural context, may be involved.

5.1. Strengths and Weaknesses

The current study had several strengths and weaknesses. This was the first attempt, given the prevalence of resilience-training interventions in Iran's marginalized adolescent population. Moreover, this investigation was one of the few studies that have examined gender differences in EI in adolescents. However, one of its shortcomings was the lack of long-term follow-up on the training programs implemented. In addition, self-administration was a weakness of this study, resulting in measurement bias and the misclassification of gender groups. Another limitation was the absence of a control group due to financial constraints and a small population.

5.2. Conclusions

The findings of the current study indicated that resilience training could significantly improve EI and its components and provides critical benefits for dealing with adversity, particularly in male adolescents. Given that the EI scores of boys were higher than female adolescents, the optimal solution for enhancing various components of EI would require reasonable intervention in families and schools.

With EI training programs, social skills and resilience may be valuable tools for reducing behavioral problems and emotional distress in students and an excellent opportunity to improve social adjustment. As a result, developing resilience as a part of educational programs enables atrisk students to better prepare for dealing with problems, adapt to challenges, and manage their reactions, particularly boys. This study suggested that resilience training in schools for adolescents living in the marginalized areas of large cities, such as Shiraz, could help them improve their EI skills. Furthermore, it would empower them in facing adversity, preventing or mitigating their participation in high-risk behaviors and their likely consequences. Finally, we recommend further investigation with control groups to adjust the confounding factors.

Footnotes

Authors' Contribution: Hassan Joulaei: Substantial contributions to the conceptualization and design of the work and supervised all the phases of the study and manuscript writing. Faranak Fathi, Mahin Nazari, and Tayebeh Rakhshani: The acquisition, analysis, and interpretation of data for the work and writing the manuscript. Zahra Hosseinkhani and Maryam Fatemi: Methodology supervising and final approval of the manuscript. Zohre Foroozanfar: Data acquisition, manuscript drafting, and critically revising for important intellectual content and final approval of the manuscript.

Conflict of Interests: None.

Data Reproducibility: The authors did not declare it.

Ethical Approval: The Ethics Committee of Shiraz University of Medical Sciences approved this study under the code IR.SUMS.REC.1398.1084.

Funding/Support: The authors would like to extend their gratitude to the Vice Chancellor's Office for Research at Shiraz University of Medical Sciences, Shiraz, Iran, for their financial support (Grant no. 18249). We also wish to thank all Education Departments in Fars Province for their cooperation and all students participating in this study.

Informed Consent: Students were asked to sign an informed consent form prior to the intervention if they wished to participate. All students voluntarily participated in the study based on announcements by schools' managers.

References

- 1. Berk LE, Petersen A. Development through the lifespan. Allyn and Bacon: Boston, United States; 2004.
- 2. World Health Organization. *Health for the world's adolescents: a second chance in the second decade: summary*. Geneva, Switzerland: World Health Organization; 2014.
- Auerswald CL, Piatt AA, Mirzazadeh A; Unicef. Research with disadvantaged, vulnerable and/or marginalized adolescents. New York, New York, United States: UNICEF; 2017.
- Mahmoodi Z, Solimannejad T, Solimannejad M, Niazi M. Social harms and social class in Iranian youth and adolescents living in slums in the suburbs: a systematic review. *Int J Adolesc Med Health*. 2021;33(3):1– 7. doi: 10.1515/ijamh-2020-0267. [PubMed: 33581002].
- Goldblatt A, Kwena Z, Lahiff M, Agot K, Minnis A, Prata N, et al. Prevalence and Correlates of HIV Infection among Street Boys in Kisumu, Kenya. *PLoS One*. 2015;**10**(10). e0140005. doi: 10.1371/journal.pone.0140005. [PubMed: 26461494]. [PubMed Central: PMC4604137].
- Rew L, Horner SD. Youth Resilience Framework for reducing healthrisk behaviors in adolescents. J Pediatr Nurs. 2003;18(6):379–88. doi: 10.1016/s0882-5963(03)00162-3. [PubMed: 15058534].
- Prince-Embury S. Risk Behavior and Personal Resiliency in Adolescents. Can J Sch Psychol. 2015;30(3):209–17. doi: 10.1177/0829573515577601.
- Dekel R, Monson CM. Military-related post-traumatic stress disorder and family relations: Current knowledge and future directions. *Aggress. Violent Beh.* 2010;15(4):303–9. doi: 10.1016/j.avb.2010.03.001.
- Cuomo C, Sarchiapone M, Giannantonio MD, Mancini M, Roy A. Aggression, impulsivity, personality traits, and childhood trauma of prisoners with substance abuse and addiction. *Am J Drug Alcohol Abuse*. 2008;**34**(3):339–45. doi: 10.1080/00952990802010884. [PubMed: 18428076].
- Richman JM, Fraser MW. The context of youth violence: Resilience, risk, and protection. Westport, Connecticut, United States: Greenwood Publishing Group; 2001.
- Newsome J, Sullivan CJ. Resilience and vulnerability in adolescents: genetic influences on differential response to risk for delinquency. *J Youth Adolesc*. 2014;43(7):1080–95. doi: 10.1007/s10964-014-0108-9. [PubMed: 24585391].
- 12. Chong AM, Lee P, Roslan S, Baba M. Emotional Intelligence and At-Risk Students. *SAGE Open*. 2015;**5**(1). doi: 10.1177/2158244014564768.
- Sanchez-Alvarez N, Extremera N, Fernandez-Berrocal P. Maintaining Life Satisfaction in Adolescence: Affective Mediators of the Influence of Perceived Emotional Intelligence on Overall Life Satisfaction Judgments in a Two-Year Longitudinal Study. *Front Psychol.* 2015;6:1892. doi: 10.3389/fpsyg.2015.01892. [PubMed: 26834654]. [PubMed Central: PMC4714630].
- Cobos-Sánchez L, Flujas-Contreras JM, Gómez-Becerra I. [The role of emotional intelligence in psychological adjustment among adolescents]. An Psicol-Spain. 2017;33(1):66–73. Spanish.
- Saffari M, Shojaeizadeh D, Ghofranipour F, Heydarnia A, Pakpour A. Health education & promotion-theories, models & methods. Tehran, Iran. Sobhan Pub; 2009.
- Molero Jurado M, Pérez-Fuentes M, Barragán Martín AB, del Pino Salvador RM, Gázquez Linares JJ. Analysis of the Relationship between Emotional Intelligence, Resilience, and Family Functioning in Adolescents' Sustainable Use of Alcohol and Tobacco. *Sustainability*. 2019;**11**(10). doi: 10.3390/su11102954.
- Ungar M, Russell P, Connelly G. School-Based Interventions to Enhance the Resilience of Students. *J Educ Develop Psychol*. 2014;4(1). doi: 10.5539/jedp.v4n1p66.
- Fouladiyan M. Studying crimes and social harms in Mashhad suburbs and evaluating their simplifying factors. *Journal of the society of Iran's* social problems. 2019;8(1).

- Rasekh K, Allapanazadeh T. Social factors affecting on drugs abuse:(Slum dwellers in Shiraz-Iran). Sociological Studies of Youth. 2012;3(7):25-42.
- Barrett DH, Ortmann LH, Dawson A, Saenz C, Reis A, Bolan G. Chapter 7: Vulnerability and Marginalized Populations. *Public Health Ethics: Cases Spanning the Globe*. London, UK: Springer Nature; 2016. doi: 10.1007/978-3-319-23847-0.
- Adibsereshki N, Hatamizadeh N, Sajedi F, Kazemnejad A. The Effectiveness of a Resilience Intervention Program on Emotional Intelligence of Adolescent Students with Hearing Loss. *Children (Basel)*. 2019;6(3). doi: 10.3390/children6030048. [PubMed: 30901951]. [PubMed Central: PMC6463041].
- Mousavi SV, Pasandideh Z, Poorabolghasem Hosseini S. The Effectiveness of emotional intelligence training on risk taking of adolescent girls. J Pediatr Nurs. 2017;4(2):1–8. doi: 10.21859/jpen-04021.
- Sweeting H, West P. Sex differences in health at ages 11, 13 and 15. Soc Sci Med. 2003;56(1):31–9. doi: 10.1016/s0277-9536(02)00010-2. [PubMed: 12435549].
- Varkevisser CM, Pathmanathan I, Brownlee AT. Designing and conducting health systems research projects. Ottawa, Canada: International Development Research Centre; 2003.
- Sedgwick P. Before and after study designs. *BMJ*. 2014;349:g5074. doi: 10.1136/bmj.g5074. [PubMed: 25106742].
- Bar-On R. The Bar-On model of emotional-social intelligence (ESI). Psicothema. 2006;18 Suppl:13–25. [PubMed: 17295953].
- Samouei R. [Standardization Emotional intelligence Questionnaire.Tehran, Iran]. Tehran, Iran: Ravan Tajhiz Sina Publication; 2002. Persian.
- Bahramnejad A, Iranpour A, Nakhaee N. Gender-based differences in risk-taking behaviors among high school students in Southeast Iran. *Int J Adolesc Med Health.* 2020;**33**(6):437–43. doi: 10.1515/ijamh-2019-0205. [PubMed: 32549159].
- Bande B, Fernández-Ferrín P, Varela JA, Jaramillo F. Emotions and salesperson propensity to leave: The effects of emotional intelligence and resilience. *Ind Market Manag.* 2015;44:142–53. doi: 10.1016/j.indmarman.2014.10.011.
- Ghahramani S, Jahromi AT, Khoshsoroor D, Seifooripour R, Sepehrpoor M. The relationship between emotional intelligence and happiness in medical students. *Korean J Med Educ*. 2019;**31**(1):29– 38. doi:10.3946/kjme.2019.116. [PubMed: 30852859]. [PubMed Central: PMC6589629].
- 31. Molero Jurado MDM, Perez-Fuentes MDC, Barragan Martin AB,

Gazquez Linares JJ, Oropesa Ruiz NF, Simon Marquez MDM. Emotional Intelligence Components as Predictors of Engagement in Nursing Professionals by Sex. *Healthcare (Basel)*. 2020;**8**(1). doi: 10.3390/healthcare8010042. [PubMed: 32098453]. [PubMed Central: PMC7151152].

- Vesel L, Waller K, Dowden J, Fotso JC. Psychosocial support and resilience building among health workers in Sierra Leone: interrelations between coping skills, stress levels, and interpersonal relationships. *BMC Health Serv Res.* 2015;**15 Suppl 1**. S3. doi: 10.1186/1472-6963-15-S1-S3. [PubMed: 26062735]. [PubMed Central: PMC4464206].
- 33. Petrides KV, Sanchez-Ruiz MJ, Siegling AB, Saklofske DH, Mavroveli S. Emotional Intelligence as Personality: Measurement and Role of Trait Emotional Intelligence in Educational Contexts. In: Keefer KV, Parker JDA, Saklofske DH, editors. Emotional Intelligence in Education: Integrating Research with Practice. Manhattan, New York City, NY, USA: Springer International Publishing; 2018. p. 49–81. doi: 10.1007/978-3-319-90633-1_3.
- Esnaola I, Revuelta L, Ros I, Sarasa M. [Desarrollo de la Inteligencia Emocional en la adolescencia: un estudio transversal y longitudinal]. An Psicol-Spain. 2017;33(2). Spanish. doi: 10.6018/analesps.33.2.251831.
- Freudenthaler HH, Neubauer AC, Haller U. Emotional Intelligence. J Individ Differ. 2008;29(2):105–15. doi: 10.1027/1614-0001.29.2.105.
- 36. Khordzanganeh F, Heidarie AR, Naderi F. The relationship between emotional intelligence, happiness and resiliency with mental health in male and female high school students of ramhormoz city. *Indian Journal of Fundamental and Applied Life Sciences*. 2014;**4**(2):633–41.
- Conversano C, Rotondo A, Lensi E, Della Vista O, Arpone F, Reda MA. Optimism and its impact on mental and physical well-being. *Clin Pract Epidemiol Ment Health*. 2010;6:25–9. doi: 10.2174/1745017901006010025. [PubMed: 20592964]. [PubMed Central: PMC2894461].
- Hosseinkhani Z, Parsaeian M, Hassanabadi HR, Khoshkchali A, Alinesaei Z, Nedjat S. Mental Health Problems and the Associated Family and School Factors in Adolescents: A Multilevel Analysis. *Iran J Psychiatry*. 2021;**16**(1):13–20. doi: 10.18502/ijps.v16i1.5373. [PubMed: 34054978]. [PubMed Central: PMC8140304].
- Furnham A, Christoforou I. Personality Traits, Emotional Intelligence, and Multiple Happiness. NAm J Psychol. 2007;9(3).
- 40. Sasanpour M, Khodabakhshi M, Nooryan K. The relationship between emotional intelligence, happiness and mental health in students of medical sciences of Isfahan University. Int J Collab Res Intern Med Public Health. 2012;4(9):0.