



A Truth or a Rumor: Effects of Addictive Substances on Prevention of COVID-19; an Investigation of Homeless Drug Abusers in Southeastern Iran

Hassan Okati-Aliabad ¹, Mahdi Mohammadi ¹, Alireza Salimi Khorashad ², Alireza Ansari-Moghaddam ¹, Mohsen Hossein Bor ³, Jalil Nejati ^{1,*}

¹ Health Promotion Research Center, Zahedan University of Medical Sciences, Zahedan, Iran

² Department of Parasitology, Research Center for Infectious and Tropical Disease, Zahedan University of Medical Sciences, Zahedan, Iran

³ Department of Midwifery, School of Nursing and Midwifery, Iranshahr University of Medical Sciences, Iranshahr, Iran

* Corresponding author: Health Promotion Research Center, Zahedan University of Medical Sciences, Zahedan, Iran. Email: jalilnejati@yahoo.com

Received 2023 November 21; Revised 2024 January 8; Accepted 2024 January 9.

Abstract

Background: In a crisis, rumors often get started, and societies use them to cope with ambiguous situations. Southeastern Iran, with its high rate of homeless addicts per population, is a suitable area to examine the accuracy of the rumor about the positive effect of addictive substances on preventing COVID-19.

Objectives: This cross-sectional study aimed to investigate the abundance of COVID-19 infection among homeless drug abusers in this area.

Patients and Methods: For each participant, the health experts completed a questionnaire including demographic characteristics, symptoms and type, frequency and methods of drug abuse. For a definite diagnosis of COVID-19, a combined oropharyngeal/nasal sampling method was used in the laboratory under standard polymerase chain reaction (RT-PCR). Data was analyzed with SPSS v. 19. The chi-square test was used to compare COVID-19 cases across substance types, abuse methods, and consumption frequencies.

Results: In total, 295 homeless drug abusers participated, and positive COVID-19 was recorded for 21(7.1%). Most participants were waste pickers (60.5%), and 3.9% had COVID-19. The statistical analysis showed no significant difference in the number of positive COVID-19 cases between narcotics, stimulants, and both combinations. Our study also did not prove the positive effect of drug abuse methods and the frequency of daily drug abuse on the incidence of this disease.

Conclusions: During the COVID-19 pandemic, unreliable, incorrect, and incomplete information on its prevention spread in societies. It is recommended that health-related rumors be investigated and officials publicly announce the results.

Keywords: COVID-19, Drug Abuse, Homelessness, Iran

1. Background

During the COVID-19 pandemic, there was a significant increase in rumors all around the world. The United Nations (UN) secretary general has identified these as a major challenge that needs to be addressed (1). In public health crises, individuals and communities may experience stress and turn to rumors to reduce it. Therefore, managing such situations is critical (2). The World Health Organization and other international health agencies have recognized rumors as a threat to pandemic preparedness and control that can decrease

community trust in governments. Consequently, they have suggested implementing systematic monitoring and control procedures to tackle this issue (3). On-time detection, assessment, and correction of rumors are challenging (1).

According to empirical research, repeating a rumor can make it more influential. Some people are more likely to believe in conspiracy theories and related rumors, even if they are logically contradictory (4).

Rumors can sometimes be mistaken for believable infection prevention and control strategies. The rumor of the significant effect of consuming highly

concentrated alcohol in disinfecting the body and eliminating the coronavirus resulted in the death of several sufferers (5). Furthermore, cases of blindness were reported after drinking methanol as a treatment for COVID-19 (6, 7).

Meanwhile, there were reports of an increase in the use of opioids after the COVID-19 pandemic. Drug overdose deaths increased in the United States in 2019, coinciding with the outbreak of COVID-19, primarily propelled by the widespread production of illicit fentanyl. It seems that the rise of COVID-19, along with disruptions in healthcare and social support, coupled with social and economic stress, worsened the drug epidemic (8). The rumors and inaccurate claims that opioid users are not at risk for COVID-19 led to a surge in demand for opioids compared to before the outbreak (9). Nevertheless, scientific evidence suggests that individuals who use drugs or smoke are more susceptible to COVID-19 (10, 11).

The COVID-19 pandemic involved a broad range of ages and social groups (12). In the meantime, homeless people were considered a high-risk group. Poverty, addiction, and mental health problems are some of the difficulties in these communities (13). Drug addiction in these individuals can be a risk factor for respiratory viral infections due to a lack of personal hygiene, lung infections, mucociliary dysfunction, weakened immunity, and disregard for social distancing (14, 15).

Like many countries, Iran is also facing the problems of homelessness and addiction (9). This country is considered a leading case in creating and expanding the model of integrating mental health and addiction into primary health care (PHC). Providing educational and health services to addicted homeless is defined in the health system (16).

Sistan and Baluchestan Province in southeastern Iran is still one of the main transit routes for drug trafficking from Afghanistan to European countries (17). In addition, the rate of addicted homeless per population is higher than in other parts of the country in this province (18). Therefore, it is a suitable area to study the accuracy of the rumor about the positive effect of opium and other addictive substances in preventing coronavirus infection.

2. Objectives

This study aimed to investigate the abundance of COVID-19 infection among homeless drug abusers in southeast Iran.

3. Patients and Methods

The data collection of this cross-sectional (descriptive-analytical) study was performed from April to May 2021, coinciding with the second-highest peak in Zahedan, the capital of Sistan and Baluchestan Province, southeastern Iran. It is the most populous city in the southeastern part of the country, with a population of 660 575. It is located near the shared border with Afghanistan and Pakistan (Figure 1).




Figure 1. Location of the study area: Zahedan, southeastern Iran.

All homeless individuals with drug addiction, gathered by Zahedan Municipality and scheduled for placement in the rehabilitation camp between April and May 2021, were included in the study. The participants' addiction was confirmed by their declaration and the laboratory examination. The research team obtained informed consent from all the participants. They were assured that the obtained information and their identities would be kept confidential.

For each participant, a questionnaire was completed by the health experts. It included demographic information, symptoms, type of addictive substances, and frequency and methods of drug abuse.

For a definite diagnosis of COVID-19, a combined oropharyngeal/nasal sampling method was used in the laboratory under standard polymerase chain reaction (RT-PCR).

The collected data were analyzed using SPSS v. 19 (IBM Corp., Armonk, NY, USA) software. Descriptive analyses were expressed as numbers and frequency percentages. The chi-square test was used to compare the number of positive COVID-19 cases in the types of substance consumed, drug abuse methods, and the number of times of consumption. The level of significance was established as $P < 0.05$.

4. Results

4.1. Demographic Characteristics

Table 1. Demographic Characteristics and Positive COVID-19 Status of Homeless Drug Abusers

Characteristic	Frequency (%)	COVID-19 ⁺	Characteristic	Frequency (%)	COVID-19 ⁺
Age (y)		Duration of homelessness			
≤ 25	20 (6.8)	0 (0)	≤ 5	37 (29.6)	4 (10.8)
26 - 35	91 (30.8)	3 (3.3)	5 - 10	25 (20.0)	0 (0)
36 - 45	114 (38.6)	11 (9.6)	> 10	63 (50.4)	5 (7.9)
≥ 46	65 (22)	7 (10.8)	Waste picker		
Sex		Yes			
Male	264 (89.5)	20 (7.6)	No		
Female	31 (10.5)	1 (3.2)	Education level		
Marital status		Illiterate			
Single	129 (44.2)	7 (5.4)	Elementary school		
Married	113 (38.7)	7 (6.2)	Junior high or high school		
Divorced	41 (14.0)	6 (14.6)	University graduate		
Widow	5 (1.7)	0 (0)	Night sleep		
Separated	1 (3)	0 (0)	Roofed place		
Others	3 (1.0)	0 (0)	Roofless place		
Night sleep		Alone			
Alone	112 (38.2)	9 (8.0)	With others		
With others	181 (61.8)	12 (6.6)			

In this study, 295 homeless drug abusers participated, most of whom were male (89.5%) and in the age group of 35 - 45 years. Totally, positive COVID-19 was recorded for 21 (7.1%) participants. People aged above 45 years (10.8%) had the highest incidence of the disease, and regarding sex, it was related to men (7.6%). Most participants were waste pickers (60.5%), and 3.9% of them had COVID-19. Although the positive cases were more seen in people who slept in roofed places and with others, the difference was not statistically significant ($P > 626$). The demographic and social characteristics of the participants are shown in [Table 1](#).

4.2. Possible Symptoms of COVID-19

Headache (33.3%), diarrhea (25%), stomachache (25%), fatigue (20%), and cough (12%) were the most common symptoms of COVID-19 positive cases in homeless drug abusers. The frequency of possible COVID-19 symptoms has been reported in [Table 2](#).

4.3. Types of Addictive Substances, Drug Abuse Methods, and Daily Use

The number of addicted participants who used narcotic analgesics, including opium and/or heroin, was 172 (59.5%), of which 10 (5.8%) people had positive COVID-19 results. Twenty patients (6.9%) used central nervous system (CNS) stimulants such as methamphetamine,

and 3 of them (15%) were positive for COVID-19. Among those who used narcotics and stimulants in combination, 7 (7.2%) were positive for the disease. The chi-square test showed that there was no significant difference in the number of positive COVID-19 cases among narcotics, stimulants, and both combinations. In other words, none of the drugs had an effect on the number of patients with COVID-19 ([Table 3](#)).

[Table 4](#) shows the frequency of drug abuse methods and the incidence of COVID-19 in homeless drug addicts. There was no significant difference in positive COVID-19 and drug abuse methods.

Additionally, the analysis did not show any significant difference in COVID-19 positivity and the number of times of drug abuse per day ([Table 5](#)).

5. Discussion

Access to addictive substances may be easier in a society with a high rate of addicts per population, so the rumor of the positive effect of addiction on the prevention or treatment of COVID-19 can encourage more people to abuse them. Therefore, an inspection of the accuracy of this opinion is necessary as a research point and can be effective in limiting or expanding drug addiction. Many studies have investigated COVID-19 in homeless people, but the literature review showed few studies on homeless drug addicts. Therefore, comparing the current study with similar ones is difficult.

Table 2. Frequency of the Possible Symptoms of COVID-19 in Homeless Drug Abusers

Symptom	Frequency (%)	COVID-19 ⁺	Symptom	Frequency (%)	COVID-19 ⁺
Fever			Nausea and vomiting		
Yes	12 (4.1)	1 (8.3)	Yes	5 (1.7)	1 (20.0)
No	283 (95.9)	20 (7.1)	No	290 (98.3)	20 (6.9)
Cough			Diarrhea		
Yes	8 (2.7)	1 (12.5)	Yes	4 (1.4)	1 (25.0)
No	287 (97.3)	20 (7.0)	No	291 (98.6)	20 (6.9)
Difficulty breathing			Stomachache		
Yes	41 (13.9)	3 (7.3)	Yes	4 (1.4)	1 (25.0)
No	254 (86.1)	18 (7.1)	No	291 (98.6)	20 (6.9)
Sore throat			Fatigue		
Yes	2 (.7)	0 (0)	Yes	5 (1.7)	1 (20.0)
No	293 (99.3)	21 (7.2)	No	290 (98.3)	20 (6.9)
Runny nose			Dizziness		
Yes	5 (1.7)	0 (0)	Yes	0 (0)	0 (0)
No	290 (98.3)	21 (7.2)	No	295 (100)	0 (0)
Chest pain			Body pain		
Yes	16 (5.4)	0 (0)	Yes	1 (.3)	0 (0)
No	279 (94.6)	21 (7.5)	No	294 (99.7)	0 (0)
Headache			Joint pain		
Yes	6 (2.0)	2 (33.3)	Yes	1 (.3)	0 (0)
No	289 (98.0)	19 (6.6)	No	294 (99.7)	0 (0)

Table 3. Frequency of Addictive Substance Types and COVID-19 Positivity in Homeless Drug Abusers

Category	No. (%)	COVID-19, No. (%)	P-Value
Narcotic analgesics (opium and/or heroin)	172 (58.3)	10 (5.8)	0.209
Central nervous system stimulants (methamphetamine)	26 (8.8)	4 (1.5)	
Both narcotics and stimulants	97 (32.9)	7 (7.2)	

Abbreviation: CNS, central nervous system.

It is expected that a high percentage of waste pickers get COVID-19 due to more exposure and contact with infectious rubbish. In addition, increased incidence and severity of airborne diseases have been recorded in homeless people. Also, the median age has been reported to be lower for the homeless than for non-homeless (19). In the present study, 7.1% of homeless drug abusers were infected with COVID-19. Most participants were in the age group of 35 - 45, and the highest incidence of COVID-19 was seen in the upper 45 years. Similarly, a study conducted on homeless drug addicts in Iran showed a 6.04% COVID-19 positivity rate (20). In another research in Ireland, the COVID-19 infection rate was 8.4% (63 out of 750 people) in homeless drug addicts (21). An investigation conducted in homeless shelters in the United States showed that the majority of participants were classified in the age

group of 35 - 59. The researchers showed the most positive COVID-19 infection in both 35 - 49 and above 60 years equally (22). In Brussels, researchers observed male predominance and a median age of about 56 years among homeless people hospitalized with COVID-19 (19).

Most participants slept in roofed places with other friends at night. In a study conducted in the United States, 86% of people with positive test results slept in public places. The researchers expressed that it can increase this group's infection probability (22). Overcrowding in shelters can enhance COVID-19 in the homeless population (19). Some studies have concluded that using prevention tools within homeless shelters can limit COVID-19 transmission (23).

In the current study, the most common symptoms of COVID-19 are similar to many other kinds of research. Other studies have also reported headaches, diarrhea,

Table 4. Frequency of Drug Abuse Methods and COVID-19 Positivity in Homeless Drug

Category	No. (%)	COVID-19, No. (%)	P-Value
Only oral	19 (6.4)	1 (0.3)	0.945
Both oral and inhalation	15 (5.1)	1 (0.3)	
All types ^a	261 (88.5)	19 (6.4)	

^a All types of drug use: Oral ingestion, inhalation, smoking, and intravenous injection.

Table 5. The Number of Drug Abuses Per Day and the Incidence of COVID-19 in Homeless Drug Abusers

Characteristic	No. (%)	COVID-19, No. (%)	P-Value
No. of drug abuses per day			0.405
1	37 (12.5)	1 (0.3)	
2	86 (29.2)	9 (3.1)	
3	113 (38.3)	8 (2.7)	
≥ 4	59 (20.0)	3 (1.0)	

stomachache, fatigue, and cough. However, the ranking was different compared to our study. A case series study showed fever, fatigue, myalgia, lack of appetite, dry cough, and anosmia as the most common COVID-19 symptoms respectively (24). Perhaps the reason for that is the different target group, in which they were not homeless addicts. Despite the broad spectrum of COVID-19, it seems that due to addictive substances' effects, some of the symptoms can be reduced or increased.

In our study, the statistical test did not show a significant difference in the number of COVID-19-positive cases between addictive substance types. However, the percentage of this disease was lower in people who used narcotics. Perhaps the last result has caused the rumor of the positive effect of traditional drug abuse on preventing COVID-19 in society. In the United States, increases in addictive substance use and drug overdoses have been documented since the COVID-19 pandemic (25-27). Some researchers have expressed that the COVID-19 pandemic or other stressful conditions can even cause small changes in the amount of drug abuse among addicts (28). Our study also did not prove the positive effect of drug abuse methods and the frequency of daily drug abuse on the incidence of this disease. The literature review did not confirm the positive role of addictive substances in the prevention and treatment of COVID-19. A COVID-19 serological study conducted in Iran on intravenous drug users showed no statistically significant between them and non-drug users (29).

5.1. Conclusions

During the COVID-19 pandemic, unreliable, incorrect, and incomplete information was spread in societies, and officials were forced to prevent and manage it (30). Rumors can be more dangerous than a pandemic, so they should be investigated and the results announced to the societies. Even with such a report, it is recommended that further laboratory and psychological studies be conducted with a focus on high-risk communities such as the homeless to determine the direction of future studies and understand the impact of rumors on society's behavior regarding healthcare.

Footnotes

Authors' Contribution: Conceptualization, literature review, and editing: Hassan Okati-Aliabad; investigation, data collection, writing - original draft: Jalil Nejati; formal analysis and data curation: Mahdi Mohammadi; technical and material support: Alireza Salimi Khorashad; methodology: Alireza Ansari-Moghaddam and Mohsen Hossein Bor.

Conflict of Interests: The authors declare no conflict of interest.

Data Availability: The dataset presented in the study is available on request from the corresponding author during submission or after publication.

Ethical Approval: This study was reviewed and approved by the Ethics Committee of Zahedan University of Medical Sciences (Ethical approval number: [IR.ZAUMS.REC.1400.162](https://doi.org/10.21860/IR.ZAUMS.REC.1400.162)).

Funding/Support: This study was supported by the Zahedan University of Medical Sciences.

References

- Islam MS, Sarkar T, Khan SH, Mostofa Kamal AH, Hasan SMM, Kabir A, et al. COVID-19-Related Infodemic and Its Impact on Public Health: A Global Social Media Analysis. *Am J Trop Med Hyg.* 2020;**103**(4):1621-9. [PubMed ID: 32783794]. [PubMed Central ID: PMC7543839]. <https://doi.org/10.4269/ajtmh.20-0812>.
- Person B, Sy F, Holton K, Govert B, Liang A, National Center for Infectious Diseases SCOT. Fear and stigma: the epidemic within the SARS outbreak. *Emerg Infect Dis.* 2004;**10**(2):358-63. [PubMed ID: 15030713]. [PubMed Central ID: PMC3322940]. <https://doi.org/10.3201/eid1002.030750>.
- World Health Organization. *Managing epidemics: key facts about major deadly diseases*. Geneva, Switzerland: World Health Organization; 2018. Available from: <https://www.who.int/publications/i/item/managing-epidemics-key-facts-about-major-deadly-diseases>.
- Berinsky AJ. Rumors and Health Care Reform: Experiments in Political Misinformation. *Br J Pol Sci.* 2015;**47**(2):241-62. <https://doi.org/10.1017/S0007123415000186>.
- World Health Organization. *Alcohol Does Not Protect against COVID-19; Access Should Be Restricted during Lockdown*. Geneva, Switzerland: World Health Organization; 2020. Available from: <https://www.who.int/europe/news/item/14-04-2020-alcohol-does-not-protect-against-covid-19-access-should-be-restricted-during-lockdown>.
- Aljazeera. *Iran: Over 700 dead after drinking alcohol to cure coronavirus*. Al Jazeera Media Network; 2020. Available from: <https://www.aljazeera.com/news/2020/4/27/iran-over-700-dead-after-drinking-alcohol-to-cure-coronavirus>.
- Hassanian-Moghaddam H, Zamani N, Kolahi AA, McDonald R, Hovda KE. Double trouble: methanol outbreak in the wake of the COVID-19 pandemic in Iran—a cross-sectional assessment. *Crit Care.* 2020;**24**(1):402. [PubMed ID: 32646475]. [PubMed Central ID: PMC7344025]. <https://doi.org/10.1186/s13054-020-03140-w>.
- Ning P, Cheng P, Li J, Zheng M, Schwebel DC, Yang Y, et al. COVID-19-Related Rumor Content, Transmission, and Clarification Strategies in China: Descriptive Study. *J Med Internet Res.* 2021;**23**(12): e27339. [PubMed ID: 34806992]. [PubMed Central ID: PMC8709421]. <https://doi.org/10.2196/27339>.
- Pirnia B, Dezhakam H, Pirnia K, Malekanmehr P, Soleimani AA, Zahiroddin A, et al. COVID-19 pandemic and addiction: Current problems in Iran. *Asian J Psychiatr.* 2020;**54**(In Press):102313. [PubMed ID: 32738562]. [PubMed Central ID: PMC7377711]. <https://doi.org/10.1016/j.ajp.2020.102313>.
- Arya S, Gupta R. COVID-19 outbreak: Challenges for Addiction services in India. *Asian J Psychiatr.* 2020;**51**(In Press):102086. [PubMed ID: 32315968]. [PubMed Central ID: PMC7194862]. <https://doi.org/10.1016/j.ajp.2020.102086>.
- Volkow ND. Collision of the COVID-19 and Addiction Epidemics. *Ann Intern Med.* 2020;**173**(1):61-2. [PubMed ID: 32240293]. [PubMed Central ID: PMC7138334]. <https://doi.org/10.7326/M20-1212>.
- Wang W, Tang J, Wei F. Updated understanding of the outbreak of 2019 novel coronavirus (2019-nCoV) in Wuhan, China. *J Med Virol.* 2020;**92**(4):441-7. [PubMed ID: 31994742]. [PubMed Central ID: PMC7167192]. <https://doi.org/10.1002/jmv.25689>.
- Lima NNR, de Souza RI, Feitosa PWG, Moreira JLS, da Silva CGL, Neto MLR. People experiencing homelessness: Their potential exposure to COVID-19. *Psychiatry Res.* 2020;**288**(In Press):112945. [PubMed ID: 32302813]. [PubMed Central ID: PMC7151321]. <https://doi.org/10.1016/j.psychres.2020.112945>.
- Saeedi M, Omrani-Nava V, Maleki I, Hedayatizadeh-Omran A, Ahmadi A, Moosazadeh M, et al. Opium Addiction and COVID-19: Truth or False Beliefs. *Iranian Journal of Psychiatry and Behavioral Sciences.* 2020;**14**(2). <https://doi.org/10.5812/ijpbs.103509>.
- Mahdavi A, Aliramezany M. Addiction and Covid-19 Disease: Risks and Misconceptions. *Addict Health.* 2021;**13**(1):66-7. [PubMed ID: 33995961]. [PubMed Central ID: PMC8080171]. <https://doi.org/10.22122/ahj.v13i1.279>.
- Dadfar M, Bahrami F. Descriptive Reports of Integration of Mental Health into the Primary Health Care (PHC) System in One of the Areas of Iran. *American Journal of Educational Research.* 2015;**3**(11):1429-32. [PubMed ID: doi:10.12691/education-3-11-12].
- Sabatelle DR. The scourge of opiates: The illicit narcotics trade in the Islamic Republic of Iran. *Trends Organ Crime.* 2011;**14**(4):314-31. <https://doi.org/10.1007/s12117-011-9142-0>.
- Rajabalipour M, Khoshab H, Baneshi MR, Nakhaee N, Sharifi H, Tavakoli F, et al. Using Social Cognitive Theory to Investigate the Risk Factors of Waterpipe Smoking among Southeastern Iranian Adolescents. *Int J Pediatr.* 2019;**7**(10):10243-53. <https://doi.org/10.22038/ijp.2019.40663.3429>.
- Schrooyen L, Delforge M, Lebout F, Vanbaelen T, Lecompte A, Dauby N. Homeless people hospitalized with COVID-19 in Brussels. *Clin Microbiol Infect.* 2021;**27**(1):151-2. [PubMed ID: 32777362]. [PubMed Central ID: PMC7411419]. <https://doi.org/10.1016/j.cmi.2020.08.002>.
- Shakib A, Mousavi MS, Panahi S, Asnaashari M, Saadabadi A, Arshadi M, et al. [COVID-19 Screening in Homeless Drug Addicts by the Deputy of Health of Tehran University of Medical Sciences]. *J Mar Med.* 2021;**3**(4):28-33. Persian. <https://doi.org/10.30491/3.4.28>.
- Collins J, O'Carroll A, Duffin T. *A milestone in drug policy: Saving the lives of people who use drugs and were homeless in Dublin during the covid-19 pandemic*. LSE Research Online; 2020. Available from: http://eprints.lse.ac.uk/105941/1/covid19_2020_07_15_a_milestone_in_drug_policy_saving_the_lives_of_pdf.
- Rogers JH, Link AC, McCulloch D, Brandstetter E, Newman KL, Jackson ML, et al. Characteristics of COVID-19 in Homeless Shelters : A Community-Based Surveillance Study. *Ann Intern Med.* 2021;**174**(1):42-9. [PubMed ID: 32931328]. [PubMed Central ID: PMC7517131]. <https://doi.org/10.7326/M20-3799>.
- Karb R, Samuels E, Vanjani R, Trimbur C, Napoli A. Homeless Shelter Characteristics and Prevalence of SARS-CoV-2. *West J Emerg Med.* 2020;**21**(5):1048-53. [PubMed ID: 32970553]. [PubMed Central ID: PMC7514394]. <https://doi.org/10.5811/westjem.2020.7.48725>.
- Macera M, De Angelis G, Sagnelli C, Coppola N, Vanvitelli C. Clinical Presentation of COVID-19: Case Series and Review of the Literature. *Int J Environ Res Public Health.* 2020;**17**(14). [PubMed ID: 32674450]. [PubMed Central ID: PMC7399865]. <https://doi.org/10.3390/ijerph17145062>.
- Friedman J, Akre S. COVID-19 and the Drug Overdose Crisis: Uncovering the Deadliest Months in the United States, January–July 2020. *Am J Public Health.* 2021;**111**(7):1284-91. [PubMed ID: 33856885]. [PubMed Central ID: PMC8493145]. <https://doi.org/10.2105/AJPH.2021.306256>.
- Niles JK, Gudín J, Radcliff J, Kaufman HW. The Opioid Epidemic Within the COVID-19 Pandemic: Drug Testing in 2020. *Popul Health Manag.* 2021;**24**(S1):S43-51. [PubMed ID: 33031013]. [PubMed Central ID: PMC7875135]. <https://doi.org/10.1089/pop.2020.0230>.
- Wainwright JJ, Mikre M, Whitley P, Dawson E, Huskey A, Lukowiak A, et al. Analysis of Drug Test Results Before and After the US Declaration of a National Emergency Concerning the COVID-19 Outbreak. *JAMA.* 2020;**324**(16):1674-7. [PubMed ID: 32945855].

- [PubMed Central ID: [PMC7501585](https://pubmed.ncbi.nlm.nih.gov/PMC7501585/)], <https://doi.org/10.1001/jama.2020.17694>.
28. Chiappini S, Mosca A, Alessi MC, Picutti E, di Giannantonio M. How the COVID-19 Pandemic Changed the World of Addiction: Considerations on the Impact on Substance Use and Treatment. *J Ital Soc Psychiatry*. 2021;7(In Press):95-9.
29. Mohammadi Z, Eghtesad S, Hashemi-Shahri SM, Tabatabaei SM, Sharafkhah M, Poustchi H. The Seroprevalence of COVID-19 in Intravenous Drug Users in Comparison to Non-drug Users. *Middle East J Dig Dis*. 2021;13(1):67-70. [PubMed ID: [34712441](https://pubmed.ncbi.nlm.nih.gov/34712441/)]. [PubMed Central ID: [PMC8531943](https://pubmed.ncbi.nlm.nih.gov/PMC8531943/)]. <https://doi.org/10.34172/mejdd.2021.206>.
30. Mohajervatan A, Rezaei F. Are rumors more dangerous than covid-19 pandemic? *Journal of Drug and Alcohol Research*. 2021;10(8).