



Mild Symptoms of SARS-CoV-2 Infection in a 3-Days-Old Newborn with Congenital Heart Disease: A Case Report

Azin Etemadimanesh ¹, Ali Ghasemi ¹, Mahmoud Khodabandeh ² and Hamid Eshaghi ^{2,*}

¹School of Medicine, Tehran University of Medical Sciences, Tehran, Iran

²Department of Infectious Diseases, Pediatric's Center of Excellence, Children's Medical Center, Tehran University of Medical Sciences, Tehran, Iran

*Corresponding author: Department of Infectious Diseases, Pediatric's Center of Excellence, Children's Medical Center, Tehran University of Medical Sciences, Tehran, Iran.
Email: h-eshaghi@sina.tums.ac.ir

Received 2020 August 02; Revised 2021 February 07; Accepted 2021 March 01.

Abstract

Background: SARS-CoV-2 as a member of Coronavirus family, caused a global pandemic in late 2019 and raised concerns about its morbidity and mortality among immune-deficient individuals. Till now, several pediatric cases infected with SARS-CoV-2 have been reported, and some have noted susceptibility to infection in infants and young children, especially those with congenital comorbidities. In this report, we discuss the clinical course, administered treatments, and outcomes of SARS-COV-2 infection in an infant suffering from cardiovascular comorbidity.

Case Presentation: We describe a newborn referred to hospital 72 hours postpartum, diagnosed with pulmonary atresia with ventricular septal defect (PA-VSD), and subsequently found to be infected with the SARS-COV-2 virus. The patient presented with tachypnea, lethargy, and a history of recent fever and myalgia in his father. He received intravenous fluid and antibiotic therapy based on an established protocol for COVID-19 treatment by Iran health ministry and was discharged after five days of hospitalization without further complications. Two weeks after discharge, he was referred to the cardiac surgery department for surgical treatment after obtaining a negative result for nasopharyngeal sample SARS-CoV-2 RT-PCR.

Conclusions: Mild symptoms and no need for excessive oxygen supports in the current case demonstrate pediatrics patients with COVID-19 have a better prognosis and fewer complications compared with adults, even at early childhood and with the presence of serious cardiac complications.

Keywords: COVID-19, Heart Septal Defects, Pulmonary Atresia, Infant, Newborn

1. Background

In late 2019, a respiratory infection caused by novel-coronavirus 2019 (COVID-19) emerged in China and led to a global pandemic. Afterward, several pediatric cases have been reported, and some have noted susceptibility to infection in young children, especially infants (1). Common symptoms of SARS-COV-2 infection include dry cough, respiratory distress, fever, myalgia, and gastrointestinal manifestations such as diarrhea. Reported pediatric cases mostly presented with mild symptoms and required intensive care and respiratory support less frequently compared with adults (2). However, more studies are mandatory for a decisive understanding of the clinical course and outcomes of SARS-COV-2 infection in pediatric patients, particularly newborns.

Here, we report a case of SARS-COV-2 infection in a 3-days-old infant with Pulmonary Atresia with Ventricular

Septal Defect (PA-VSD) who was presented to a tertiary referral hospital with tachypnea.

2. Case Presentation

The patient was a boy, born at 37 weeks of gestation. He presented with tachypnea 72 hours after birth and was hospitalized for further investigations. Physical examination on admission revealed a lethargic newborn. He was tachypneic (respiratory rate of 46 breaths/minute), but the rest of the vital signs were within the normal range. The patient's oxygen saturation -measured with pulse oximetry- was 98%. He underwent cardiac echocardiography, which showed evidence of PA-VSD.

Pediatric cardiovascular surgery department scheduled the patient for heart surgery. However, during pre-operative preparations, the patient was diagnosed with COVID-19. Surgery was canceled due to the positive SARS-

COV-2 RT-PCR test, and the patient was admitted to the infectious diseases ward. Supportive care and antibiotics (based on an approved protocol of COVID-19 management published by Iran health ministry at the time) were initiated after obtaining blood and urine samples for bacterial cultures.

The patient's mother had no symptoms of any infection before and during labor. However, his father mentioned a history of fever and lethargy in the last few days without a definitive diagnosis of SARS-COV-2 infection.

The respiratory examination was unremarkable, and the chest X-ray radiography showed no definitive signs of pneumonia (Figure 1). Routine laboratory tests were obtained, and the results are listed in Table 1. Blood and urine culture results were negative. Venous blood gas (VBG) analysis demonstrated a pH of 7.36 and oxygen partial pressure (PO₂) of 46.8 mmHg, and carbon dioxide (PCO₂) of 26.3 mmHg.

Table 1. Routine Laboratory Parameters

Laboratory Parameters	Result
White blood cells ($\times 10^3/\mu\text{L}$)	9.63
Neutrophils (%)	57.3
Lymphocytes (%)	30
Hemoglobin (g/dL)	17.3
Platelets ($\times 10^3/\mu\text{L}$)	301
Urine analysis	Negative
Urine culture	Negative
Blood culture	Negative
C-reactive protein (mg/L)	1

His clinical status was monitored closely every day, and there was no record of fever, cough, gastrointestinal symptoms, or hemodynamic instability in the patient during hospitalization. He merely developed mild respiratory distress and tachypnea that required no additional respiratory support and was supposed to be related to the aforementioned congenital heart disease.

The patient was discharged on the 5th day of admission without complications. He did not show any signs of pneumonia during hospitalization and was in no need for excessive ventilation or oxygen support. Two weeks after discharge, a nasopharyngeal swab test for SARS-COV-2 RT-PCR was repeated and reported negative, and he was referred to a cardiovascular surgeon to reschedule his surgery.

3. Discussion

Here, we described a confirmed SARS-COV-2 infection in a 3-day-old full-term male with congenital heart disease (PA-VSD) who was hospitalized due to tachypnea.

The possibility of vertical transmission of SARS-CoV-2 has been mentioned previously in some newborn case reports (3). However, the absence of symptoms or definite SARS-COV-2 diagnosis in mother, and a history of recent fever and myalgia in father raises the possibility that the disease was transmitted from father to the infant through close contacts after birth.

In a recent report, less than 1% of all SARS-COV-2 cases were younger than ten years old (2), about 16% of them did not have any respiratory symptoms or radiologic changes suggestive of pneumonia, and most cases, which required intensive care and respiratory support had preexisting comorbidities (4). Some case reports of neonatal SARS-COV-2 infection demonstrated that it could potentially present as respiratory failure in neonates (5). However, progression to pneumonia requires oxygen supply, and intensive care support that occurs in most severe cases of adults was not seen in this patient despite the underlying cardiovascular disease.

3.1. Conclusions

More data is needed to reach a comprehensive understanding of clinical features and outcomes of SARS-COV-2 infection among younger children. In this report, we described a 3-day-old newborn with congenital heart disease and SARS-CoV-2 infection without any serious complications of COVID-19. Based on available reports, contrary to adults, pediatric cases mostly present with a spectrum of asymptomatic to mild symptoms, even for patients with critical comorbidities, namely congenital heart diseases.

Footnotes

Authors' Contribution: Study concept and design: H. E., and M. K.; Drafting of the manuscript: A. E.; Critical revision of the manuscript for important intellectual content: A.G.

Conflict of Interests: The authors had no conflict of interest in this study.

Funding/Support: The authors received no funding for this study.

Informed Consent: Informed consent was obtained from the patient's parents.




Figure 1. Chest X-ray graph of the patient. It has no radiological findings suggestive of pneumonia.

References

1. Dong Y, Mo X, Hu Y, Qi X, Jiang F, Jiang Z, et al. Epidemiology of COVID-19 among children in China. *Pediatrics*. 2020;**145**(6). doi: [10.1542/peds.2020-0702](https://doi.org/10.1542/peds.2020-0702). [PubMed: [32179660](https://pubmed.ncbi.nlm.nih.gov/32179660/)].
2. Wu Z, McGoogan JM. Characteristics of and important lessons from the coronavirus disease 2019 (COVID-19) outbreak in China: Summary of a report of 72314 cases from the Chinese center for disease control and prevention. *JAMA*. 2020;**323**(13):1239-42. doi: [10.1001/jama.2020.2648](https://doi.org/10.1001/jama.2020.2648). [PubMed: [32091533](https://pubmed.ncbi.nlm.nih.gov/32091533/)].
3. Wang S, Guo L, Chen L, Liu W, Cao Y, Zhang J, et al. A Case report of neonatal 2019 coronavirus disease in China. *Clin Infect Dis*. 2020;**71**(15):853-7. doi: [10.1093/cid/ciaa225](https://doi.org/10.1093/cid/ciaa225). [PubMed: [32161941](https://pubmed.ncbi.nlm.nih.gov/32161941/)]. [PubMed Central: [PMC7108144](https://pubmed.ncbi.nlm.nih.gov/PMC7108144/)].
4. Lu X, Zhang L, Du H, Zhang J, Li YY, Qu J, et al. SARS-CoV-2 infection in children. *N Engl J Med*. 2020;**382**(17):1663-5. doi: [10.1056/NEJMc2005073](https://doi.org/10.1056/NEJMc2005073). [PubMed: [32187458](https://pubmed.ncbi.nlm.nih.gov/32187458/)]. [PubMed Central: [PMC7121177](https://pubmed.ncbi.nlm.nih.gov/PMC7121177/)].
5. Precit MR, Yee R, Anand V, Mongkolrattanothai K, Pandey U, Dien Bard J. A case report of neonatal acute respiratory failure due to severe acute respiratory syndrome coronavirus-2. *J Pediatric Infect Dis Soc*. 2020;**9**(3):390-2. doi: [10.1093/jpids/piaa064](https://doi.org/10.1093/jpids/piaa064). [PubMed: [32441743](https://pubmed.ncbi.nlm.nih.gov/32441743/)]. [PubMed Central: [PMC7313921](https://pubmed.ncbi.nlm.nih.gov/PMC7313921/)].