



Computerized Tomography Scan Findings in Pediatric Patients with Coronavirus Disease 2019

Hamid Dahmardeh ¹, Gholamreza Soleimani ², Alireza Ahmadi¹ and Alireza Teimouri ^{2,*}

¹Department of Radiology, School of Medicine, Zahedan University of Medical Sciences, Zahedan, Iran

²Children and Adolescents Health Research Center, Research Institute of Cellular and Molecular Science in Infectious Diseases, Zahedan University of Medical Science's, Zahedan, Iran

*Corresponding author: Children and Adolescents Health Research Center Office, Ali Ibne Abitalib Hospital, Zahedan University of Medical Science's, Zahedan, Iran. Email: alirezateimouri260@gmail.com

Received 2023 June 13; Revised 2023 August 10; Accepted 2023 August 31.

Abstract

Background: This study aimed to ascertain the computerized tomography findings of 56 children diagnosed with coronavirus disease 2019 (COVID-19) at Ali Ibn Abitalib Hospital in Zahedan, Iran.

Methods: This descriptive, cross-sectional study was conducted on 56 pediatric patients with laboratory-confirmed COVID-19 who were treated at Ali Ibn Abitalib Hospital in Zahedan from March to August 2020. The computerized tomography (CT) scans were performed for these children because they did not show signs of improvement, had decreased oxygen levels during the treatment, had severe respiratory symptoms, or had a breathing disorder. The CT findings were reviewed to determine the frequency of normal and abnormal scans. The distribution of lesions (unilateral, bilateral, peripheral, central), type of opacity (consolidation, ground glass, etc.), and presence or absence of bronchiectasis, effusion, cavity, and lymphadenopathy were recorded. The frequency of each type of lesion was then calculated.

Results: Of the patients, 30.36% had a normal CT scan. The most common CT scan finding was ground-glass opacity (42.86%), followed by consolidation (35.71%). The distribution of lesions was mostly bilateral (64.10%) and mainly peripheral (66.67%).

Conclusions: One-third of children with COVID-19 had normal CT features. Ground-glass opacity and consolidation were more common than cavity or pneumatocele, pleural effusion, bronchiectasis, and lymphadenopathy.

Keywords: COVID-19, Pediatrics, Chest Computed Tomography

1. Background

In December 2019, an outbreak of pneumonia caused by coronavirus disease 2019 (COVID-19) emerged in Wuhan, the capital of Hubei province, China (1). Since then, the infection has spread globally, with significant numbers occurring in Asia, Europe, and other regions of the world (2, 3). Initially, those infected outside China were reported in individuals who had traveled to China or had been in contact with tourists from China. However, there are now increasing reports of community outbreaks, including within the United States of America (4). As of April 2020, a total of 2,878,196 confirmed cases of COVID-19 have been reported in over 100 countries worldwide, resulting in 198,668 deaths (4). The average age of affected individuals was 47 years (5), with a disproportionate incidence in older men with co-morbidities (6). Approximately 2% of reported cases were pediatric cases (7, 8), and 0.9% of all

affected patients were under the age of 15 (5). Although early investigations described the medical and imaging features of COVID-19 in adults (9-12), computerized tomography (CT) findings in children have only been reported in small cohorts (13-15). Rapid diagnosis is challenging in children due to milder symptoms and lower hospitalization rates than adult patients (8, 13-17). Rapid and accurate diagnosis is particularly important, as severe symptoms and deaths have been reported in pediatric patients (8). Additionally, the social and psychological impact of isolating children, as well as the economic impact on caregivers, must be taken into consideration.

2. Objectives

This study aimed to identify CT features in children with COVID-19.

3. Methods

This descriptive cross-sectional study was conducted on inpatient children aged 1 month to 18 years hospitalized with a diagnosis of COVID-19 at Ali Ibn Abitalib Hospital, a comprehensive medical center that integrates medical care, education, and scientific research affiliated with Zahedan University of Medical Sciences (ZaUMS). The study was done from March 1, 2020, to August 30, 2020. During this period, 56 pediatric patients staying in the hospital needed a CT scan. This was because their RT-PCR test was negative or they showed no improvement. They also had low oxygen levels, severe respiratory symptoms, or trouble breathing.

For these children, COVID-19 was confirmed by a positive RT-PCR result. Confirmed or suspected children were referred to the radiology department for CT scans. Two experienced radiologists looked at the CT scans separately. They did not know anything about the patient's medical history. A special CT scanner was used to take images inside the body. The machine used was made by Siemens and is called the SOMATOM multi-slice spiral scanner. They did not inject any medicine into the body using a needle for this scan. For the imaging machine, the tube voltage was between 100 and 120 kV, there was automatic exposure control, the collimation was set to 2.0 mm, and the pitch was set to 1. We looked at CT scans from different angles to see both the lungs and the area in the middle of the chest. We found out how often normal and abnormal CT scans occurred.

A form was used to collect information about where and how the lesions were distributed in the body (one side or both sides, on the edges or in the center) and what kind of blockage was seen (hardening, hazy appearance, etc.). We also noted whether bronchiectasis, exudate, cavity, or lymphadenopathy was present. Then, we worked out how often each type of problem occurred. This study was approved by the university's ethics committee with a code of IR.ZAUMS.REC.1399.532. Data were aggregated and reported using SPSS 20.0 (SPSS Inc., Chicago, IL, USA). Mean and SD were used for quantitative variables, and frequency and percentage were used for categorical variables.

4. Results

This study obtained CT scans from 56 pediatric patients infected or suspected of COVID-19 who met the inclusion criteria. The patients' ages ranged from 1 month to 16 years, with a median age of 6 years and a male-to-female ratio of 1.54:1. The demographic features of patients and age group statistics are presented in [Table 1](#). [Table 2](#) displays the chest CT features of COVID-19 pediatric patients, indicating that

17 patients (30.36%) had normal CT findings. Ground-glass opacity (42.86%) and consolidation (35.71%) were the most prevalent CT findings in these patients. The distribution of lesions was mostly bilateral (64.10%) and mainly peripheral (66.67%). Other findings included cavity or pneumatocele (5.36%), pleural effusion (3.57%), bronchiectasis (1.79%), and lymphadenopathy (1.79%).

Table 1. Demographic Features of Coronavirus Disease 2019 Confirmed Patients (N = 56)

| Variables | Mean ± SD or No. (%) |
|----------------|---------------------------------|
| Age (y) | 6 ± 4.15 (3 months to 16 years) |
| < 1 | 6 (10.71) |
| 1-5 | 16 (28.57) |
| 5-10 | 23 (41.07) |
| > 10 | 11 (19.64) |
| Gender | |
| Boy | 34 (60.71) |
| Girl | 22 (39.29) |

5. Discussion

This study found that 30.36% of the 56 patients had normal CT findings, while those with positive CT findings exhibited typical features such as ground-glass opacities and airspace disease, which presented as a thrombus with one or more lobes.

Notably, ground-glass opacities were more prevalent in this study, and a few patients developed pleural effusion, lymphadenopathy, and bronchiectasis. Approximately 44.64% of patients had bilateral pulmonary involvement, and 46.43% had peripheral pulmonary findings. These imaging findings suggest that COVID-19 may primarily manifest as ground-glass opacities in bilateral and peripheral lobe distributions in the pediatric population. This study is one of the largest and most comprehensive in this geographical area to demonstrate imaging findings in pediatric patients with COVID-19. The imaging results and their distribution are consistent with those published in the literature on COVID-19 in both children and adults ([9-14, 18](#)).

In this study, the detection rate of abnormal CT findings in children was only 69.64% compared to adults, which is similar to the findings of Shelmerdine et al. (64%) and Wang et al.'s study (63.2%) ([19, 20](#)). Ground-glass opacities (42.86%) were the most common CT findings in children with COVID-19 in this study, followed by consolidation (35.71%). The prevalence of these two features was slightly lower in children

Table 2. Chest Computerized Tomography Findings of Enrolled Patients

| Variables | No. (%) |
|--------------------------|------------|
| Normal | 17 (30.36) |
| Abnormal | 39 (69.64) |
| Opacities | |
| Ground-glass opacity | 24 (42.86) |
| Consolidation | 20 (35.71) |
| Nodular | 12 (21.43) |
| Reticular | 6 (10.71) |
| Crazy paving | 2 (3.57) |
| Laterality | |
| Normal | 17 (30.36) |
| Unilateral | 14 (25) |
| Bilateral | 25 (44.64) |
| Distribution | |
| Normal | 17 (30.36) |
| Peripheral | 26 (46.43) |
| Central | 11 (19.64) |
| Diffuse | 2 (3.57) |
| Other CT findings | |
| Cavity/pneumatocele | 3 (5.36) |
| Pleural effusion | 2 (3.57) |
| Bronchiectasis | 1 (1.79) |
| Lymphadenopathy | 1 (1.79) |

Abbreviation: CT, computerized tomography.

than in adult patients. Ojha et al. (21) reported 50.2% and 39% for ground-glass opacity and consolidation, respectively, while Wan et al. (22) reported 86% and 47% for ground-glass opacity and consolidation, respectively. Ebrahimpour and colleagues (23) examined the frequency of CT findings in pediatrics and found that the incidence of ground-glass opacity was 57.4% and consolidation was 10.2%, comparable to the present study's results. Additionally, the incidence of ground-glass opacity in Kumar et al. (24) and Wang et al. (20) studies was 40% and 39.5%, respectively, similar to our study. The rate in this study was 64.10%, compared to 61.5% in the study by Wang et al. (20) and 53.62% in the Ebrahimpour et al. study (23).

Ghods et al. (25) conducted the same study on children under the age of one and found a specific rate of 57.63%, and pulmonary involvement was predominantly peripheral (66.67%). In the Kumar et al. (24) study, 80% of the opacities were peripheral, and in the Ebrahimpour and colleagues (23) study, 81.2%, 15.6%, and 3.1% of children had a peripheral, diffuse, and central pattern, respectively. Other

common findings in this study were pulmonary nodules and reticular opacities (21.43% and 10.71%, respectively). In Ebrahimpour et al.'s study, pulmonary nodules were observed in 2.6% of patients, and reticulation was observed in 1.7%, less than in the present study. The nodular and reticular opacity frequencies in Wang et al.'s study were 25.7% and 9.7%, respectively (20, 23). Pleural effusion (3.57%) and lymphadenopathy (1.79%) were rare findings in the current study. Other studies reported pleural effusion rates ranging from 2.7% to 9.1% and mediastinal lymphadenopathy at 0.5% (20-23). The 'crazy paving' pattern, one of the hallmarks of severe lung disease, was found less frequently than in adult patients. In our study, it occurred in only 3.57% of the children, whereas in adults, it has been reported as 12%, 15%, and 19.5% (21, 22, 26). In Ebrahimpour et al.'s study, the frequency of crazy patch patterns in children was only 0.6% (23). Because the CT manifestations of pneumonia caused by different pathogens can overlap, and COVID-19 pneumonia can also overlap with pneumonia caused by other pathogens, it may present with more severe and complex imaging manifestations. Therefore, combining epidemiological, etiological, and radiological investigations is important. The small sample size was the limitation of the study.

5.1. Conclusions

In conclusion, the study determined that approximately one-third of the children with COVID-19 exhibited normal CT features. Ground-glass opacity and consolidation were more prevalent than cavity or pneumatocele, pleural effusion, bronchiectasis, and lymphadenopathy. Most COVID-19-positive children exhibit only mild CT findings, and given the potential risk of radiation exposure, CT imaging results may not significantly impact treatment decisions. Therefore, CT scans may not be necessary for most patients and may be replaced by CXR, except for individual cases.

Acknowledgments

The authors would like to acknowledge the children's parents.

Footnotes

Authors' Contribution: H. D. and G.H. S.: Main concepts, design, and supervision; A. A.: Data acquisition; A. T.: Data analysis, literature search, interpretation, drafting of the primary version of the manuscript, and final approval.

Conflict of Interests: The authors declare no conflict of interest. All working at Zahedan University of Medical

Sciences have no support from any organizations for their research and projects, no financial interests, and no consultation fees. They are not editorial board members, but Gholamreza Soleimani has been an active reviewer of this journal within the last five years.

Data Reproducibility: The dataset presented in the study is available on request from the corresponding author during submission or after publication. The data are not publicly available due to being part of a big database related to pediatric patients with COVID-19.

Ethical Approval: This study was approved by the university's ethics committee with a code of IR.ZAUMS.REC.1399.532.

Funding/Support: The study was not supported by any organization or agencies.

Informed Consent: It was not declared by the authors.

References

- Chen N, Zhou M, Dong X, Qu J, Gong F, Han Y, et al. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet*. 2020;395(10223):507-13. [PubMed ID: 32007143]. [PubMed Central ID: PMC7135076]. [https://doi.org/10.1016/S0140-6736\(20\)30211-7](https://doi.org/10.1016/S0140-6736(20)30211-7).
- European Centre for Disease Prevention and Control. *Geographical distribution of COVID-19 cases*. 2020, [cited 11 March 2020]. Available from: www.ecdc.europa.eu/en/geographical-distribution-2019-ncov-cases.
- Centers for Disease Control and Prevention. *Coronavirus disease (2019) (COVID-19): cases in the US*. 2020, [cited 18 March 2020]. Available from: www.cdc.gov/coronavirus/2019-ncov/cases-in-us.html.
- World Health Organization. *Coronavirus disease 2019 (COVID-19): Situation Report - 98*. 2020. Available from: <https://www.who.int/docs/default-source/coronavirus/situation-reports/20200427-sitrep-98-covid-19.pdf>.
- Guan WJ, Ni ZY, Hu Y, Liang WH, Ou CQ, He JX, et al. Clinical Characteristics of Coronavirus Disease 2019 in China. *N Engl J Med*. 2020;382(18):1708-20. [PubMed ID: 32109013]. [PubMed Central ID: PMC7092819]. <https://doi.org/10.1056/NEJMoa2002032>.
- Del Rio C, Malani PN. 2019 Novel Coronavirus-Important Information for Clinicians. *JAMA*. 2020;323(11):1039-40. [PubMed ID: 32022836]. <https://doi.org/10.1001/jama.2020.1490>.
- Wu Z, McGoogan JM. Characteristics of and Important Lessons From the Coronavirus Disease 2019 (COVID-19) Outbreak in China: Summary of a Report of 72314 Cases From the Chinese Center for Disease Control and Prevention. *JAMA*. 2020;323(13):1239-42. [PubMed ID: 32091533]. <https://doi.org/10.1001/jama.2020.2648>.
- CDC COVID-19 Response Team. Coronavirus Disease 2019 in Children - United States, February 12-April 2, 2020. *MMWR Morb Mortal Wkly Rep*. 2020;69(14):422-6. [PubMed ID: 32271728]. [PubMed Central ID: PMC7147903]. <https://doi.org/10.15585/mmwr.mm6914e4>.
- Chung M, Bernheim A, Mei X, Zhang N, Huang M, Zeng X, et al. CT Imaging Features of 2019 Novel Coronavirus (2019-nCoV). *Radiology*. 2020;295(1):202-7. [PubMed ID: 32017661]. [PubMed Central ID: PMC7194022]. <https://doi.org/10.1148/radiol.2020200230>.
- Zhao W, Zhong Z, Xie X, Yu Q, Liu J. Relation Between Chest CT Findings and Clinical Conditions of Coronavirus Disease (COVID-19) Pneumonia: A Multicenter Study. *AJR Am J Roentgenol*. 2020;214(5):1072-7. [PubMed ID: 32125873]. <https://doi.org/10.2214/AJR.20.22976>.
- Bernheim A, Mei X, Huang M, Yang Y, Fayad ZA, Zhang N, et al. Chest CT Findings in Coronavirus Disease-19 (COVID-19): Relationship to Duration of Infection. *Radiology*. 2020;295(3):200463. [PubMed ID: 32077789]. [PubMed Central ID: PMC7233369]. <https://doi.org/10.1148/radiol.2020200463>.
- Salehi S, Abedi A, Balakrishnan S, Gholamrezaezhad A. Coronavirus Disease 2019 (COVID-19): A Systematic Review of Imaging Findings in 919 Patients. *AJR Am J Roentgenol*. 2020;215(1):87-93. [PubMed ID: 32174129]. <https://doi.org/10.2214/AJR.20.23034>.
- Chen A, Huang JX, Liao Y, Liu Z, Chen D, Yang C, et al. Differences in Clinical and Imaging Presentation of Pediatric Patients with COVID-19 in Comparison with Adults. *Radiol Cardiothorac Imaging*. 2020;2(2):e200117. [PubMed ID: 33778567]. [PubMed Central ID: PMC7233434]. <https://doi.org/10.1148/ryct.2020200117>.
- Xia W, Shao J, Guo Y, Peng X, Li Z, Hu D. Clinical and CT features in pediatric patients with COVID-19 infection: Different points from adults. *Pediatr Pulmonol*. 2020;55(5):1169-74. [PubMed ID: 32134205]. [PubMed Central ID: PMC7168071]. <https://doi.org/10.1002/ppul.24718>.
- Jiehao C, Jin X, Daojiong L, Zhi Y, Lei X, Zhenghai Q, et al. A Case Series of Children With 2019 Novel Coronavirus Infection: Clinical and Epidemiological Features. *Clin Infect Dis*. 2020;71(6):1547-51. [PubMed ID: 32112072]. [PubMed Central ID: PMC7108143]. <https://doi.org/10.1093/cid/ciaa198>.
- Shen KL, Yang YH. Diagnosis and treatment of 2019 novel coronavirus infection in children: a pressing issue. *World J Pediatr*. 2020;16(3):219-21. [PubMed ID: 32026147]. [PubMed Central ID: PMC7091265]. <https://doi.org/10.1007/s12519-020-00344-6>.
- Dong Y, Mo X, Hu Y, Qi X, Jiang F, Jiang Z, et al. Epidemiology of COVID-19 Among Children in China. *Pediatrics*. 2020;145(6):e20200702. [PubMed ID: 32179660]. <https://doi.org/10.1542/peds.2020-0702>.
- Simpson S, Kay FU, Abbara S, Bhalla S, Chung JH, Chung M, et al. Radiological Society of North America Expert Consensus Document on Reporting Chest CT Findings Related to COVID-19: Endorsed by the Society of Thoracic Radiology, the American College of Radiology, and RSNA. *Radiol Cardiothorac Imaging*. 2020;2(2):e200152. [PubMed ID: 33778571]. [PubMed Central ID: PMC7233447]. <https://doi.org/10.1148/ryct.2020200152>.
- Shelmerdine SC, Lovrenski J, Caro-Dominguez P, Toso S; Collaborators of the European Society of Paediatric Radiology Cardiothoracic Imaging Taskforce. Coronavirus disease 2019 (COVID-19) in children: a systematic review of imaging findings. *Pediatr Radiol*. 2020;50(9):1217-30. [PubMed ID: 32556807]. [PubMed Central ID: PMC7300372]. <https://doi.org/10.1007/s00247-020-04726-w>.
- Wang JG, Mo YF, Su YH, Wang LC, Liu GB, Li M, et al. Computed tomography features of COVID-19 in children: A systematic review and meta-analysis. *Medicine (Baltimore)*. 2021;100(38):e22571. [PubMed ID: 34559092]. [PubMed Central ID: PMC8462638]. <https://doi.org/10.1097/MD.00000000000022571>.
- Ojha V, Mani A, Pandey NN, Sharma S, Kumar S. CT in coronavirus disease 2019 (COVID-19): a systematic review of chest CT findings in 4410 adult patients. *Eur Radiol*. 2020;30(11):6129-38. [PubMed ID: 32474632]. [PubMed Central ID: PMC7261039]. <https://doi.org/10.1007/s00330-020-06975-7>.
- Wan S, Li M, Ye Z, Yang C, Cai Q, Duan S, et al. CT Manifestations and Clinical Characteristics of 1115 Patients with Coronavirus Disease 2019 (COVID-19): A Systematic Review and Meta-analysis. *Acad Radiol*. 2020;27(7):910-21. [PubMed ID: 32505599]. [PubMed Central ID: PMC7200137]. <https://doi.org/10.1016/j.acra.2020.04.033>.
- Ebrahimpour L, Marashi M, Zamanian H, Abedi M. Computed tomography findings in 3,557 COVID-19 infected children: a systematic review. *Quant Imaging Med Surg*. 2021;11(11):4644-60. [PubMed ID: 34737930]. [PubMed Central ID: PMC8511729]. <https://doi.org/10.21037/qims-20-1410>.

-
24. Kumar J, Meena J, Yadav A, Yadav J. Radiological Findings of COVID-19 in Children: A Systematic Review and Meta-Analysis. *J Trop Pediatr*. 2021;**67**(3):fmaa045. [PubMed ID: [32692815](#)]. [PubMed Central ID: [PMC7454935](#)]. <https://doi.org/10.1093/tropej/fmaa045>.
25. Ghodsi A, Azarfar A, Ghahremani S. A Review of Coronavirus Disease (COVID-19) in Children. *J Ped Nephrol*. 2020;**8**(3):1-6. <https://doi.org/10.22037/jpn.v8i3.31060>.
26. Awulachew E, Diriba K, Anja A, Getu E, Belayneh F. Computed Tomography (CT) Imaging Features of Patients with COVID-19: Systematic Review and Meta-Analysis. *Radiol Res Pract*. 2020;**2020**:1023506. [PubMed ID: [32733706](#)]. [PubMed Central ID: [PMC7378588](#)]. <https://doi.org/10.1155/2020/1023506>.