



Home Mechanical Ventilation in Children: Complications and Benefits—A Two-Year Experience at a Referral Children's Hospital

Majid Keivanfar^{1,2}, Mahnaz Kheiri^{3,2}, Sharareh Babaie^{4,2}, Mohsen Reisi^{1,2,*}

¹ Department of Pediatric Pulmonology, Pediatrics Department, Emam Hossein Children's Hospital, Isfahan University of Medical Sciences, Isfahan, Iran

² Child Growth and Development Research Center, Research Institute for Primordial Prevention of Non-communicable Disease, Isfahan University of Medical Sciences, Isfahan, Iran

³ Pediatrics Department, Emam Hossein Children's Hospital, Isfahan University of Medical Sciences, Isfahan, Iran

⁴ Department of Pediatric Intensive Care Unit, Pediatrics Department, Emam Hossein children's hospital, Isfahan University of Medical Sciences, Isfahan, Iran

*Corresponding author: Department of Pediatric Pulmonology, Pediatrics Department, Emam Hossein children's hospital, Isfahan University of Medical Sciences, Isfahan, Iran. Child Growth and Development Research Center, Research Institute for Primordial Prevention of Non-communicable Disease, Isfahan University of Medical Sciences, Isfahan, Iran. Email: mohsenreisi72@yahoo.com

Received 2023 December 19; Revised 2024 March 18; Accepted 2024 April 28.

Abstract

Background: The rising prevalence of chronic respiratory failure in pediatric patients necessitates a detailed evaluation of home mechanical ventilation (HMV).

Objectives: This study assesses the advantages of HMV devices, focusing on cases from Imam Hussein Children's Hospital.

Methods: This research involved 20 children who required long-term ventilation and were admitted to both general and intensive care units at Imam Hussein Children's Hospital in Isfahan. Before discharge, a team of HMV experts trained the parents on the proper use of the ventilation devices. Post-discharge, the educational needs of the parents were assessed continuously, with regular home visits conducted by a nurse and a representative from the ventilator supplier.

Results: Approximately 73% of the children demonstrated therapeutic effectiveness following HMV implementation. About 35% of parents observed no adverse effects in their children. The average cost savings from reduced ICU rehospitalizations was significant at 347.59%, and there was a 65.09% reduction in the costs associated with general ward hospitalizations.

Conclusions: The adoption of HMV has led to improved health outcomes, substantial reductions in medical expenses, and shorter hospital stays. It is advisable to plan for broader implementation of HMV in children who require long-term ventilation.

Keywords: Mechanical Ventilation, Children, Medical Costs, Hospital Stay

1. Background

Home mechanical ventilation (HMV) represents an innovative approach to providing respiratory support in children with various chronic respiratory diseases. Mechanical ventilation is crucial for the stable support of individuals with respiratory failure, particularly those with chronic neuromuscular or neurological disorders (1). Additionally, HMV has been proven to enhance quality of life and reduce hospitalization burdens for these patients (2). By allowing extended ventilation periods, HMV significantly enhances life quality and survival rates for infants and children from one month of age and older (3-5).

The American respiratory care guidelines emphasize several long-term objectives for HMV, such as enhancing life sustainability, improving overall quality of life, reducing complication rates, promoting optimal growth and development, and providing a cost-effective therapeutic option (6). Candidates for HMV include patients aged three months and older (6, 7). The escalating costs of hospital care and advancements in positive pressure ventilation technology for home use have encouraged the transition of patients from hospital settings to home care. Nevertheless, this transition can fail or lead to ICU readmissions if parents are not adequately trained in HMV management or lack sufficient support (8).

The application of HMV varies; for instance, a study in Italy by Ottonella et al. involved 20 children needing long-term mechanical ventilation. Among these, 65% utilized non-invasive ventilation (NIV), while 35% required invasive mechanical ventilation (IMV). Of these, only 10% used their devices during the day, 20% used them for 8 to 12 hours, and the majority, 70%, used them solely at night (9). Given the substantial benefits of HMV and the scarcity of in-depth studies on the pediatric population in Iran, a comprehensive evaluation of this demographic is critical.

2. Objectives

This study aimed to assess the advantages of using home mechanical ventilators at Imam Hossein Children's Hospital.

3. Methods

In this case study, children requiring mechanical ventilation in the hospital for over three months were evaluated by an expert team consisting of a pediatric ICU subspecialist, a pediatric pulmonologist, and a pediatric anesthesiology subspecialist. They were deemed suitable for HMV, following ethical approval by the esteemed Deputy for Research and Technology at Isfahan University of Medical Sciences, under ethical code [IR.MUI.MED.REC.1400.577](#). Inclusion criteria included a willingness to participate, a need for long-term ventilator support, the capability of parents to provide HMV, medical stability, no changes in mechanical ventilation settings and oxygen requirements for two weeks prior to discharge, and parental cooperation. Exclusion criteria encompassed the death of a child during the study, relocation from Isfahan to another region, and parental withdrawal from the study.

Key aspects assessed were the age and gender of the patients, disease prevalence, treatment costs, and a cost comparison between hospital and home care. After patient selection by a pediatric pulmonologist, additional measures included home visits by a nurse and a ventilator company representative when necessary. The oversight and execution of the process involved a team comprising a pediatric pulmonologist, a pediatric resident, a nurse specialized in respiratory care, a physical therapist, a speech therapist, a nutritionist, and a ventilator company technician.

The required equipment included (1) a ventilator, either non-invasive for use with a mask or invasive via a tracheostomy; (2) an oxygen source; (3) a pulse oximeter; (4) an Ambo bag; (5) humidifiers and ventilator accessories for tracheostomy use; (6) a manual suction device, if needed; (7) feeding equipment for non-oral intake; and (8) a first aid kit with resuscitation supplies. Parents and caregivers received comprehensive training on operating the ventilator, suction devices, tracheostomy care (if applicable), nebulizers, and non-oral feeding methods. This training ensured high-quality, effective care. Before discharge, the team assessed and confirmed the caregivers' proficiency in these skills to ensure a safe and gradual transition to home care.

In addition, caregivers received training on strategies to enhance the child's mental and physical health during care. The care team supported both the patient and parents during the transition from hospital to home. The settings of the home ventilator, installed by hospital experts, were fixed and non-adjustable to prevent unauthorized adjustments. The care team educated parents on how to respond to specific alarm situations as necessary. After discharge, nurses provided regular reports, assessments, and tests. Given these factors and the importance of educating parents, efforts were intensified to enhance parents' understanding and proficiency in operating the home ventilator, thereby gradually reducing the frequency of care team visits to the child's home. The frequency of home visits was determined based on the needs expressed by the caregivers and their competence in managing the alarms of the HMV device. For data analysis, descriptive statistics such as mean, standard deviation, maximum, minimum, frequency, and percentage were utilized, employing SPSS software version 23.

4. Results

This study encompassed 20 patients with an average age of 4.092 ± 695.8 years, ranging from 15 months to 16 years. The mean weight of the children was 23.85 ± 14.914 kg, their average height was 115.3 ± 28.2 cm, and the mean BMI was 16.9 ± 6.27 (Table 1). Twelve children (60%) were female, and eight (40%) were male.

Table 2 indicates that 22.22% of the cases requiring HMV were due to myopathy, 16.67% to sleep apnea, 44.44% to spinal muscular atrophy, and 27.78% to laryngomalacia-tracheomalacia. The distribution of device use based on the method of treatment showed

Table 1. Characteristics of the Studied Children Requiring Long-Term Mechanical Ventilation

Variables	Min-Max	Mean \pm SD	Median (IQR)
Age, y	1.3 - 16	8.695 \pm 4.092	7 (8.7)
Weight, kg	10 - 64	23.85 \pm 14.914	21.5(16.8)
Height, m	0.7 - 1.52	1.153 \pm 0.282	1.23 (0.43)
BMI, kg/m ²	10 - 32.6	16.9 \pm 6.27	13.74 (8)

Table 2. Side Effects of Home Mechanical Ventilation (HMV) and the Cost of Long-Term HMV in Children Requiring Prolonged Mechanical Ventilation

Sex	Age	Conditional Disease	Type of HMV	Cost of HMV (Million Rial)	Side Effects	Type of Side Effects
Female	3.1	Myopathy	Tracheostomy	25	Yes	Abdominal bloating or abdominal pain that causes intolerance to feeding or stopping the use of the device.
Female	15	Sleep apnea	Musk	6	Yes	Abdominal bloating or abdominal pain that causes intolerance to feeding or stopping the use of the device/ Dry or watery eyes.
Female	16	Spinal muscular atrophy	Musk	10	No	-
Male	2.1	Laryngeomalasy-Tracheomalasy	Musk	25	Yes	Intolerance of masks
Male	1.3	Laryngeomalasy-Tracheomalasy	Musk	10	Yes	Dry or watery eyes
Male	5	Spinal muscular atrophy	Tracheostomy	60	No	-
Female	11.5	Sleep apnea	Musk	10	No	-
Female	10	Spinal muscular atrophy	Musk	30	No	-
Female	6	Spinal muscular atrophy	Musk	30	Yes	Intolerance of masks
Female	7	Laryngeomalasy-Tracheomalasy	Musk	30	Yes	Dry or watery eyes
Female	7	Spinal muscular atrophy	Tracheostomy	9	Yes	Abdominal bloating or abdominal pain that causes intolerance to feeding or stopping the use of the device.
Male	10	Sleep apnea	Musk	10	No	-
Male	5	Myopathy	Musk	30	No	-
Male	8	Laryngeomalasy-Tracheomalasy	Musk	60	Yes	Dry or watery eyes
Male	10	Spinal muscular atrophy	Musk	6	Yes	Dry or watery eyes
Male	9	Spinal muscular atrophy	Tracheostomy	25	Yes	Abdominal bloating or abdominal pain that causes intolerance to feeding or stopping the use of the device/Dry or watery eyes
Female	2	Laryngeomalasy-Tracheomalasy	Musk	30	No	-
Female	7	Spinal muscular atrophy	Tracheostomy	10	Yes	Dry or watery eyes
Female	5	Myopathy	Musk	15	No	-
Female	10	Myopathy	Tracheostomy	30	Yes	Abdominal bloating or abdominal pain that causes intolerance to feeding or stopping the use of the device.

that 30% of the patients utilized tracheostomy, and 70% used mask ventilation. Additionally, 65% of the patients experienced side effects while using the HMV, while 35% reported no side effects. The most commonly reported side effect was dryness or irritation of the eyes. These findings underscore the importance of vigilant monitoring and management of potential side effects associated with HMV use, especially concerning eye health (Table 2).

Approximately 73% of patients demonstrated therapeutic effectiveness after utilizing HMV. However, 27% of patients showed no improvement, suggesting that factors such as comorbidities, disease severity, or individual patient characteristics might influence their response to HMV. This underscores the necessity for tailored care and ongoing monitoring (note: The content is a general translation as the exact context of "Figure 1" and the medical condition's specifics are not provided.).

Table 3. Average and Standard Deviation of Re-hospitalization Costs in General and ICU Wards Before and After Using HMV^a

Variables	Before Utilizing the HMV	After Utilizing the HMV	Total Change Percent
Ward	35.95 ± 45.36	23.4 ± 32.45	65.09
ICU	280.5 ± 663.92	9.75 ± 14.25	347.59

^a Values are expressed as mean ± SD.


**Figure 1.** Comparing the results of using home mechanical ventilation (HMV)

Table 3 reveals that the average cost of re-hospitalization in the ICU decreased significantly by 347.59%, a notable reduction. Similarly, the cost of re-hospitalization in the general ward was reduced by 65.09%.

5. Discussion

This study was conducted to assess the benefits of using HMV devices at Imam Hussein Children's Hospital. The findings highlight the importance of regular assessments and adjustments to treatment parameters to optimize patient outcomes and provide the highest quality of care. Moreover, continuous advancements in HMV technology and further research are crucial for improving response rates and addressing the needs of patients who do not initially benefit from this treatment. For instance, a study from Massachusetts in 2005 observed that the incidence of long-term ventilation tripled in patients who required respiratory support for over 15 years (10). Additionally, Povitz et al. reported that the annual incidence of mechanical home

ventilation authorization increased from 1.8 per 100 000 individuals in 2000 to 5.0 per 100 000 in 2012, marking an annual increase of about 0.3 per 100 000 people per year (11).

Rose et al. (12) reported a 61% efficacy rate for HMV, whereas this study observed an improvement of up to 73% in children using the device. One potential reason for these differing outcomes might be the age range of the participants; Rose et al.'s study included all age groups, which could have affected their results. Additionally, the variability in outcomes might stem from inadequate caregiver training. In this study, caregivers received educational support and regular visits from the care team, enhancing their ability to manage HMV effectively. In contrast, feedback from participants in Rose et al.'s study indicated that only 45% received skills assessments and follow-up training, suggesting that insufficient ongoing education may contribute to inconsistent results.

Fauroux et al. (13), Windisch et al. (14), Sterni et al. (7), and Kwak (15) have highlighted that while HMV allows for a more normal daily life compared to institutional or hospital-based ventilation, it also introduces significant psychological, physiological, and social challenges for children and their families. Valko et al. (16) observed a 10.5% improvement in quality of life over six months, underscoring the potential benefits of HMV.

Povitz et al. (11) noted a decrease in the prehospitalization rate from 39.8% to 29.9% after the introduction of HMV, which signifies a substantial reduction in the need for re-hospitalization, falling to less than 90%. This improvement could largely be attributed to the continuous educational support and frequent home visits provided by the care team. Correspondingly, Amin et al. (17) reported that the median length of stay in acute and rehabilitation clinics decreased significantly after tracheostomy insertion, from an initial median of 162.0 days (IQR 98.0 to 275.0 days) to 97.0 days (IQR 69.0 to 210.0 days). Remarkably, in this study, the average duration of hospital stays in

intensive care units was reduced from 394 days to just 13 days, illustrating a significant reduction.

The research by Hazenberg et al. (18) suggested that patients using HMV experience fewer side effects and enjoy a better quality of life than others. This aligns with the findings of the current study, where 35% of participants reported no side effects. For the remaining 65%, side effects were typically minor and manageable through education and proper care. However, Mattson et al. (19) noted that "children on mechanical ventilation at home reported a lower quality of life than healthy children and those with other chronic diseases." This disparity may stem from inadequate follow-up and parental support in some cases. A crucial and exemplary aspect of this study was the presence of a supportive care team that began its involvement before discharge, ensuring parents were well-prepared to manage their children's care at home with HMV. Regular follow-up visits were conducted to monitor the child's condition, with ongoing parental support until they felt confident in their caregiving abilities.

Regarding medical costs, this study found that hospital expenses for patients totaled 349.05 million rials, which significantly decreased to 34.7 million rials with home care. Similarly, Ottonello et al. (9) reported, "We compared the cost of home care with the actual hospital costs (1324 €/day in the intensive care unit of the G. Gaslini Children's Hospital) and found a significant economic advantage of home care (7593 €/week)."

5.1. Study Limitations

A key limitation of this study was the absence of a control group. It is suggested that future research replicates this study as a clinical trial to better control confounding variables and strengthen the generalizability of the findings.

5.2. Conclusions

This study has provided valuable insights into the use of HMV for patients requiring long-term ventilation. The findings indicate the effectiveness of HMV in improving health outcomes for the majority of patients, significantly reducing medical treatment costs, and shortening hospital stays. These results underscore the importance of HMV and suggest the need for more detailed research in this area. Considering the growing number of children requiring HMV, these findings can

guide the development of essential support programs for these patients and their families.

Acknowledgements

This study is the result of a research project with ethical code [IR.MUI.MED.REC.1400.577](#), approved by Isfahan University of Medical Sciences. We appreciate both the financial and moral support provided.

Footnotes

Authors' Contribution: Study concept and design by M. K.; analysis and interpretation of data by M.KH.; manuscript drafting by SH. B.; critical manuscript revision for important intellectual content by M.R.

Conflict of Interests Statement: No conflicts of interests were reported.

Data Availability: The dataset from this study is available upon request from the corresponding author either during submission or post-publication. The data are not publicly available due to the presence of unpublished results.

Ethical Approval: The study protocol received approval from the Institutional Review Board of Isfahan University of Medical Sciences, Isfahan, Iran (code: [IR.MUI.MED.REC.1400.577](#)). Informed consent was obtained from all participants.

Funding/Support: This study was funded by Isfahan University of Medical Sciences, Isfahan, Iran.

Informed Consent: Informed consent was obtained.

References

- Lewarski JS, Gay PC. Current issues in home mechanical ventilation. *Chest*. 2007;**132**(2):671-6. [PubMed ID: [17699139](#)]. <https://doi.org/10.1378/chest.07-0558>.
- Wise MP, Hart N, Davidson C, Fox R, Allen M, Elliott M, et al. Home mechanical ventilation. *BMJ*. 2011;**342**:d1687. [PubMed ID: [21471174](#)]. <https://doi.org/10.1136/bmj.d1687>.
- Ferreira C, Moita J. Home Mechanical Ventilation and Quality of Life in Neuromuscular Patients During Noninvasive Mechanical Ventilation: New Trends and Key Practical Topics. In: Antonio M E, editor. *Noninvasive Mechanical Ventilation Theory, Equipment, and Clinical Applications*. New York City, USA: Springer; 2016. p. 781-90. https://doi.org/10.1007/978-3-319-21653-9_91.
- Sahetya S, Allgood S, Gay PC, Lechtzin N. Long-Term Mechanical Ventilation. *Clin Chest Med*. 2016;**37**(4):753-63. [PubMed ID: [27842754](#)]. [PubMed Central ID: [PMC7273179](#)]. <https://doi.org/10.1016/j.ccm.2016.07.014>.

5. Agarwal A, Gupta P. Home Mechanical Ventilation in Ventilator-Dependent Children: Criteria, Outcome, and Health Organization. In: Antonio M. Esquinas, editor. *Noninvasive Mechanical Ventilation and Difficult Weaning in Critical Care Key Topics and Practical Approaches*. New York City, USA: Springer; 2016. p. 439-47. https://doi.org/10.1007/978-3-319-04259-6_54.
6. American Association for Respiratory Care. Long-Term Invasive Mechanical Ventilation in the Home—2007 Revision & Update. *Respiratory Care*. 2007;**52**(8):1056-62.
7. Sterni LM, Collaco JM, Baker CD, Carroll JL, Sharma GD, Brozek JL, et al. An Official American Thoracic Society Clinical Practice Guideline: Pediatric Chronic Home Invasive Ventilation. *Am J Respir Crit Care Med*. 2016;**193**(8):e16-35. [PubMed ID: 27082538]. [PubMed Central ID: PMC5439679]. <https://doi.org/10.1164/rccm.201602-0276ST>.
8. Paulides FM, Plotz FB, Verweij-van den Oudenrijn LP, van Gestel JP, Kampelmacher MJ. Thirty years of home mechanical ventilation in children: escalating need for pediatric intensive care beds. *Intensive Care Med*. 2012;**38**(5):847-52. [PubMed ID: 22476447]. [PubMed Central ID: PMC3332376]. <https://doi.org/10.1007/s00134-012-2545-9>.
9. Ottonello G, Ferrari I, Pirroddi IM, Diana MC, Villa G, Nahum L, et al. Home mechanical ventilation in children: retrospective survey of a pediatric population. *Pediatr Int*. 2007;**49**(6):801-5. [PubMed ID: 18045275]. <https://doi.org/10.1111/j.1442-200X.2007.02463.x>.
10. Graham RJ, Flegler EW, Robinson WM. Chronic ventilator need in the community: a 2005 pediatric census of Massachusetts. *Pediatr*. 2007;**119**(6):e1280-7. [PubMed ID: 17485451]. <https://doi.org/10.1542/peds.2006-2471>.
11. Povitz M, Rose L, Shariff SZ, Leonard S, Welk B, Jenkyn KB, et al. Home Mechanical Ventilation: A 12-Year Population-Based Retrospective Cohort Study. *Respir Care*. 2018;**63**(4):380-7. [PubMed ID: 29208755]. <https://doi.org/10.4187/respcare.05689>.
12. Rose L, McKim DA, Katz SL, Leasa D, Nonoyama M, Pedersen C, et al. Home mechanical ventilation in Canada: a national survey. *Respir Care*. 2015;**60**(5):695-704. [PubMed ID: 25587173]. <https://doi.org/10.4187/respcare.03609>.
13. Fauroux B, Abel F, Amaddeo A, Bignamini E, Chan E, Corel L, et al. ERS statement on paediatric long-term noninvasive respiratory support. *Eur Respir J*. 2022;**59**(6). [PubMed ID: 34916265]. <https://doi.org/10.1183/13993003.01404-2021>.
14. Windisch W, Geiseler J, Simon K, Walterspacher S, Dreher M; on behalf of the Guideline Commission. German National Guideline for Treating Chronic Respiratory Failure with Invasive and Non-Invasive Ventilation: Revised Edition 2017 - Part 1. *Respir*. 2018;**96**(1):66-97. [PubMed ID: 29945148]. <https://doi.org/10.1159/000488001>.
15. Kwak S. Home mechanical ventilation in children with chronic respiratory failure: a narrative review. *J Yeungnam Med Sci*. 2023;**40**(2):123-35. [PubMed ID: 35618662]. [PubMed Central ID: PMC10076918]. <https://doi.org/10.12701/jyms.2022.00227>.
16. Valko L, Baglyas S, Gyarmathy VA, Gal J, Lorx A. Home mechanical ventilation: quality of life patterns after six months of treatment. *BMC Pulm Med*. 2020;**20**(1):221. [PubMed ID: 32807149]. [PubMed Central ID: PMC7433042]. <https://doi.org/10.1186/s12890-020-01262-z>.
17. Amin R, Sayal A, Syed F, Daniels C, Hoffman A, Moraes TJ, et al. How long does it take to initiate a child on long-term invasive ventilation? Results from a Canadian pediatric home ventilation program. *Can Respir J*. 2015;**22**(2):103-8. [PubMed ID: 25848720]. [PubMed Central ID: PMC4390005]. <https://doi.org/10.1155/2015/107914>.
18. Hazenberg A, Kerstjens HA, Prins SC, Vermeulen KM, Wijkstra PJ. Initiation of home mechanical ventilation at home: a randomised controlled trial of efficacy, feasibility and costs. *Respir Med*. 2014;**108**(9):1387-95. [PubMed ID: 25081652]. <https://doi.org/10.1016/j.rmed.2014.07.008>.
19. Mattson J, Lunnellie J, Lofholm T, Andersson ES, Aune RE, Bjorling G. Quality Of Life in Children With Home Mechanical Ventilation - A Scoping Review. *SAGE Open Nurs*. 2022;**8**:23779608221094500. [PubMed ID: 35493542]. [PubMed Central ID: PMC9047042]. <https://doi.org/10.1177/23779608221094522>.