



# Umbilical Cord Zinc Levels and Neonatal Anthropometric Measures: A Cross-Sectional Study in Iran

Mahbod Kaveh <sup>1</sup>, Marzieh Darabi <sup>2</sup>, Razieh Sangsari <sup>3</sup>, Roksana Moeini <sup>4</sup>, Maryam Veysizadeh <sup>1</sup>, Mohammad Reza Zarkesh <sup>2, 4, \*</sup>, Mamak Shariat <sup>2</sup>

<sup>1</sup> Department of Pediatrics, Bahrami Children's Hospital, Tehran University of Medical Sciences, Tehran, Iran

<sup>2</sup> Maternal, Fetal and Neonatal Research Center, Family Health Research Institute, Tehran University of Medical Sciences, Tehran, Iran

<sup>3</sup> Children's Medical Center, Pediatrics Centre of Excellence, Tehran University of Medical Sciences, Tehran, Iran

<sup>4</sup> Department of Neonatology, Yas Hospital Complex, Tehran University of Medical Sciences, Tehran, Iran

\*Corresponding Author: Maternal, Fetal and Neonatal Research Center, Family Health Research Institute, Tehran University of Medical Sciences, Tehran, Iran; Department of Neonatology, Yas Women Hospital, Tehran University of Medical Sciences, Tehran, Iran. Email: zarkesh@tums.ac.ir

Received: 24 June, 2024; Revised: 16 November, 2024; Accepted: 3 December, 2024

## Abstract

**Background:** Zinc (Zn) is one of the trace elements in the body that, despite its low level, plays important roles in molecular and cellular activities. A few investigations have assessed the correlations between cord blood Zn levels and neonatal anthropometric measures; however, the results are conflicting.

**Objectives:** The purpose of the present study was to investigate the relationships between the Zn level in the umbilical cord blood and the birth weight, height, and head circumference of term newborns.

**Methods:** A cross-sectional study was conducted on full-term newborns in Tehran, Iran, in 2021. Neonates' Zn levels in umbilical cord blood were evaluated. Data related to perinatal and neonatal variables such as the number of pregnancies (gravida), birth height, weight, and head circumference were recorded, and their relationships with umbilical cord Zn levels were assessed. P-values < 0.05 were considered statistically significant.

**Results:** One hundred mothers with a mean age of 30.792 ± 5.908 years and a parity of 2.693 ± 1.181 were included. Of all the newborns, 68% were boys, and 2% had low birth weight. The mean level of cord blood Zn was 107.703 ± 27.843 µg/dL. Evaluating the correlations between cord blood Zn levels and maternal and neonatal qualitative variables, data analysis showed no relationships between Zn levels and neonatal sex (P = 0.70), low birth weight (P = 0.87), and maternal underlying disease (P = 0.49). Data analysis regarding correlations between quantitative variables also showed no significant correlations between cord blood Zn levels and birth weight (P = 0.466), height (P = 0.466), and head circumference (P = 0.925).

**Conclusions:** Although adequate Zn levels were observed in all neonates, these levels did not significantly influence neonatal anthropometric measures or immediate health indicators. Further studies with larger sample sizes are recommended to provide more comparative data, particularly in relation to long-term growth outcomes.

**Keywords:** Umbilical Cord, Newborn, Zinc

## 1. Background

Zinc (Zn) is an essential micronutrient that plays an important role at different molecular and cellular levels. This element is involved in protein synthesis, gene expression, cellular metabolism, nuclear division, signal transduction, hormonal structure, and enzymatic function (1-3). Despite its abundant functions, the level of Zn is constantly maintained within narrow biological limits (4, 5). Studies have shown the role of Zn in fetal

and neonatal growth and development (6). Maternal blood Zn levels decrease until 35 weeks of gestation due to increased blood volume, organ metabolism, and urinary excretion, as well as hormonal alterations and active maternal-placental transport. In addition to inadequate Zn intake, perinatal co-morbidities, such as digestive complications, may put pregnant women at a higher risk of Zn deficiency (6). Previous studies have pointed to the possible teratogenic effects related to Zn deficiency, which can result in adverse pregnancy

outcomes (7, 8). Preterm birth, preterm premature rupture of membranes, preeclampsia, small for gestational age, intrauterine growth restriction, and low Apgar scores are several reported maternal and neonatal morbidities associated with prenatal Zn deficiency (8, 9).

Evidence has also shown low serum Zn concentrations in neonates with higher weight gain due to increased cellular metabolism and enzymatic activities. Accordingly, Zn deficiency may result in delayed neonatal growth (10). It has also been demonstrated that Zn intake during the first 12 months could affect growth outcomes in both term and preterm infants (11). As Zn deficiency may limit infant growth, this diagnosis should be considered, particularly for premature infants with failure to thrive despite apparently receiving enough calories and macronutrients (12).

Concerning the relationship between Zn levels and anthropometric measures, Gomez et al. in 2015 showed a positive significant correlation between cord blood Zn levels and birth weight ( $P = 0.283$ ;  $P = 0.005$ ) (13). Khoushabi et al. also demonstrated that of 60 included neonates, 12% were low birth weight, while the others had normal birth weight. The results of the study indicated that the mean Zn levels in the cord blood of neonates with normal birth weight ( $88.3 \pm 12$  mg/dL) were significantly higher ( $P < 0.05$ ) than those of their low-birth-weight counterparts ( $70.0 \pm 9.9$  mg/dL) (14). This significant relationship between cord blood Zn levels and newborn birth weight was also confirmed by a systematic review and meta-analysis by Atazadegan et al. (15). Nanbakhsh and Tabrizi in 2017 demonstrated the effects of maternal serum Zn concentrations on neonatal birth weight. According to the findings of this study, mothers with serum Zn levels  $> 70$   $\mu\text{g/dL}$  had neonates with birth weights  $> 3500$  g, while mothers with serum Zn levels  $< 60$   $\mu\text{g/dL}$  had neonates with birth weights  $< 3000$  g. In addition, low birth weight neonates had significantly lower umbilical cord Zn levels than neonates with normal birth weights ( $19.86 \pm 79.16$  vs.  $95.14 \pm 17.56$   $\mu\text{g/dL}$ ;  $P < 0.02$ ) (16).

## 2. Objectives

Zinc deficiency is one of the most frequent deficiencies among women, particularly pregnant women in developing countries, including Iran (17, 18). Accordingly, the World Health Organization promotes prenatal micronutrient supplementation containing Zn for all women (19). Moreover, despite a few investigations that have assessed the correlations

between cord blood Zn levels and anthropometric measures, the results are conflicting (13, 20-25). Hence, the present study was carried out to examine the potential influence of cord blood Zn on term neonates' growth parameters.

## 3. Methods

A cross-sectional study was conducted at Yas Complex Hospital, affiliated with Tehran University of Medical Sciences (Tehran, Iran), in 2021. The study population included mothers and their singleton, term neonates (gestational age  $> 37$  weeks). All mothers received supplements during their pregnancy. Written consent was obtained before the participants' enrollment. Exclusion criteria included multiple gestations, preterm birth, placental abnormalities, and a positive history of maternal complications such as severe anemia and chronic disease.

All demographic and clinical data related to the mothers and neonates were extracted from medical records and recorded in a checklist. Neonatal anthropometric values, including birth weight, height, and head circumference, were also determined using a digital weight scale (g) and measuring tape (cm), respectively.

Immediately after birth, blood sampling was performed, and 3 milliliters of umbilical cord blood were drawn, centrifuged, stored at  $-20^\circ\text{C}$ , and sent to the hospital laboratory. Using the atomic absorption spectrophotometry method, Zn levels in cord blood ( $\mu\text{g/dL}$ ) were measured (9, 26). Serum Zn levels below 65  $\mu\text{g/dL}$  were considered deficient, while levels above 65  $\mu\text{g/dL}$  were considered normal (21).

All gathered data related to Zn levels in cord blood, neonatal measurements, and anthropometric values were analyzed to determine possible correlations between Zn levels and neonatal anthropometric measures. Moreover, correlations between Zn levels and other variables were also assessed to identify any potential influencing factors.

### 3.1. Sample Size

Mohamed et al. in 2019 (24) demonstrated a Pearson correlation of 0.03 between Zn and weight. Using the following formula and considering a power of 80% and a confidence interval of 95%, the sample size was estimated to be 95.

$$n = \left[ \frac{z_{1-\frac{\alpha}{2}} + z_{1-\beta}}{C} \right]^2 \quad (1)$$

$$C = 0.5 \times L_n \left[ \frac{1+r}{1-r} \right] \quad (2)$$

$$r = 0.09; Z_{1-\alpha/2} = 1.96; Z_{1-\beta} = 0.84; n = 95.$$

### 3.2. Statistical Analysis

Data analysis was performed using the Statistical Package for the Social Sciences (SPSS Inc., Chicago, USA), version 26.0. Descriptive data were presented in numbers (percent) and Mean  $\pm$  SD. The results of the non-parametric Kolmogorov-Smirnov test showed that the variables did not follow a normal distribution. Therefore, the Mann-Whitney and Pearson Correlation tests were used to analyze all data without normal distribution. P-values  $<$  0.05 were considered significant.

### 3.3. Ethical Consideration

This study was approved in accordance with the Helsinki Declaration by the Ethics Board of Tehran University of Medical Sciences (ID Code: [IR.TUMS.IKHC.REC.1400.282](#)). Written consent was obtained from the included participants, no extra cost was imposed, and all data were treated as confidential.

## 4. Results

One hundred mothers with a mean age of  $30.792 \pm 5.908$  years and a parity of  $2.693 \pm 1.181$  were included. Sixty-eight of their neonates were male, and two newborns had low birth weight. The mean level of cord blood Zn was  $107.703 \pm 27.843$   $\mu\text{g/dL}$  (Min: 65 and Max: 225), and none of the neonates were Zn deficient. The mean Apgar scores at the first and fifth minutes, as well as birth weight, height, and head circumference, were  $8.945 \pm 0.271$ ,  $9.956 \pm 0.253$ ,  $3301.020 \pm 354.163$  g,  $50.816 \pm 2.549$  cm, and  $35.319 \pm 1.297$  cm, respectively. The mean level of Zn in the neonates was  $107.240 \pm 27.590$  (65.00 - 225.00)  $\mu\text{g/dL}$ . [Figure 1](#) shows cord blood Zn levels in neonates with different birth weight, height, and head circumference parameters.

Evaluating the correlations between cord blood Zn levels and maternal and neonatal qualitative variables, data analysis showed no relationships between Zn levels and neonates' sex ( $P = 0.70$ ), low birth weight ( $P = 0.87$ ), and maternal underlying disease ( $P = 0.49$ ). Detailed data are shown in [Table 1](#).

Analysis of data regarding correlations between quantitative variables also showed no significant correlations between cord blood Zn levels and maternal age ( $P = 0.491$ ), number of gravidity ( $P = 0.587$ ), and neonates' first ( $P = 0.118$ ) and fifth minute ( $P = 0.207$ )


Apgar scores. Neonates with higher anthropometric measures had lower cord blood Zn concentrations; however, the Zn values did not significantly alter neonates' birth weight ( $P = 0.466$ ), height ( $P = 0.466$ ), and head circumference ( $P = 0.925$ ). Cord blood Zn levels were also not correlated with blood pH ( $P = 0.532$ ) or base excess ( $P = 0.068$ ). Detailed data are shown in [Table 2](#).

## 5. Discussion

The present study aimed to assess the potential correlations between umbilical cord blood Zn concentrations and neonatal anthropometric parameters. Although the beneficial effects of micronutrients like Zn on pregnancy and neonatal outcomes have been reported by previous evidence ([27-29](#)), our results align with the results of a meta-analysis ([19](#)) suggesting that cord blood Zn levels may not exert a significant influence on neonatal growth outcomes.

The mean cord blood Zn level was found to be 107.7  $\mu\text{g/dL}$ , with none of the neonates showing Zn deficiency, suggesting that maternal supplementation during pregnancy was effective in maintaining adequate Zn status in newborns. Fortunately, most pregnant women in our center received prenatal care from the first trimester of gestation. Supplements, including folic acid and iron, were administered to them based on the national protocol. In addition, during the second and third trimesters, pregnant women routinely took prenatal multivitamins containing vitamins and minerals. While this aligns with studies that emphasize the benefits of maternal Zn supplementation for fetal health ([30, 31](#)), the lack of a significant association between cord blood Zn and neonatal anthropometric measures may indicate that Zn's role in fetal growth is complex. It is possible that Zn may play a more prominent role in postnatal growth and development ([30](#)), rather than exerting measurable effects on birth parameters. It also seems that larger sample sizes, through affecting variables, may influence the results.

The results of this study were in agreement with the results of Daniali et al. ([22](#)), who, through a cross-sectional study, evaluated the relationships between cord blood Zn concentrations and anthropometric measures in 226 neonates. The authors demonstrated no significant correlations between cord blood Zn levels and neonate height ( $P = 0.792$ ), head circumference ( $P = 0.697$ ), abdominal circumference ( $P = 0.785$ ), chest circumference ( $P = 0.498$ ), sex ( $P = 0.632$ ), or Apgar score ( $P = 0.673$ ). However, cord blood Zn level was significantly correlated with the neonate's birth weight ( $\beta = 0.178$ ;  $P = 0.02$ ). In accordance with our findings, the



**Figure 1.** Cord blood Zn levels in neonates with different birth weight (A), height (B), and head circumference (C) parameters

**Table 1.** The Relationships Between Cord Blood Zinc Levels and Maternal/Neonatal Qualitative Variables

Variables	Mean $\pm$ SD	P-Value
<b>Neonate's gender</b>		0.70
Male	106.647 $\pm$ 27.100	
Female	108.500 $\pm$ 29.006	
<b>Low birth weight</b>		0.87
Yes	109.500 $\pm$ 31.819	
No	108.010 $\pm$ 28.195	
<b>Maternal underlying disease</b>		0.49
Yes	105.1064 $\pm$ 25.031	
No	109.9630 $\pm$ 30.129	

results of a systematic review (32) have shown that maternal Zn supplementation did not significantly improve neonatal outcomes.

Amini et al. (23) demonstrated an inverse correlation between cord blood Zn levels and neonate birth weight ( $P = 0.008$ ); however, no significant relationships were observed between blood Zn concentrations and other anthropometric measures like length or head circumference ( $P > 0.05$ ). In contrast to our findings,

several studies have shown diverse results. A systematic review (33) delineated a significant correlation between maternal Zn supplementation and neonate birth weight or length in 3 out of 9 included studies, while these findings were not confirmed by the other 6 studies. The results of a systematic review and meta-analysis (15) also demonstrated a significant correlation between cord blood Zn and neonate birth weight. Bayomy et al. (34) showed a significant and positive correlation between

**Table 2.** Correlations Between Cord Blood Zinc Levels and Different Variables

Variables	Correlation Coefficient	P-Value
Maternal age	-0.069	0.491
Gravidity	-0.055	0.587
First minute Apgar score	-0.164	0.118
Fifth minute Apgar score	-0.133	0.207
Weight	-0.030	0.768
Height	-0.074	0.466
Head circumference	-0.010	0.925
PH	-0.064	0.532
Base excess	-0.185	0.068

serum Zn levels and term neonate anthropometric measures, including weight, height, and head circumference.

Gomez et al. (13), in a case-control study, evaluated umbilical cord blood Zn levels in 123 neonates with different gestational ages and birth weights. The results of their study indicated significant correlations between umbilical cord Zn status and neonate birth weight and gestational age. Seriana et al. (35) showed significant relationships between Zn status at term pregnancy and the neonate's height ( $P = 0.026$ ) and head circumference ( $P = 0.012$ ). On the other hand, this significant correlation was not observed with the neonate's birth weight. Nanbakhsh and Tabrizi (16) also evaluated umbilical cord blood Zn levels in pregnant women. According to their findings, mothers with Zn levels  $> 70$   $\mu\text{g/dL}$  had neonates with birth weights  $> 3500$  g, while mothers with Zn levels  $< 60$   $\mu\text{g/dL}$  had neonates with birth weights  $< 3000$  g. Their results also showed that the mean Zn level in low-birth-weight neonates was significantly lower than in their normal-birth-weight counterparts.

These diverse results across studies might relate to differences in sample size, study design, including Zn assessment methods, study population, Zn metabolism, genetic factors, supplementation status, or participants' gestational age. It is suggested that more investigations with larger sample sizes, considering more variables like prenatal supplementation status, maternal underlying diseases, perinatal complications, placental abnormalities, neonates' gestational age, and health status, are needed.

According to the results, cord blood Zn level was not correlated with neonates' sex, Apgar scores, blood pH, or base excess. However, this also underscores the need for further research.

One of the confounding factors in the present study might be related to the use of supplements from

different commercial brands. We assessed the available prenatal supplements in the market and found that the level of zinc was consistent (25 mg Zn). The study results should be interpreted with consideration of its limitations, such as the cross-sectional study design and relatively homogeneous population. Since non-supplemented mothers were not included in the study and the results showed no Zn-deficient mothers, we could not compare the neonates' growth parameters in Zn-sufficient and Zn-deficient mothers.

Case-control studies with larger sample sizes and longitudinal follow-up are recommended to provide more informative insights into post-natal growth and development. Moreover, including multiple gestations and preterm neonates, as well as maternal lifestyle habits (nutritional and smoking status), biomarkers related to Zn metabolism, hematologic disorders like anemia, and placental complications, are factors that may influence the findings. Future investigations including such variables are suggested.

### 5.1. Conclusions

Although adequate Zn levels were observed in all neonates, these levels did not significantly influence neonatal anthropometric measures or immediate health indicators. Further studies with larger sample sizes are recommended to provide more comparative data, particularly regarding long-term growth outcomes.

### Acknowledgements

This study was supported by Tehran University of Medical Sciences (TUMS) and Maternal Fetal and Neonatal Research Center. We acknowledge their kindly supports in this study. The authors would like to thank Dr. Z. K. F. for her kind cooperation.



## Footnotes

**Authors' Contribution:** Conception: M. K., M. R. Z.; Design: M. R. Z., M. K.; Supervision: M. R. Z.; Data collection: M. D., R. S.; Processing: M. R. Z., M. V.; Analysis and interpretation: R. M., M. Sh.; Literature review: M. R. Z., M. K.; Writing: M. R. Z., M. D.; Critical review: M. R. Z., M. K.

**Conflict of Interests Statement:** The authors declared no conflict of interest.

**Data Availability:** Data are available according reasonable request.

**Ethical Approval:** IR.TUMS.IKHC.REC.1400.282

**Funding/Support:** This study was supported by Tehran University of Medical Sciences (TUMS) and Maternal Fetal and Neonatal Research Center.

**Informed Consent:** Written informed consent was obtained from included participants, no extra cost was imposed, and all data were considered confidential.

## References

- Luo J, Wu W, Zhang P, Chen X, Feng Y, Ma N, et al. Zinc Levels and Birth Weight in Pregnant Women with Gestational Diabetes Mellitus: A Matched Cohort Study in China. *J Clin Endocrinol Metab.* 2020;**105**(7). [PubMed ID: 32285111]. <https://doi.org/10.1210/clinem/dgaa171>.
- Chasapis CT, Ntoupa PA, Spiliopoulou CA, Stefanidou ME. Recent aspects of the effects of zinc on human health. *Arch Toxicol.* 2020;**94**(5):1443-60. [PubMed ID: 32394086]. <https://doi.org/10.1007/s00204-020-02702-9>.
- Patil R, Sontakke T, Biradar A, Nalage D. Zinc: an essential trace element for human health and beyond. *Food Health J.* 2023;**5**(3). <https://doi.org/10.53388/fh2023013>.
- Squitti R, Pal A, Picozza M, Avana A, Ventriglia M, Rongioletti MC, et al. Zinc Therapy in Early Alzheimer's Disease: Safety and Potential Therapeutic Efficacy. *Biomolecul J.* 2020;**10**(8). [PubMed ID: 32784855]. [PubMed Central ID: PMC7466035]. <https://doi.org/10.3390/biom10081164>.
- Costa MI, Sarmento-Ribeiro AB, Goncalves AC. Zinc: From Biological Functions to Therapeutic Potential. *Int J Mol Sci.* 2023;**24**(5). [PubMed ID: 36902254]. [PubMed Central ID: PMC10003636]. <https://doi.org/10.3390/ijms24054822>.
- Brion LP, Heyne R, Lair CS. Role of zinc in neonatal growth and brain growth: review and scoping review. *Pediatr Res.* 2021;**89**(7):1627-40. [PubMed ID: 33010794]. <https://doi.org/10.1038/s41390-020-01181-z>.
- Uriu-Adams JY, Keen CL. Zinc and reproduction: effects of zinc deficiency on prenatal and early postnatal development. *Birth Defects Res B Dev Reprod Toxicol.* 2010;**89**(4):313-25. [PubMed ID: 20803691]. <https://doi.org/10.1002/bdrb.20264>.
- DeĠer U, Turan G, Peker N. Is there any connection between zinc deficiency and poor obstetric outcomes in pregnancy? *J Gynecol Obstetr Neonatol Med.* 2022;**19**(3):1424-30. <https://doi.org/10.38136/jgon.1117596>.
- Akdas S, Yazihan N. Cord blood zinc status effects on pregnancy outcomes and its relation with maternal serum zinc levels: a systematic review and meta-analysis. *World J Pediatr.* 2020;**16**(4):366-76. [PubMed ID: 31446568]. <https://doi.org/10.1007/s12519-019-00305-8>.
- Terrin G, Berni Canani R, Di Chiara M, Pietravalle A, Aleandri V, Conte F, et al. Zinc in Early Life: A Key Element in the Fetus and Preterm Neonate. *Nutr J.* 2015;**7**(12):10427-46. [PubMed ID: 26690476]. [PubMed Central ID: PMC4690094]. <https://doi.org/10.3390/nu7125542>.
- Islam MN, Chowdhury MA, Siddika M, Qurishi SB, Bhuiyan MK, Hoque MM, et al. Effect of oral zinc supplementation on the growth of preterm infants. *Indian Pediatr J.* 2010;**47**(10):845-9. [PubMed ID: 20308765]. <https://doi.org/10.1007/s13312-010-0145-8>.
- Laureano A, Bras S, Carvalho R, Amaro C, Cardoso J. Transient symptomatic zinc deficiency in a preterm exclusively breast-fed infant. *Dermatol Online J.* 2014;**20**(2). <https://doi.org/10.5070/d3202021547>.
- Gomez T, Bequer L, Mollineda A, Gonzalez O, Diaz M, Fernandez D. Serum zinc levels of cord blood: relation to birth weight and gestational period. *J Trace Elem Med Biol.* 2015;**30**:180-3. [PubMed ID: 25617232]. <https://doi.org/10.1016/j.jtemb.2014.12.009>.
- Khoushabi F, Shadan MR, Miri A, Sharifi-Rad J. Determination of Maternal Serum Zinc, Iron, Calcium and Magnesium during Pregnancy in Pregnant Women and Umbilical Cord Blood and Their Association with Outcome of Pregnancy. *Mater Sociomed.* 2016;**28**(2):104-7. [PubMed ID: 27147914]. [PubMed Central ID: PMC4851502]. <https://doi.org/10.5455/msm.2016.28.104.107>.
- Atazadegan MA, Heidari-Beni M, Riahi R, Kelishadi R. Association of selenium, zinc and copper concentrations during pregnancy with birth weight: A systematic review and meta-analysis. *J Trace Elem Med Biol.* 2022;**69**:126903. [PubMed ID: 34823102]. <https://doi.org/10.1016/j.jtemb.2021.126903>.
- Nanbakhsh F, Tabrizi FM. The Relationship between Cord Blood and Maternal Serum Zinc Levels and Birth Weight. *Iran J Neonatol.* 2017;**8**(3).
- Samimi M, Asemi Z, Taghizadeh M, Azarbad Z, Rahimi-Foroushani A, Sarahroodi S. Concentrations of Serum Zinc, Hemoglobin and Ferritin among Pregnant Women and their Effects on Birth Outcomes in Kashan, Iran. *Oman Med J.* 2012;**27**(1):40-5. [PubMed ID: 22359724]. [PubMed Central ID: PMC3282140]. <https://doi.org/10.5001/omj.2012.08>.
- Eslami MJ, Khoshhali M, Kelishadi R. A Systematic Review and Meta-analysis on the Prevalence of Zinc Deficiency in Iranian Population. *J Pediatr Rev.* 2023;**11**(3):209-20. <https://doi.org/10.32598/jpr.11.3.451.1>.
- Carducci B, Keats EC, Bhutta ZA. Zinc supplementation for improving pregnancy and infant outcome. *Cochrane Database Syst Rev.* 2021;**3**(3). CD000230. [PubMed ID: 33724446]. [PubMed Central ID: PMC8094617]. <https://doi.org/10.1002/14651858.CD000230.pub6>.
- Ofakunrin A, Collins J, Diala U, Afolaranmi T, Okolo S. Relationship between maternal serum zinc, cord blood zinc and birth weight of term newborn infants in Jos, Plateau State, Nigeria, Jos. *J Med.* 2017;**11**(2):12-20.
- Jyotsna S, Amit A, Kumar A. Study of serum zinc in low birth weight neonates and its relation with maternal zinc. *J Clin Diagn Res.* 2015;**9**(1):SC01-3. [PubMed ID: 25738050]. [PubMed Central ID: PMC4347414]. <https://doi.org/10.7860/JCDR/2015/10449.5402>.
- Daniali SS, Shayegh S, Tajaddini MH, Goodarzi-Khoigani M, Kelishadi R. Association of cord blood zinc level and birth weight in a sample of Iranian neonates. *Int J Prevent Med.* 2020;**11**. [PubMed ID: 32089803]. [https://doi.org/10.4103/ijpvm.IJPVM\\_160\\_19](https://doi.org/10.4103/ijpvm.IJPVM_160_19).
- Amini N, Mousavi S, Vard B, Daniali SS, Kelishadi R. The Assessment of Neonatal Anthropometric Indices Association with Umbilical Cord Blood Zinc and Magnesium Levels. *Adv Biomed Res.* 2023;**12**:253.

- [PubMed ID: 38192895]. [PubMed Central ID: PMC10772790]. [https://doi.org/10.4103/abr.abr\\_90\\_23](https://doi.org/10.4103/abr.abr_90_23).
24. Mohamed S, Kumar S, Kumutha J. A study of zinc level in cord blood of neonates and its relationship with antropometric measurements in the neonates: a descriptive cross-sectional study. *Int J Contemporary Pediatr*. 2019;**6**(2). <https://doi.org/10.18203/2349-3291.ijcp20190718>.
  25. Boskabadi H, Maamouri G, Akhondian J, Ashrafzadeh F, Boskabadi A, Faramarzi R, et al. Comparison of birth weights of neonates of mothers receiving vs. not receiving zinc supplement at pregnancy. *BMC Pregnanc Childbirth*. 2021;**21**(1):187. [PubMed ID: 33676424]. [PubMed Central ID: PMC7936407]. <https://doi.org/10.1186/s12884-021-03598-8>.
  26. Noela OC. The relationship between maternal serum zinc level and cord serum zinc. *Inst Appl Autonom J Biologic Sci*. 2021;**7**(1).
  27. Sairoz Prabhu K, Dastidar R, Aroor AR, Rao M, shetty S, Poojari VG, et al. Micronutrients in Adverse Pregnancy Outcomes. *F1000Research*. 2024;**11**. <https://doi.org/10.12688/f1000research.124960.4>.
  28. Karim S, Fazal S, Naeem M, Ali H, Fazal R, Karim A, et al. Frequency of low birth weight and its relationship with maternal serum zinc: a cross-sectional study. *Ann Med Surg (Lond)*. 2023;**85**(6):2469-73. [PubMed ID: 37363444]. [PubMed Central ID: PMC10289759]. <https://doi.org/10.1097/MS9.0000000000000687>.
  29. Thakur GK, Shankar H, Arora TK, Kulkarni B. Role of mineral nutrients other than iron in pregnancy: under recognized opportunities to improve maternal/fetal outcomes: a literature review. *Arch Gynecol Obstet*. 2024;**309**(3):895-905. [PubMed ID: 37698602]. <https://doi.org/10.1007/s00404-023-07183-6>.
  30. Shahraki Jazinaki M, Rashidmayvan M, Safarian M, Norouzy A. Investigating the Effects of Zinc Supplementation on Growth-Related Factors in Infants With Failure to Thrive: A Systematic Review and Meta-analysis of Randomized Controlled Trials. *J Pediatr Rev*. 2024;**12**(1):15-26. <https://doi.org/10.32598/jpr.12.1.1150.1>.
  31. Gonzalez-Fernandez D, Muralidharan O, Neves PA, Bhutta ZA. Associations of Maternal Nutritional Status and Supplementation with Fetal, Newborn, and Infant Outcomes in Low-Income and Middle-Income Settings: An Overview of Reviews. *Nutr J*. 2024;**16**(21). [PubMed ID: 39519557]. [PubMed Central ID: PMC11547697]. <https://doi.org/10.3390/nu16213725>.
  32. Keats EC, Oh C, Chau T, Khalifa DS, Imdad A, Bhutta ZA. Effects of vitamin and mineral supplementation during pregnancy on maternal, birth, child health and development outcomes in low- and middle-income countries: A systematic review. *Campbell Syst Rev*. 2021;**17**(2). e1127. [PubMed ID: 37051178]. [PubMed Central ID: PMC8356361]. <https://doi.org/10.1002/cl2.1127>.
  33. Syakur R, Syam A, Hadju V, Palutturi S. The impact of maternal zinc supplementation on pregnancy outcomes and maternal serum zinc: A systematic review. *Nveo-Natur Volatil Essent Oils J*. 2021;**2021**:8301-13.
  34. Bayomy A, Kassem YT, Younis M, Solaiman AM. Cord Serum Zinc Level and Its Impact on Birth Weight, Length and Growth Velocity in the First Three Months of Life. *America J Med Med Sci*. 2017;**7**(2):47-54.
  35. Seriana I, Yusrawati Y, Lubis G. Serum zinc level at term pregnancy and newborn anthropometry. *Indonesia J Obstetr Gynecol*. 2015;**2015**:190-5.