



# Risk of Attention Deficit Hyper Activity Disorder After Early Exposure to General Anesthesia; A Case Control Study

Abbas Sedighnejad<sup>1</sup>, Soheil Soltanipour<sup>2</sup>, Alia Saberi<sup>2</sup>, Maryam Kousha<sup>2</sup>, Elham Bidabadi<sup>2</sup>, Gelareh Biazar<sup>1,\*</sup> and Novin Naderi<sup>2</sup>

<sup>1</sup>Anesthesiology Department, Anesthesiology Research Center, Alzahra Hospital, Guilan University of Medical Sciences, Rasht, Iran

<sup>2</sup>Guilan University of Medical Sciences (GUMS), Rasht, Iran

\*Corresponding author: Anesthesiology Department, Anesthesiology Research Center, Alzahra Hospital, Guilan University of Medical Sciences, Rasht, Iran. Email: gelarehbiazar1386@gmail.com

Received 2019 December 06; Revised 2020 March 29; Accepted 2020 March 31.

## Abstract

**Background:** Over the past decade, following the discovery that developing brain of immature animals was affected by anesthetic agents, the safety of general anesthesia (GA) in early life has been questioned.

**Objectives:** We investigated the association between anesthesia exposure in children and ADHD development.

**Methods:** This case-control study was conducted at pediatric psychology clinic of our institution and a pediatric neurology private clinic during 2019. Firstly the responsible resident of anesthesiology separated new ADHD cases. Then a questionnaire was filled out through an almost 10 minute's telephone interview. Finally, frequency distribution of GA was compared between ADHD cases and controls.

**Results:** Finally, the data from 210 children were analyzed. Among 105 ADHD cases, 19% had a history of a procedure requiring GA while it was 3.8% in control group. Comparing the two groups a significant difference was observed regarding the age of receiving GA ( $P = 0.004$ ), gender ( $P < 0.001$ ), the history of receiving GA ( $P = 0.001$ ) and the number of anesthesia exposures ( $P = 0.001$ ). According to logistic regression analysis, male gender ( $P = 0.001$ ) OR 3.11 (95CI = 1.63 - 5.93) and age ( $P = 0.003$ ) OR 0.92 (95CI = 0.87 - 0.97) were significant predictors of early exposure to GA and ADHD development.

**Conclusions:** It was revealed that early exposure to GA might be a risk factor for later developing ADHD. Boys might be more sensitive to the long term adverse effects of anesthetic agents than girls. Further prospective well-planned studies are needed to confirm these findings.

**Keywords:** Attention Deficit Hyper Activity Disorder, General Anesthesia, Early Exposure

## 1. Background

Alone in the United States annually six million children undergo surgeries under GA while 1.5 million of them are infants. Anesthesia benefits in pediatric surgeries including maintaining stable hemodynamic state, reduction of pain and anxiety, providing proper conditions for surgeon are not deniable. However recently the safety of GA in young children has been questioned (1, 2). Experimental studies have shown that early exposure of developing brain to general anesthesia results in neurodegenerative changes (3). The current available anesthetic drugs which act as N-methyl-D-aspartic acid receptor antagonists (NMDA) and  $\gamma$ -amino butyric acid modulator interfere with CNS development. Since all anesthetic drugs except of opioids and agonists act as above, GA and deep sedation can result in apoptotic neuro-degeneration. These

agents affect primarily cortical regions through apoptotic phenomena (4). These changes cause deficits in different aspects of behavior. Indeed the findings of animal researches induce the concern that the mentioned risk might be also in human brain. Up to now human studies have discussed the risk of neurotoxicity related GA with controversial results (5). In spite of a number of researches with different outcome measures, (e.g. intelligence, academic achievements neuro-psychological statue biomarkers and neuro-imaging) that have assessed the GA harm in developing brain, there are a lot of gaps in our knowledge (6). Due to the lack of an agreement on the topic, whether GA causes neurodevelopment impairment, investigations continue to reach a definite answer (7). The concern regarding the risks of anesthesia agents is not restricted just to anesthesiologists' society, it has involved other fields, United States Food and Drug Administration (FDA), the Eu-

ropean Medicines Agency and even partly public society (8). A wide range of time from pregnancy up to 4 years have been considered as unsafe age for GA (9). In an experimental research, Fredriksson and Archer (10) examined the effects of induction of general anesthesia with ketamine on rodents and found that they developed hyperactivity which responded well to dextroamphetamine.

Although translation of the clinical significance of these data to human brain is difficult, it was supposed that this pattern may mimic human attention-deficit/hyperactivity disorder (ADHD) in children (11). ADHD as the most common neuropsychological disease of children may continue into adulthood. The child suffers from uncontrolled impulses, motor restlessness and impaired ability to pay attention and concentrate (12).

The predisposing factors for this disease are not well known yet. However, available information strongly points to the important role of both gene and environment for clinical manifestation of ADHD (13). Due to the scarcely limited researches in our country and the importance of the issue, this study was planned.

## 2. Objectives

In the present study we investigated whether GA exposure before four years of age was associated with behavioral problems or not.

## 3. Methods

This case control study was conducted at pediatric psychology clinic of our institution and a pediatric neurology private clinic, from February to July 2019. Firstly the responsible resident of anesthesiology screened all the files of children who were referred to the mentioned centers and ADHD cases that were recently diagnosed were identified and sorted out.

Inclusion criteria consisted of newly diagnosed ADHD cases (within the last year) having a healthy brother or sister, whose parents were contactable and also accepted to enroll in the survey. An experienced pediatric psychiatrist and a pediatric neurologist identified our cases, according to the DSM-5 diagnostic criteria for ADHD.

### 3.1. Exclusion Criteria (Families with No Cooperation)

After sorting out the files, the resident of anesthesiology telephoned the families of these children and explained the aim of the survey. When they agreed to participate, a questionnaire was filled out through an almost 10 minute's telephone interview for both ADHD cases and the healthy sister or brother. The parents were asked to answer

the questions including the children's history of exposure to GA in the first four years of age, the age of exposure, birth status; term or preterm and single or multi-exposure to GA. The main outcome of the study is development of ADHD when exposed to general anesthesia in childhood.

Finally the frequency distribution of GA exposure was compared between ADHD cases and controls and the data were compared between healthy and ADHD groups to find any association between receiving GA in the first four years of age and ADHD.

### 3.2. Sample Size

Based on a pilot study we found that 104 cases could be a proper sample size for this survey. From 30 ADHD cases, six (20%) had a history of general anesthesia exposure in the first four years of age. Study power was considered as 90%,  $Z = 1.96$  and  $\beta = 1.28$ . The used formula is presented below.

### 3.3. Ethics

After approval of the study protocol by the Ethics in Research Committee of the University (no.: IR.GUMS.REC.1397.524), informed consent was obtained from parents.

## 4. Results

The data from 210 children were analyzed. Among 105 ADHD cases, 19% had a history of procedures requiring general anesthesia while it was 3.8% in control group. The characteristics of cases and controls are presented in [Table 1](#). Comparing the two groups a significant difference was observed regarding the age of receiving GA ( $P = 0.004$ ), gender ( $P < 0.001$ ), the history of receiving GA ( $P = 0.001$ ) and the number of anesthesia exposures ( $P = 0.001$ ). Hence in order to adjust the variables with significant difference in univariate analysis, (sex, history of anesthesia exposure, number of anesthesia exposures and the age when received anesthesia), backward likelihood ratio method was applied and after three steps, age, sex remained statistically significant. According to logistic regression analysis, male gender ( $P = 0.003$ ) OR 0.92 (95CI = 0.87 - 0.97) and age ( $P = 0.001$ ) OR 3.11 (95CI = 1.63 - 5.93) were significant predictors of early exposure to GA and ADHD development ([Table 2](#)).

Statistics analysis: Statistical package for social science (SPSS) version 16 software was used to analyze the data. According to Kolmogorov-Smirnov test the age distribution was not normal among cases and controls. Mann-Whitney U and chi-square tests were used. In multivariate analysis,

**Table 1.** Characteristics of Cases and Controls<sup>a</sup>

	Case	Control	Value	P value
Age, y	11 (4)	15 (11)	4114.50 <sup>b</sup>	0.001
Sex, %				< 0.001
Male	79	54.3	14.48 <sup>c</sup>	
Female	21	45.7		
Birth weight, %				0.685
Normal	85.7	87.6	0.16 <sup>c</sup>	
Low birth weight	14.3	12.4		
Situation at birth, %				0.390
Term	92.4	95.2	0.73 <sup>c</sup>	
Preterm	7.6	4.8		
History of anesthesia exposure, %				0.001
No	81	96.2	11.98 <sup>d</sup>	
Yes	19	3.8		
Number of anesthesia exposures, %				0.001
Zero	81	96.2	12.38 <sup>d</sup>	
Once	15.2	3.8		
> Once	3.8	0		
Age when receiving anesthesia, %				P = 0.004
One ye	5.7	1.9	13.16 <sup>d</sup>	
Two years	5.7	0		
Three years	4.8	1.9		
> Three years	2.9	0		

<sup>a</sup>Values are expressed as No. (%) or median (interquartile range).<sup>b</sup>Mann-Whitney U.<sup>c</sup>Pearson chi-square.<sup>d</sup>Fisher's exact.**Table 2.** Odds Ratio and Confidence Interval of Predictors for ADHD Development<sup>a</sup>

Step 3	Values, B (SE)	95% CI <sup>b</sup> for Exp b		
		Lower	Exp b	Upper
Included				
Constant	0.03 (0.41)			
Age	-0.07 (0.02) <sup>c</sup>	0.87	0.92	0.97
Sex	1.13 (0.32) <sup>d</sup>	1.63	3.11	5.93

<sup>a</sup>R<sup>2</sup> = 0.17 (Cox & Snell), 0.22 (Nagelkerke). Model  $\chi^2$  (5) = 39.11, P < 0.001.<sup>b</sup>Confidence interval.<sup>c</sup>P = 0.003.<sup>d</sup>P = 0.001.

we used logistic regression analysis for predicting our dichotomous outcome (ADHD). Statistical significance was considered as P < 0.05.

## 5. Discussion

Neurodevelopment abnormalities in ADHD cases have been described well. Studies have found that in these cases, hypofunction of N-methyl d-aspartate receptors induce inattention. Prefrontal cortex (PFC) is responsible for thoughts, analysis and regulating behavioral, emotion focus and attention. PFC helps to predict the outcomes of a behavior and determining right from wrong. This vital part of the brain is unregulated in ADHD cases and dendritic spine density in PFC significantly change (14). According to our search, it was the first study in Iran evaluating the association between early GA exposure and later behavioral disorders. Indeed, increasing interest in this topic as a big concern judged by numerous published articles, has not been observed in our country and the limited available studies indicate the lack of enough attention to the issue (15, 16). The main finding of this work was that chil-

dren exposed to anesthetic agents in the first four years of age, had a higher incidence of ADHD than those without this history. In other words we supported previous studies which indicated the neurotoxicity of anesthetic drugs in developing brain. We found that the age of receiving GA, male gender, the history of receiving GA and the number of exposures were significantly associated with ADHD. However they could not be a strong predictor for this behavioral disorder. After our data was stratified by sex, we found a strong association between ADHD and male gender. Searching the current literature, some of them supported our findings and some other were in contrast. In line with our paper, Tsai et al. (17) in a birth cohort study reported that exposure to GA before the age of three years had an increased risk for later ADHD. DiMaggio et al. (18) in two retrospective studies found that exposure to GA in the first 3 years of age increased the risk of developmental or behavioral abnormalities. Ing et al. (19) in 2012 indicated that children who were exposed to GA in the first three years of age showed more language deficits than unexposed ones. Furthermore studies reported that neonates delivered by cesarean section under GA were more likely to develop behavioral deficits compared with those delivered vaginally without anesthesia (20). Flick et al. (21) 2011 demonstrated that early exposure to GA could be an independent risk for neurological disorders affecting both learning and behavior abilities. In contrast to our findings, in a pilot study Kalkman et al. (22) reported that there was a non-significant association between exposed children and non-exposed to GA before 24 months regarding behavioral disorders. Bartels et al. (23) did not report the mentioned association either. O'leary J et al. (24), reported that children who received GA before age 5 to 6 were at a higher risk of early neurodevelopment vulnerability and long term adverse outcomes. However they found that multiple exposure or age under 2 were not recognized as additional risks. Sprung et al. (11) studied the association between GA before 2 years of age and the development of ADHD. They found that children, who underwent repeated surgeries under GA, had a higher risk of development of ADHD. Opposite to this work, Ko et al. (25) in a retrospective matched-cohort study in Taiwan reported that there was no association between early life anesthesia exposure before three years of age and ADHD. Creagh et al. (26) also did not observe any positive correlation. Bong et al. (27) in a retrospective study found that the incidence of learning disability among children with a history of GA exposure before one year was 4.5 times greater than that of not exposed peers. As discussed above, a discrepancy among the findings of human studies is observed which could be justified by the differences regarding socioeconomic status, genetics, familial conditions, parenteral characteris-

tics such as age, comorbidities, dosage and timing of GA, all of which might affect the results. Indeed the reason of this disagreement could be differences in study design such as choice of study population, sample size, the length of follow-ups, different assessment tools (e.g. intelligence, academic success, behavioral disorders) (28). The definition of ADHD and case selection strategies might be different among studies. In Sprung et al.'s study (11) the majority of cases came from schools, that had referred the children for behavioral problems and a questionnaire was filled out by teacher or parent. Tasi et al. (17) used a nationwide population-based sample and in our research we had a regional participation. Furthermore studies which select cases based on ICD-9-CM code 314.01 that presents a combined type of disease, may miss ADHD cases who clinically express hyperactivity or inattention, not both of them. In Tasi et al.'s study (17) with ICD-9-CM 314 a broader criterion was considered for ADHD diagnosis. Furthermore; selected exposure period was not the same among studies. Sprung considered this time before the age of two years (11). Tasi et al. (17) before 3 and in our study it was extended to before 4. The other noticeable factor was duration of follow-up periods. In Ko et al. (25) study children aged 5 - 10 years were focused on. Therefore cases diagnosed after 10 years of age could be missed. Due to different interpretations among observational studies and the multifactorial nature of the mentioned criteria, focusing on other modalities such as biomarkers and neuroimaging might provide more reliable results. There are still several unanswered questions: anesthetic drugs, doses, anesthesia duration, age at exposure and proper evaluation criteria. We acknowledge that to achieve more meaningful results, cohort studies with an adequate sample size is required. Surely, neuro behavioral disorders are multifactorial and similar to other supporting studies we cannot claim that we have found a single causative factor. However, despite the inconstant results of clinical studies and unanswered questions in this field, based on accumulating evidence suggesting irreversible neuronal damage and lasting neurodevelopmental sequels, it is wise to avoid any unnecessary procedure requiring GA in early life. Obviously children's deprivation of anesthesia and analgesia is not legally or ethically accepted (29). Definitely to achieve the desired goals, not only anesthesiologists but also other specialists should be aware from the potential risks of GA administration during early life (15). Indeed proper communication with other involved physicians who refer the children for an elective surgery or invasive diagnostic procedure requiring GA which could be postponed is crucial (30). Providing sufficient knowledge in general society especially among parents should be considered as well. As such parents frequently question the physicians about the

safety of GA in their children (31).

### 5.1. Limitations

We admit that there are several limitations for this work. Indeed due to the nature of this study, our data was achieved via a telephone interview and in many cases parents did not have a proper communication or might not remember the required data.

Therefore we were not able to infer causality either and clarify the pure effects of GA because it was hard to differentiate how the potential confounding factors such as maternal smoking, alcoholism, mental statue, child nutrition, lead exposure, child hood systemic and inflammatory diseases could affect the results.

### 5.2. Conclusions

This study showed that early exposure to GA might be a risk factor for later developing ADHD. Boys might be more sensitive to the long term adverse effects of anesthetic agents than girls. However, we believe that to confirm what is reported here and to determine the possible mechanisms for this association, future studies are required.

## Acknowledgments

Authors would like to thank all the children's families for their kind collaboration.

## Footnotes

**Authors' Contribution:** Abbas Sedighnejad contributed in study design and quality assurance. Gelareh Biazar contributed in manuscript writing and critical revision. Soheil Soltanipour performed data analysis and peerred the paper. Alia Saberi helped in study design and peerred the manuscript. Maryam Kousha, Elham Bidabadi, and Novin Naderi contributed in data collection.

**Conflict of Interests:** No conflict of interest.

**Ethical Approval:** The Institutional Review Board (IRB) of Guilan University of Medical Sciences approved study protocol.

**Funding/Support:** This study was not supported.

**Informed Consent:** Informed consent was obtained from parents.

## References

1. Sun L. Early childhood general anaesthesia exposure and neurocognitive development. *British journal of anaesthesia*. 2010;**105**(suppl\_1):i61-8.
2. Barton K, Nickerson JP, Higgins T, Williams RK. Pediatric anesthesia and neurotoxicity: what the radiologist needs to know. *Pediatric radiology*. 2018;**48**(1):31-6.
3. Walters JL, Paule MG. Developmental Neurotoxicity of General Anesthetics. *Handbook of Developmental Neurotoxicology*. USA: Elsevier; 2018. p. 477-85.
4. Sun LS, Li G, Miller TL, Salorio C, Byrne MW, Bellinger DC, et al. Association between a single general anesthesia exposure before age 36 months and neurocognitive outcomes in later childhood. *Jama*. 2016;**315**(21):2312-20.
5. O'leary JD, Janus M, Duku E, Wijeyesundera DN, To T, Li P, et al. Influence of surgical procedures and general anesthesia on child development before primary school entry among matched sibling pairs. *JAMA pediatrics*. 2019;**173**(1):29-36.
6. Zanghi CN, Jevtovic-Todorovic V. A holistic approach to anesthesia-induced neurotoxicity and its implications for future mechanistic studies. *Neurotoxicology and teratology*. 2017;**60**:24-32.
7. Clausen NG, Kähler S, Hansen TG. Systematic review of the neurocognitive outcomes used in studies of paediatric anaesthesia neurotoxicity. *British Journal of Anaesthesia*. 2018;**120**(6):1255-73.
8. US Food, Drug Administration. *FDA Drug Safety Communication: FDA review results in new warnings about using general anesthetics and sedation drugs in young children and pregnant women*. FDA; 2017.
9. Disma N, Mondardini MC, Terrando N, Absalom AR, Bilotta F. A systematic review of methodology applied during preclinical anesthetic neurotoxicity studies: important issues and lessons relevant to the design of future clinical research. *Pediatric Anesthesia*. 2016;**26**(1):6-36.
10. Fredriksson A, Archer T. Hyperactivity following postnatal NMDA antagonist treatment: reversal by D-amphetamine. *Neurotoxicity research*. 2003;**5**(7):549-64.
11. Sprung J, Flick RP, Katusic SK, Colligan RC, Barbaresi WJ, Bojanić K, et al. Attention-deficit/hyperactivity disorder after early exposure to procedures requiring general anesthesia. *Mayo Clinic Proceedings*. Elsevier; 2012. p. 120-9.
12. Nazari MA TJMSH. Do Computer Games Affect Arousal Level in Children With Attention/Deficit Hyperactivity Disorder? *Caspian Journal of Neurological Sciences*. 2018;**4**(4). doi: 10.29252/CJNS.4.15.144.
13. Luo Y, Weibman D, Halperin J, Li X. A Review of Heterogeneity in Attention Deficit/Hyperactivity Disorder (ADHD). *Frontiers in human neuroscience*. 2019;**13**:42.
14. Depue BE, Burgess GC, Willcutt EG, Ruzic L, Banich MT. Inhibitory control of memory retrieval and motor processing associated with the right lateral prefrontal cortex: evidence from deficits in individuals with ADHD. *Neuropsychologia*. 2010;**48**(13):3909-17.
15. Sedighinejad A, Soltanipour S, Rimaz S, Biazar G, Chaibakhsh Y, Kouhi MB. General Anesthesia-Related Neurotoxicity in the Developing Brain and Current Knowledge and Practice of Physicians at Guilan Academic Hospitals. *Anesthesiology and Pain Medicine*. 2019;**9**(4).
16. Nabi B, Lima S. General Anesthesia-Related Neurotoxicity: Status of Pediatric Surgeries at an Academic Hospital in the North of Iran. *Journal of Comprehensive Pediatrics*. 2019;**10**(4).
17. Tsai C, Lee CT, Liang SH, Tsai P, Chen VC, Gossop M. Risk of ADHD after multiple exposures to general anesthesia: a nationwide retrospective cohort study. *Journal of attention disorders*. 2018;**22**(3):229-39.
18. DiMaggio C, Sun L, Li G. Early childhood exposure to anesthesia and risk of developmental and behavioral disorders in a sibling birth cohort. *Anesthesia and analgesia*. 2011;**113**(5):1143.
19. Ing C, DiMaggio C, Whitehouse A, Hegarty MK, Brady J, von Ungern-Sternberg BS, et al. Long-term differences in language and cognitive function after childhood exposure to anesthesia. *Pediatrics-English Edition*. 2012;**130**(3). e476.
20. Chien L, Lin H, Shao YJ, Chiou S, Chiou H. Risk of autism associated with general anesthesia during cesarean delivery: a population-based birth-cohort analysis. *Journal of autism and developmental disorders*. 2015;**45**(4):932-42.

21. Flick RP, Katusic SK, Colligan RC, Wilder RT, Voigt RG, Olson MD, et al. Cognitive and behavioral outcomes after early exposure to anesthesia and surgery. *Pediatrics*. 2011;**128**(5). e1053.
22. Kalkman CJ, Peelen L, Moons KG, Veenhuizen M, Bruens M, Sinnema G, et al. Behavior and development in children and age at the time of first anesthetic exposure. *Anesthesiology: The Journal of the American Society of Anesthesiologists*. 2009;**110**(4):805-12.
23. Bartels M, Althoff RR, Boomsma DI. Anesthesia and cognitive performance in children: no evidence for a causal relationship. *Twin research and human genetics*. 2009;**12**(3):246-53.
24. O'leary JD, Janus M, Duku E, Wijeyesundera DN, To T, Li P, et al. A population-based study evaluating the association between surgery in early life and child development at primary school entry. *Anesthesiology: The Journal of the American Society of Anesthesiologists*. 2016;**125**(2):272-9.
25. Ko W, Liaw Y, Huang J, Zhao D, Chang H, Ko P, et al. Exposure to general anesthesia in early life and the risk of attention deficit/hyperactivity disorder development: a nationwide, retrospective matched-cohort study. *Pediatric Anesthesia*. 2014;**24**(7):741-8.
26. Creagh O, Torres H, Rivera K, Morales-Franqui M, Altieri-Acevedo G, Warner D. Previous Exposure to Anesthesia and Autism Spectrum Disorder (ASD): A Puerto Rican Population-Based Sibling Cohort Study. *Boletín de la Asociación Médica de Puerto Rico*. 2015;**107**(3):29-37.
27. Bong CL, Allen JC, Kim JTS. The effects of exposure to general anesthesia in infancy on academic performance at age 12. *Anesthesia & Analgesia*. 2013;**117**(6):1419-28.
28. Castellheim A, Lundström S, Molin M, Kuja-Halkola R, Gillberg C, Gillberg C. The role of general anesthesia on traits of neurodevelopmental disorders in a Swedish cohort of twins. *Journal of Child Psychology and Psychiatry*. 2018;**59**(9):966-72.
29. Wang X, Xu Z, Miao C. Current clinical evidence on the effect of general anesthesia on neurodevelopment in children: an updated systematic review with meta-regression. *PloS one*. 2014;**9**(1). e85760.
30. Gart MS, Suresh S, Adkinson JM. Anesthetic neurotoxicity in congenital hand surgery: an overview of the evidence and advice for counseling parents. *The Journal of hand surgery*. 2017;**42**(7):564-8.
31. Nemergut ME, Aganga D, Flick RP. Anesthetic neurotoxicity: what to tell the parents? *Pediatric Anesthesia*. 2014;**24**(1):120-6.