

# Factors Influencing Object Relation Therapy Based on Transference and Potential Space: Analyzing the Experience and Understanding of Clients Suffering from Major Depressive Disorder

Samaneh Ghafoori,<sup>1,2</sup> Mohammad-Kazem Atef-Vahid,<sup>1,2,\*</sup> Frank Summers,<sup>3</sup> Mahmood Dehghani,<sup>1</sup>

Farahnaz Mohammadi Shahboulaghi,<sup>4</sup> and Aliasghar Asgharnejad Farid<sup>1,2</sup>

<sup>1</sup>School of Behavioral Sciences and Mental Health (Tehran Institute of Psychiatry), Iran University of Medical Sciences, Tehran, Iran

<sup>2</sup>The Center of Excellence in Psychiatry and Clinical Psychology, Iran University of Medical Sciences, Tehran, Iran

<sup>3</sup>Feinberg School of Medicine, Northwestern University Chicago, IL, P

<sup>4</sup>Social Determinants of Health Research Center, Nursing Department, University of Social Welfare and Rehabilitation Sciences, Tehran, Iran

\*Corresponding author: Mohammad-Kazem Atef-Vahid, School of Behavioral Sciences and Mental Health (Tehran Institute of Psychiatry) and the Center of Excellence in Psychiatry and Clinical Psychology, Iran University of Medical Sciences, Tehran, Iran. Tel: +98-9123459593, E-mail: kazemv@yahoo.com

Received 2017 January 05; Revised 2017 March 16; Accepted 2017 April 07.

## Abstract

**Background:** Depression is a widespread disorder that affects all aspects of a person's life. Various psychological and medical interventions have been proposed to help treat depressive symptoms. One effective treatment is object relation therapy based on transference and potential space.

**Objectives:** In the current study, we focus on how clients who suffer depressive disorder understand this type of therapy.

**Methods:** This qualitative research was conducted in 2015 - 2016 using a content analysis with 18 Iranian participants who were diagnosed with depressive disorder based on the structured clinical interview for DSM-IV, the clinician version (SCID-I/CV) and who had received object relation therapy based on transference and potential space. The study was carried out in an outpatient clinic of the school of behavioral sciences and mental health (formerly Tehran Institute of psychiatry). The data were collected through conducting in-depth semi-structured interviews, which continued until data saturation was achieved. The data were analyzed with the induction method.

**Results:** The analysis of the data resulted in the identification of 2 main categories and 4 subcategories. "Gradual formation of therapeutic alliance" and "use of non-interpretative techniques" were the subcategories of "analytic dyad formation" as the main category; and "in-depth examination of client experience" and "client's emotional expression" were listed under "development of self-awareness and self-expression".

**Conclusions:** Gradual formation of therapeutic alliance is consistent with literature that emphasizes the necessity of new relational experiences and creation of a safe attachment base in therapy. The findings showed that the exploration and awareness of patterns in the context of a secure relationship and transference and extra-transference can produce therapeutic changes. In addition, such safe space can provide a suitable context for releasing built-up emotions.

**Keywords:** Depressive Disorder, Object Relation, Qualitative Research, Transference

## 1. Background

Depression is one of the most common mental disorders and is the fourth major cause of disability in the world (1). A variety of psychological therapies have been proposed. One of the psychoanalytic therapies is object relation therapy.

In various object relation approaches, depression is conceptualized in different ways such as reactivation of depressive position (2). In one recent approach of object relation, depression has been regarded as a kind of defect in self, which arises from the lack on necessary appropriate to create an authentic self rooted in emotions, or in other words, what self needs to be lively (3, 4).

In this approach, Summers (4-7), based on the object relation model, emphasizes on 2 critical phases in his therapy. The first one is the interpretation phase, which puts an emphasis on the client's defense mechanisms, client-analyst relation, transference, the client's primary attachment, internalization of these relations, and insight. In the second phase, an opportunity is provided for new patterns to be created and for arrested potentials to be developed using the potential space. He believes that, in addition to interpretation, the analytic space should provide a chance for new patterns to be created and replace the current pathological ones. Potential space does not bear a meaning and any meaning in this space is potential and

future-oriented. It emphasizes on self creation by client.

Understanding the nature of therapy the clients suffering from depressive disorder can play a vital role in the facilitation of therapy and boosting its effectiveness. Whilst the current theory tries to shed light on the nature of therapy for clients suffering from depressive disorder, there is still ambiguity about factors influencing this kind of therapy, which is exacerbated by the lack of evidence on how clients perceive this therapy.

## 2. Objectives

Reviewing recent quantitative studies (e.g. (8, 9)) regarding the effect impact of psychoanalysis on depression reveals the correlation between the quantitative approach and pre-determined mind frames. Since such positivistic paradigm lacks the capability to offer an understanding of the nature of therapy and does not consider this on the basis of client's understanding, the present study was conducted in order to understand factors which affect this type of therapy.

## 3. Materials and Methods

This study is a qualitative research, which was carried out using the content analysis method proposed by Graneheim and Lundman (10).

### 3.1. Participants

The participants in this study included 18 clients who were purposefully selected from clients who were diagnosed with depressive disorder SCID-I/CV, had received object relation therapy based on transference and potential space on a weekly basis for at least one year, and were willing to participate in the study.

Data collection continued until it was determined that data saturation has been reached. The demographic characteristics of participants are shown in [Table 1](#).

### 3.2. Procedure

Data were collected through in-depth semi-structured interview. Subjects were interviewed at the clinic of the school of behavioral sciences and mental health (formerly Tehran Institute of psychiatry) and their workplace. Interviews were recorded and transcribed. Each interview session lasted from 1 to 2.5 hours in once or twice.

To adhere to ethical principles, approval of the ethics committee of Iran's University of Medical Sciences was obtained. In addition, the participants were informed of the goals and methodology of the study, they were informed that the sessions will be audio-recorded, and were also

asked to fill out and sign a written consent. They were also informed of their right to withdraw from the study at any time during the course of the study.

### 3.3. Data Analysis

The method consists of 4 stages: choosing a unit of analysis, detecting the meaning units and referring to a key word, the process of shortening while preserving the core, and description and interpretation on a higher logical level and creation of categories.

To prepare the data for analysis, each transcript was read several times in order to identify its meaning units. Then, they were condensed with a close description of data. After that, a list of codes was developed and reviewed from a semantic point of view. Afterwards, by specifying the differences and similarities in meanings, common codes were determined on a more abstract level using the method of reduction and induction. Data analysis was performed by the research team and all the codes and categories were approved during consecutive sessions.

For the purpose of trustworthiness, member check, peer check, and external checks were done. In order to understand the participants' statements better, findings were presented to them so that fitness of data labeling to participants' experience was approved (member check). In addition, some of the writings were reviewed by those faculty members who were familiar with the qualitative research method (peer check). Dependability was achieved through audit trail by 2 experts who supervised data documentation and analysis. Effort was also put in achieving data credibility through sampling people with highest variation, different demographic classes, and people who received therapy for different lengths of time.

## 4. Results

Data analysis showed that influential factors are 2 main categories of "formation of analytic dyad" and "development of self-expression and self-awareness" as well as 4 subcategories ([Table 2](#)).

### 4.1. Formation of Analytic Dyad

This category refers to the creation of therapeutic relationship. The subcategories are "gradual formation of therapeutic alliance" and "use of non-interpretive techniques".

#### 4.1.1. Gradual Formation of Therapeutic Alliance

Formation of analytic dyad is one of the influential factors. Providing a safe space by the therapist, holding and

**Table 1.** Characteristics of Participants

Code	Gender	Age	Marital Status	Education	Duration of Therapy, y
1	M	40	Single	Ph.D	4.5
2	F	32	Single	Post-doctorate	2
3	F	34	Married	Graduate	3.5
4	M	32	Married	Post-graduate	4
5	M	39	Married	Graduate	9
6	F	34	Married	Ph.D	7
7	F	34	Married	Graduate	4
8	F	32	Married	Ph.D	2.5
9	F	29	Single	Post-graduate	3
10	F	32	Single	Post-graduate	1
11	F	26	Single	Post-graduate	2.5
12	F	39	Single	Ph.D	5
13	M	25	Single	Ph.D	3
14	F	35	Married	Graduate	3
15	F	38	Married	Post-graduate	1.5
16	M	28	Single	Graduate	3.5
17	F	36	Divorced	Post-graduate	3
18	F	30	Single	Graduate	1

**Table 2.** Emerged Categories and Subcategories Related to Content Analysis of Therapeutic Process

Main Categories	Subcategories
<b>Formation of analytic dyad</b>	1. Gradual formation of therapeutic alliance 2. Use of non-interpretive techniques
<b>Development of self-awareness and self-expression</b>	1. In-depth examination of client's experience 2. Client's emotional expression

containing the client, optimal responsiveness of the therapist towards the client's emotional needs, and managing the time of sessions by the therapist affect the formation of analytic dyad.

The safe space, in turn, is created by several actions of the therapist, which are confidentiality and non-judgmental behavior of the therapist, careful listening and remembering by the therapist, paying attention to the client's readiness for presenting crucial topics, and considering the client's mental capacity.

For clients, containing and holding the client by the therapist when are needed, helped to stay away from unnecessary confrontation with intense and unbearable emotions until the time was ripe.

This optimal responsiveness was based on the client's frustration and satisfaction and put the therapist in a situ-

ation where he or she would be good enough for the client, however it never ended in satisfaction.

The clients' psychological needs include the need to be accepted and respected, acceptance of all of his or her painful feelings and emotions and their talks, the therapist's empathy with the client and viewing the world from the his or her point of view, the therapist's availability, and attunement with the client's needs and conditions.

Managing the time included starting and finishing sessions on time, integrating appropriate time intervals between clients, and not bringing up critical issues near the end of the sessions, which conveyed the sense of respect and punctuality of the therapist and provided a stable and safe model, which in turn increased predictability and gave the client a sense of security.

#### 4.1.2. Use of Non-Interpretive Techniques

These techniques included the therapist's supportive interventions, emphatic confrontation by gradual presentation of questions related to conflicting words, thoughts, or feelings without causing guilt feelings in the client or attacking him or her, and avoiding sudden confrontation; direct intervention of the therapist in necessary conditions in the sense of inhibiting the client from acting out their feelings; the therapist's reflective functioning in the sense of returning what the client has brought to the session, especially reflecting the therapist's vision of the client's potentials and capacities.

The therapist's supportive interventions included joining with the client, reassurance, normalization of thoughts, feelings, and fantasies.

#### 4.2. Development of Self-Awareness and Self-Expression

For clients, the prominent factor in therapy is the development of self-awareness and self-expression. In-depth exploration of the clients' experiences and expression of feelings falls under this category.

##### 4.2.1. In-Depth Exploration of the Client's Experiences

In this subcategory, all feelings, thoughts, behaviors, and fantasies of the client in the analytic dyad are examined. This subcategory includes the therapist's neutral stance, in-depth and insightful inquiries, interpretations, and addressing client's free associations.

Having no family relation with the therapist, not taking sides in the client's conflicts, and taking an unbiased stance, from which they can examine the client's experiences and conflicts accurately, constitute the neutral stance. The therapist's silence and avoiding countertransference provides a noise-free condition, which facilitates the development of self-awareness.

In-depth examination of all the client's thoughts, feelings and behaviors from now to the past, from the surface to the depth, and here and now, and in transference relation and its ups and downs, too extra-transference relations and relation with important people from the past, sorting out and discovering the relations between them, and recognizing the constant patterns constitute the in-depth and insightful inquiry.

Interpretation of the client's experiences, which sheds light on patterns, was done considering several factors: timing of the interpretation, the dose of interpretations, which started by connecting small topics and continued to deep and general pattern, and interpretation tact, which was accurately based on the client's mental experiences.

##### 4.2.2. Client's Emotional Expression

This includes catharsis by the client, verbalization of the deepest parts of self, and expression of feelings at the time of regression.

Expression of deep, repressed, and painful feelings at the time of inquiry and regression can comfort the client and lead to connection to the deepest parts of self, which had never been reached before. Verbalizing these feelings not only makes the clients hear themselves for the first time, but also helps the development of self-awareness.

## 5. Discussion

Analyzing the experiences of clients suffering from depressive disorder about object relation therapy based on transference and potential space resulted in 2 categories. From the patient's viewpoint, the formation of analytic dyad creates a safe space for the development of self-awareness and self-expression, which in turn through creating a safe space filled with trust facilitates self-inquiry, helps to consolidate the analytic dyad and advances the therapy.

The formation of analytic dyad created a safe space for the patient through an intimate, safe and trustful relation, optimal responsiveness toward the client's emotional needs, and use of non-interpretive techniques in order to create therapeutic alliance. This helped the patient to gradually talk about feelings, events or fantasies, which were unpleasant, dangerous or even forbidden. The therapist could facilitate the acceptance of such feelings and open the space to talk about such topics through keeping and tolerating them. This was in accordance with previous experimental research of importance of therapeutic relation (11, 12). The emphasis of the present study on analytic dyad conforms to psychoanalytic writings about the significance of new relational experiences in therapy (13, 14), especially in interpersonal viewpoint of psychoanalysis (15-17). Therapeutic relationship is considered as the re-creation of child-parent relationship, which aims to mend deficits (18). According to Wallin (19), "psychotherapy is a kind of self-transformation through relationship in which the attachment of the client to the therapist is a fundamental issue because it provides a secure base for inquiry, development and change". In this study, patients had the opportunity to discover, test, and revise the internal representation of themselves and others with the therapist in a trustful space. The client's roots of patterns were discovered with active participation of the therapist. As a result of this, the patient found the therapeutic space safer, which facilitated self-awareness and self-expression.

In addition, the clients described the category of development of self-awareness and self-expression as a pro-

cess in which they made a joint inquiry with the therapist. Here, the therapist was considered as an active participant who intervened with asking questions, relating topics, examining feelings, behaviors and fantasies, and interpreting. The results of this study are in line with the findings of the studies (11, 12, 20), which emphasize the development of self-awareness and inquiry into patterns. The significance of discovering patterns in a person's behavior and relating them to the past, as well as, addressing transference and countertransference conforms to the psychoanalytic literature, which considers the development of self-expression as a principle in therapy (21, 22). Moreover, the emphasis of the current study on the patient's ability for self-expression and free association agrees with the strategies of psychoanalysis in therapeutic action (23, 24).

There has long been a gap between 2 approaches: the ones that emphasize interpretation, unconscious conflicts, memories of early experiences, unconscious fantasies, defense mechanisms, resistance, transference, and insight in therapeutic action, as well as the ones that emphasize on therapeutic relationship in therapeutic action and, as an example, gaining insight by the client was related to the gaining insight in the relationship. This gap has had an effect on the therapist's listening and intervention. A number of studies have addressed both approaches (for example, (25-30)), however, in none of them, have these concepts studied as interactive concepts with mutual effect. In other words, each study has focused on one aspect of the psychoanalytic therapy and has either ignored the other essential aspect or considered it as less important. Since the present study emphasizes the importance of both therapeutic relationship and insight and unconscious interpretation, and conceptualizes them in an integrated and uniform manner, it offers a new perspective on the therapeutic process. This new prospect is achieved in the context of transitional space and is derived from the mental experiences of the clients and, more than the merely theoretical approaches, benefits from clients' metaphors and words in understanding the therapeutic action. From the clients' point of view, the therapeutic process, in object relation therapy based on transference and potential space uses therapeutic relation and expansion of self-consciousness and discovering patterns in the therapeutic action, both of which interact with each other. Contrary to the articles mentioned above, in which some theorists emphasize on the relational experience in therapy and consider the interpretation as a form of relational experience, and others highlight client's intrapsychic experiences and interpretation of subconscious conflicts, the results of the present study show that this approach, from clients' point of view, integrates these 2 important aspects of therapy. The findings showed that formation of analyti-

cal dyad and creation of a safe therapeutic space provide a space for exploring habitual and pathological patterns and emotional impulses that have never found the opportunity to be expressed. Moreover, investigation of such patterns and developing insight into them lead to an increase in clients' trust in the presence of the therapist and the safe therapeutic space and facilitate the positive self-inquiry cycle. Both of these approaches can lead to new ways of being and relating to others.

The study has some limitations. Since the object relation therapy based on transference and potential space has been newly introduced in Iran, not many therapists are trained in it and use it in their work. This limited the selection of participants for the study. In addition, because the majority of the clients are from the socially and economically upper-class, there was a limitation in choosing people from the lower-class.

Future researches can address facilitating and hindering factors of therapy in this model. Conducting a research based on grounded theory and developing a theory, which originates from clients' mental experience of therapeutic process, can play a significant role in gaining insight into various aspects of this model of therapy. In addition, in future studies, integrating a qualitative and quantitative approach can lead to a better understanding of therapeutic action in object relation therapy based on transference and potential space. Investigating therapeutic action from therapists' point of view can play a significant role in making this model more applicable.

### 5.1. Conclusion

The present study confirmed that the formation of safe therapeutic space can result in a sense of security due to the trustworthiness of the other individuals presence. This space creates the appropriate context for exploring habitual and pathological patterns in transference and extra-transference space and provides space for expression of repressed emotions. The importance of safe therapeutic space becomes more evident when insight is gained into pathological patterns and is challenged in the therapeutic process. In this circumstance, clients need another person who can feel and contain their sense of threat so that they can explore new ways of being and relating.

### Acknowledgments

The authors would like to thank all the participants in the study.

## Footnotes

**Authors' Contribution:** Samaneh Ghafoori, Mohammad-Kazem Atef-Vahid, Frank Summers, Mahmood Dehghani designed and conducted the study. Samaneh Ghafoori collected the clinical data. Samaneh Ghafoori, Mohammad-Kazem Atef-Vahid, Mahmood Dehghani, Farahnaz Mohammadi Shahboulaghi, Aliasghar Asgharnejad Farid performed data analysis. Samaneh Ghafoori and Mohammad-Kazem Atef-Vahid drafted the manuscript. Mahmood Dehghani and Farahnaz Mohammadi Shahboulaghi revised the manuscript. All the authors read and approved the final manuscript.

**Declaration of Interests:** None declared.

**Funding/Support:** None declared.

## References

- WHO . Depression 2016. Available from: <http://www.who.int/mediacentre/factsheets/fs369/en/>.
- Klein M. In: Love, guilt and reparation. 1st ed. New York: Free Press; 1937. pp. 306-43. Love, guilt and reparation.
- Summers F. In: Comparative treatments of depression. Reinecke M DM, editor. United States of America: Springer Publishing Company; 2002. pp. 112-43. An object relations view of depression.
- Summers F. Self creation: Psychoanalytic therapy and the art of the possible. 75. Hillsdale, NJ: Analytic Press; 2005.
- Summers F. Object Relations Theories and Psychopathology: A Comprehensive Text. New Jersey: The Analytic Press; 1994.
- Summers F. Kohut's vision and the nuclear program of the self. *Int J Psychoanal Self Psychol*. 2011;**6**(3):289-305. doi: [10.1080/15551024.2011.583334](https://doi.org/10.1080/15551024.2011.583334).
- Summers F. The psychoanalytic vision: the experiencing subject, transcendence, and the therapeutic process. New York: Routledge: Taylor & Francis Group; 2013. p. 211.
- Bastos AG, Guimaraes LS, Trentini CM. The efficacy of long-term psychodynamic psychotherapy, fluoxetine and their combination in the outpatient treatment of depression. *Psychother Res*. 2015;**25**(5):612-24. doi: [10.1080/10503307.2014.935519](https://doi.org/10.1080/10503307.2014.935519). [PubMed: 25041333].
- Knekt P, Lindfors O, Laaksonen MA. Review: long term psychodynamic psychotherapy improves outcomes in people with complex mental disorders. *Evid Based Ment Health*. 2009;**12**(2):56. doi: [10.1136/ebmh.12.2.56](https://doi.org/10.1136/ebmh.12.2.56). [PubMed: 19395615].
- Graneheim UH, Lundman B. Qualitative content analysis in nursing research: concepts, procedures and measures to achieve trustworthiness. *Nurse Educ Today*. 2004;**24**(2):105-12. doi: [10.1016/j.nedt.2003.10.001](https://doi.org/10.1016/j.nedt.2003.10.001). [PubMed: 14769454].
- Elliott R, James E. Varieties of client experience in psychotherapy: An analysis of the literature. *Clin Psychol Rev*. 1989;**9**(4):443-67. doi: [10.1016/0272-7358\(89\)90003-2](https://doi.org/10.1016/0272-7358(89)90003-2).
- Lilliengren P, Werbart A. A model of therapeutic action grounded in the patients' view of curative and hindering factors in psychoanalytic psychotherapy. *Psychother Theory Res Pract Train*. 2005;**42**(3):324-39. doi: [10.1037/0033-3204.42.3.324](https://doi.org/10.1037/0033-3204.42.3.324).
- Balint M. In: Primary love and psychoanalytic technique. Balint M , editor. London: Karnac Books; 1985. pp. 159-73. Character analysis and new beginning.
- Alexander F. In: Psychoanalytic therapy: Principles and application. Alexander F, French TM, editors. New York: Roland Press; 1946. pp. 66-70. The principle of corrective emotional experience.
- Aron L. One person and two person psychologies and the method of psychoanalysis. *Psychoanal Psychol*. 1990;**7**(4):475-85. doi: [10.1037/0736-9735.7.4.475](https://doi.org/10.1037/0736-9735.7.4.475).
- Mitchell SA. Relational concepts in psychoanalysis: an integration. Cambridge, Massachusetts: Harvard University Press; 1988. p. 326.
- Stern DN, Sander LW, Nahum JP, Harrison AM, Lyons-Ruth K, Morgan AC, et al. Non-interpretive mechanisms in psychoanalytic therapy. The 'something more' than interpretation. The Process of Change Study Group. *Int J Psychoanal*. 1998;**79** ( Pt 5):903-21. [PubMed: 9871830].
- Mayes LC, Spence DP. Understanding therapeutic action in the analytic situation: a second look at the developmental metaphor. *J Am Psychoanal Assoc*. 1994;**42**(3):789-817. doi: [10.1177/000306519404200306](https://doi.org/10.1177/000306519404200306). [PubMed: 7963231].
- Wallin DJ. Attachment in Psychotherapy. New York: The Guilford Press; 2007.
- Leuzinger-Bohleber M. In: Outcomes of psychoanalytic treatments: Perspectives for therapists and researchers. Leuzinger-Bohleber M, Target M, editors. London: Whurr; 2002. pp. 143-73. A follow-up study: Critical inspiration for our clinical practice.
- Freud S. In: The standard edition of the complete psychological works of Sigmund Freud. Strachey J, editor. 7:122. London: Hogarth Press; 1905. Fragment of an analysis of a case of hysteria.
- Freud S. In: The standard edition of the complete psychological works of Sigmund Freud. Strachey J, editor. London: Hogarth Press; 1917. pp. 239-58. Mourning and melancholia.
- Freud S. In: The standard edition of the complete psychological works of Sigmund Freud. Strachey J, editor. London: Hogarth Press; 1895. pp. 255-305. Psychotherapy of hysteria.
- Freud S. In: The standard edition of the complete psychological works of Sigmund Freud. Strachey J , editor. London: Hogarth Press; 1920. pp. 7-64. Beyond the pleasure principle.
- Aisenstein M. On therapeutic action. *Psychoanal Q*. 2007;**76** Suppl:1443-61. doi: [10.1002/j.2167-4086.2007.tb00315.x](https://doi.org/10.1002/j.2167-4086.2007.tb00315.x). [PubMed: 18286756].
- Eizirik CL. On the therapeutic action of psychoanalysis. *Psychoanal Q*. 2007;**76** Suppl:1463-78. doi: [10.1002/j.2167-4086.2007.tb00316.x](https://doi.org/10.1002/j.2167-4086.2007.tb00316.x). [PubMed: 18286757].
- Friedman L. Who needs theory of therapeutic action? *Psychoanal Q*. 2007;**76** Suppl:1635-62. doi: [10.1002/j.2167-4086.2007.tb00324.x](https://doi.org/10.1002/j.2167-4086.2007.tb00324.x). [PubMed: 18286765].
- Greenberg J. Therapeutic action: convergence without consensus. *Psychoanal Q*. 2007;**76** Suppl:1675-88. doi: [10.1002/j.2167-4086.2007.tb00326.x](https://doi.org/10.1002/j.2167-4086.2007.tb00326.x). [PubMed: 18286767].
- Kernberg OF. The therapeutic action of psychoanalysis: controversies and challenges. *Psychoanal Q*. 2007;**76** Suppl:1689-723. doi: [10.1002/j.2167-4086.2007.tb00327.x](https://doi.org/10.1002/j.2167-4086.2007.tb00327.x). [PubMed: 18286768].
- Smith HF. In search of a theory of therapeutic action. *Psychoanal Q*. 2007;**76** Suppl:1735-61. doi: [10.1002/j.2167-4086.2007.tb00329.x](https://doi.org/10.1002/j.2167-4086.2007.tb00329.x). [PubMed: 18286770].