



# Evaluation of Psychiatric Counseling for Patients Hospitalized with COVID-19 in a University-Affiliated General Hospital in Tehran, Iran

Razieh Salehian <sup>1</sup>, Mehdi Nasr Esfahani <sup>2</sup>, Daem Roshani <sup>3,4</sup> and Mahnaz Abdi <sup>1,\*</sup>

<sup>1</sup>Rasoul-e-Akram Hospital, Iran University of Medical Sciences, Tehran, Iran

<sup>2</sup>Mental Health Research Center, School of Behavioral Sciences and Mental Health, Tehran Psychiatry Institution, Iran University of Medical Sciences, Tehran, Iran

<sup>3</sup>Social Determinants of Health Research Center, Research Institute for Health Development, Kurdistan University of Medical Sciences, Sanandaj, Iran

<sup>4</sup>Department of Epidemiology and Biostatistics, Medical School, Kurdistan University of Medical Sciences, Sanandaj, Iran

\*Corresponding author: Rasoul-e-Akram Hospital, Iran University of Medical Sciences, Tehran, Iran. Email: heroabdi@yahoo.com

Received 2021 July 04; Revised 2022 April 30; Accepted 2022 July 10.

## Abstract

**Background:** Coronavirus disease 2019 (COVID-19) is associated with a number of psychiatric problems in patients; however, mental healthcare in these patients does not appear to be adequate.

**Objectives:** This study aimed to evaluate the extent of psychological care provided to COVID-19 patients admitted to the hospital and the most common reasons for psychiatric counseling.

**Methods:** This study examined the electronic medical records of all COVID-19-positive patients admitted to Rasoul-e-Akram hospital in Tehran, Iran, within April 1 and September 30, 2020 (approximately 2200 cases).

**Results:** Psychiatric consultation was requested for 5.45% of admitted patients with COVID-19 diagnosis based on a positive real-time reverse transcription-polymerase chain reaction. More than half of those who received psychiatric counseling had a previous psychiatric disorder. The most common reason for requesting psychiatric counseling in patients with a previous psychiatric disorder was the patient's evaluation and management; the most common reason in patients without a psychiatric disorder was agitation and lack of cooperation. The most common diagnosis in patients with a positive history of psychiatric disorder was bipolar disorders (33.9%), followed by opioid use disorder (29%). Adjustment disorder (54%) and delirium (26%) were the most common diagnoses in patients without a history of psychiatric disease.

**Conclusions:** The current study showed that psychiatric counseling in COVID-19 patients was limited to a small number, including those with a positive history of psychiatric illnesses and prominent psychiatric symptoms.

**Keywords:** Adjustment Disorder, Counseling, COVID-19, Mental Health

## 1. Background

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) is highly transmissible and causes coronavirus disease 2019 (COVID-19), which is often lethal and has reached a pandemic scale worldwide in 2020 (1). In December 2019, the first case of COVID-19 was reported in Wuhan, China, and the disease rapidly spread to the rest of the world. In January 2020, the World Health Organization declared COVID-19 as a public health emergency threatening all individuals of the world (2). Therefore, the COVID-19 pandemic represents a threatening and potentially traumatic event, as it can lead to the hospitalization of numerous patients and even death.

The COVID-19 is associated with multiple psychiatric problems in several groups, including patients with COVID-19 and patients with psychiatric disorders pre-dating the pandemic, requiring quick and immediate

management of mental health (3-5). A few studies describe that psychiatric symptoms and disorders arise during the COVID-19 epidemic due to biological and psychosocial factors (6-8). A wide variety of neuropsychiatric symptoms can appear during COVID-19 infection and might be related to direct damage to neurological tissues or due to environmental issues, cytokine release, respiratory insufficiency, critical illness, and side effects of pharmacological treatment or exacerbation of previous neuropsychiatric illness (9).

Currently, the focus is mainly on preventing COVID-19 infection and discussing issues related to physical health consequences. Therefore, COVID-19 clinics are being established in different places to address the physical health consequences of the pandemic. However, it is important to note that although some infected patients face life-threatening physical consequences, a considerable num-

ber of infected patients face negative mental consequences as a result of the outbreak. It seems that mental health providers have not been highly involved in the treatment of those affected by the pandemic, and mental healthcare in these patients does not appear to be adequate (10, 11).

## 2. Objectives

Since there is limited information on psychiatric assessments in patients with COVID-19, this study aimed to evaluate the extent of psychological care provided to COVID-19 patients admitted to a hospital, the most common reasons for psychiatric counseling, and the results of psychiatric counseling retrospectively. It is hoped that by evaluating psychiatric counseling and identifying the main psychiatric problems in these patients, more comprehensive interventions for patients during hospitalization can be considered, along with more comprehensive psychiatric assessments and interventions for new patients.

## 3. Methods

### 3.1. Study Design and Data Collection

Following the approval of the Ethics Committee of Iran University of Medical Sciences, Tehran, Iran, this retrospective observational study was conducted in one of Tehran's largest referral hospitals for COVID-19 patients (Rasoul-e-Akram hospital). This study examined the electronic medical records of all COVID-19-positive patients admitted to the hospital within April 1 and September 30, 2020 (approximately 2200 cases).

The patients who requested psychiatric consultation while being hospitalized and had a confirmed laboratory diagnosis of COVID-19 (either a positive result for immunoglobulin G/immunoglobulin M antibodies against SARS-CoV-2 in a blood test or identification of SARS-CoV-2 ribonucleic acid with a real-time reverse transcription-polymerase chain reaction of throat swab samples) were chosen and analyzed in this study. Suspected COVID-19 patients were excluded from the study. Demographic data, such as age, gender, marital status, educational level, previous psychiatric and/or medical comorbidities (e.g., hypertension, diabetes, dyslipidemia, heart disease, chronic kidney disease, immunosuppression, cancer, and neurological diseases), substance abuse and smoking habit, and the reason for seeking psychiatric consultation from the physician (i.e., primary psychiatric symptom), were reported. Final psychiatric diagnosis according to the diagnostic and statistical manual of mental disorders, 5th edition (DSM-5), psychiatrist's recommendations for pharmacological and

nonpharmacological interventions in response to counseling, taken actions, and results were documented. Finally, the information collected from 112 patients was examined. Two experienced psychiatrists collaborated to collect the data using a standardized case-report form. Before the formal study, the psychiatrists were trained on how to collect the data from electronic medical records.

### 3.2. Ethical Statement

This study was approved by the Ethics Committee of the Iran University of Medical Sciences, Tehran, Iran (code: IR.IUMS.REC.1399.594). This was a retrospective and non-interventional study. No experiments were carried out on human subjects. A waiver of written informed consent was obtained since all the data were collected retrospectively and anonymously.

### 3.3. Statistical Analysis

The characteristics of COVID-19 patients with and without previous psychiatric disease were analyzed and compared using descriptive statistics. The chi-square test or Fisher's exact test was used to compare two groups of COVID-19 patients with and without previous psychiatric illness. The independent-sample *t*-test and Mann-Whitney *u* test were used to compare the continuous variables between these two groups of patients according to their normal distribution and deviation distribution (a skewed distribution), respectively. The statistical significance level was set at less than 0.05 (two-sided). Confidence intervals (CI) were calculated if indicated. All data analyses were carried out using SPSS software (version 24.0).

## 4. Results

After a detailed review of medical records of 2200 COVID-19 patients during that period, 120 patients were identified for whom psychiatric consultation was requested. In other words, psychiatric consultation was requested for 5.45% of admitted patients with COVID-19. Eight patients were removed due to uncertain COVID-19 infection and/or incomplete data. Finally, 112 cases were included in the study, among whom psychiatric consultation was requested for 62 patients due to a positive history of psychiatric diseases, including substance use disorders, and for 50 patients due to new-onset psychiatric symptoms during hospitalization without any known psychiatric illness.

According to the DSM-5, occasional consumption of substances, including alcohol or tobacco smoking, was not considered a psychiatric disorder. In the current study, the most common reason for requesting consultation was

a positive history of psychiatric disorder. Opioid use disorder and agitation were ranked second (Figure 1). The most common diagnosis in patients with a positive history of a psychiatric disease was bipolar disorders (33.9%), followed by opioid use disorder (29%). Adjustment disorder (54%) and delirium (26%) were the most common diagnoses in patients without a history of psychiatric disease. It should be noted that the mean age of patients with a positive history of psychiatric disorders was significantly lower than other patients (54.95 years in comparison to 63.02 years). Furthermore, the prevalence of medical comorbidities, such as hypertension, heart diseases, dyslipidemia, and kidney diseases, in the patients without a history of mental illness was higher than in the other group (Table 1). The study results showed that the mortality rate was higher in patients with a positive history of psychiatric issues; nevertheless, there was no statistically significant difference in mortality between psychiatric illnesses (Table 1). Figure 1 depicts the distribution of the reasons for counseling requests and primary psychiatric symptoms. Tables 1 and 2 show the characteristics of patients and final diagnosis and interventions, respectively.

## 5. Discussion

This new study was conducted to determine the most common reasons for psychiatric counseling, evaluate psychiatric counseling performed for COVID-19 patients, and compare the common psychiatric symptoms, psychiatric diagnoses, and outcomes among COVID-19 patients who received psychiatric counseling between patients with and without previous psychiatric diseases.

In the current study, psychiatric counseling was requested for only 5.45% of admitted COVID-19 patients, more than half of which were related to cases already known to have psychiatric illnesses. According to previous studies, numerous COVID-19 patients have psychological symptoms and issues, and COVID-19 is frequently correlated with neuropsychiatric diagnosis during the acute phase of the disease (12, 13). A cross-sectional study conducted on 199 COVID-19 patients admitted to Sari hospitals in Iran to examine posttraumatic stress disorder (PTSD) symptoms through the PTSD Checklist for DSM-5 showed that the prevalence of PTSD was 19.1% (14). Another cross-sectional study that evaluated the prevalence and severity of depression, anxiety, stress (based on the Depression Anxiety Stress Scale-21), and stress perception (based on Perceived Stress Scale 4) among 106 inpatients with COVID-19 infection in Alborz, Iran, revealed that 97.2% of patients had some degree of depression and 85.8% of the patients had severe depression. All patients (100%) had anxiety, and

97.1% had some degree of stress. Moreover, 73.6% of patients had high levels of perceived stress (12). Therefore, the low percentage of psychiatric counseling in this study might indicate neglect of psychological issues in these patients.

Agitation, lack of cooperation with care providers, and anxiety were the most common psychiatric symptoms in patients with new-onset psychiatric symptoms, leading to requesting psychiatric counseling, and adjustment disorder was the most common psychiatric diagnosis in this group. Since 6% of patients with agitation and lack of cooperation did not receive a formal diagnosis during a psychiatric evaluation, and 20% of patients did not need pharmacotherapy, their symptoms were justified by a lack of awareness and information about their disease's condition. In the present pandemic, this issue could justify the importance of psychoeducation and psychological care for these patients.

One retrospective cohort study of 62354 COVID-19 patients in the USA revealed that a psychiatric diagnosis in the previous year was associated with a higher incidence of COVID-19 diagnosis (relative risk = 1.65, 95% CI 1.59 - 1.71;  $P < 0.0001$ ), and the risk was independent of known physical health risk factors for COVID-19 (15). Only 12% of patients with a prior documented psychiatric condition were in complete remission in the current study, which might suggest a higher incidence of COVID-19 in uncontrolled psychiatric patients, and psychiatric diagnosis might be an independent risk factor for COVID-19 (15, 16).

In the current study, despite the higher mean age and higher prevalence of comorbidities, such as hypertension, heart diseases, dyslipidemia, and kidney diseases, in the population without a history of mental illness than in the other group, the mortality rate was higher in patients with a positive history of psychiatric issues. This result is consistent with the results of one cohort study conducted on 1685 hospitalized COVID-19 patients in the northeast region of the United States, indicating that after controlling for demographic characteristics, medical comorbidities, and hospital location, the risk of mortality remained significantly greater among patients with a psychiatric disorder than those with no psychiatric diagnosis (hazard ratio = 1.5; 95% CI, 1.1 - 1.9;  $P = 0.003$ ) (17). Unlike another study that showed that COVID-19 mortality was higher in patients with schizophrenia than in patients with other psychiatric disorders (18), in the current study, there was no statistically significant difference in mortality between psychiatric illnesses.


Although the present study provides important data on the psychological problems of hospitalized COVID-19 patients, there are still several limitations. Since the data were collected retrospectively, there might be selection

**Table 1.** Characteristics and Comorbidities of 112 Inpatients with Coronavirus Disease 2019<sup>a</sup>

Variables	Patients with a Positive History of Psychiatric Illness (n = 62)	Patients Without Previous Psychiatric Illness (n = 50)
Age (y)	(54.95 ± 16.51) <sup>b</sup>	(63.02 ± 17.46) <sup>b</sup>
<b>Gender</b>		
Male	40 (64.5)	31 (62)
Female	22 (35.5)	19 (38)
<b>Marital status</b>		
Single	14 (22.6)	1 (2)
Married	34 (54.8)	37 (74)
Divorced/widowed	14 (22.6)	12 (24)
<b>Educational level</b>		
Illiterate	12 (19.4)	13 (26)
Under diploma	29 (46.8)	18 (36)
Diploma/associate degree	17 (27.4)	13 (26)
Bachelor's degree and higher	4 (6.5)	6 (12)
<b>Employment</b>		
Housewife	18 (29.0)	17 (34)
Unemployed	19 (30.6)	12 (24)
Freelance	11 (17.7)	11 (22)
Employee	2 (3.2)	5 (10)
Retired	12 (19.4)	5 (10)
<b>Alcohol and/or substance abuse</b>		
Tobacco smoking <sup>b</sup>	26 (41.9)	10 (20)
Alcohol use/abuse	6 (9.7)	2 (4)
Opioid use disorder	18 (29)	-
Stimulant/Inhalant use disorder	7 (11.3)	-
<b>Comorbidities</b>		
Hypertension <sup>b</sup>	17 (27.4)	29 (58)
Diabetes mellitus	15 (24.2)	19 (38)
Dyslipidemia <sup>b</sup>	9 (14.5)	19 (38)
Heart disease <sup>b</sup>	6 (9.7)	15 (30)
Chronic kidney disease <sup>b</sup>	3 (4.8)	7 (14)
Malignancy	3 (4.8)	1 (2)
Neurological disease	10 (16.1)	9 (18)
Rheumatologic disease	1 (1.6)	2 (4)
Pulmonary disease	1 (1.6)	1 (2)
Thyroid disease	4 (6.5)	4 (8)
<b>Previous psychiatric illness status at the time of hospitalization</b>		
Active	35 (56.5)	-
Partial remission	19 (30.6)	-
Complete remission	8 (12.9)	-
<b>Positive history of psychiatric admission</b>	31 (50)	-
<b>Transfer from psychiatric service to COVID-19 ward</b>	5 (8.1)	-
<b>ICU admission</b>	16 (25.8)	20 (40)
<b>Outcome</b>		
Death	17 (27.4) <sup>b</sup>	9 (18) <sup>b</sup>
Discharge	39 (62.9) <sup>b</sup>	41 (82) <sup>b</sup>
Transfer to psychiatric service	6 (9.7) <sup>b</sup>	-

Abbreviations: COVID-19, coronavirus disease 2019; ICU, intensive care unit

<sup>a</sup> Values are expressed as mean ± SD or No. (%).<sup>b</sup> P < 0.05



**Figure 1.** Distribution of reasons for psychiatric counseling requests

bias and some missing crucial details. Furthermore, since the study was a descriptive one, it could not be concluded that the patients' psychological symptoms were triggered by the SARS-CoV-2 infection or by other factors, such as cross-immunity, inflammatory reactions, or side effects of treatments. Finally, the data were collected from only one hospital, and the small sample size might limit the generalization of the results. Since this study was carried out in a hospital, the prevalence of psychological issues among COVID-19 patients in the community could differ, and any

findings should be interpreted accordingly.

### 5.1. Conclusions

Despite the high prevalence of psychiatric problems in patients with COVID-19, in the current study, psychiatric counseling was limited to a small number of patients, including those with a positive history of psychiatric illnesses and prominent psychiatric symptoms. It is suggested that first-line physicians be aware of possible psychiatric complications since the global emergency

**Table 2.** Diagnosis and Psychiatric Interventions after Psychiatric Counseling in Patients with Coronavirus Disease 2019 <sup>a</sup>

Variables	Patients with a Positive History of Psychiatric Disease (n = 62)	Patients Without a Previous Psychiatric Disease (n = 50)
<b>Diagnosis</b>		
Bipolar disorders	21 (33.9)	-
Adjustment disorder	-	27 (54)
Opioid use disorder	18 (29)	-
Psychotic disorders	11 (17.7)	1 (2)
Depressive disorders	9 (14.5)	3 (6)
Delirium	7 (11.3)	13 (26)
Anxiety disorders	-	3 (6)
Stimulant/inhalant use disorder	7 (11.3)	-
Ruling out psychiatric disorders	-	3 (6)
Cluster B personality disorder	5 (8.1)	-
Others <sup>b</sup> (some patients had multiple simultaneous diagnoses)	8 (12.9)	-
<b>Pharmacological intervention</b>		
No need for pharmacotherapy	3 (4.8)	10 (20)
Antipsychotics	39 (62.9)	23 (46)
Mood stabilizer	27 (43.5)	-
SSRIs	16 (25.8)	8 (16)
Benzodiazepines	15 (24.2)	18 (36)
Methadone	14 (22.6)	-
Anticonvulsants	-	1 (2)
TCAs	1 (1.6)	-
Antihistamines	1 (1.6)	2 (4)
Others	2 (3.2)	-
<b>Nonpharmacological intervention</b>	11 patients (17.7%) received nonpharmacological interventions (e.g., psychoeducation and supportive psychotherapy)	42 patients (84%) received nonpharmacological interventions (e.g., psychoeducation and supportive psychotherapy)

Abbreviations: SSRIs, selective serotonin reuptake inhibitors; TCAs, tricyclic antidepressants

<sup>a</sup> Values are expressed as No. (%).

<sup>b</sup> Obsessive-compulsive disorder, anxiety disorders, posttraumatic stress disorder and intellectual disability

is expected to last. All COVID-19 patients should be assessed mentally, according to the recommendations of this study. It is also recommended that psychiatrists be included in COVID-19 response teams on a regular basis to ensure the adequate early detection and treatment of psychiatric manifestations, thereby improving psychiatric outcomes. Accordingly, it is required to further investigate the mechanisms of mental injuries and their long-term effects.

### Acknowledgments

The authors would like to express their heartfelt respect to all healthcare workers who are fighting against the COVID-19 pandemic on the front line.

### Footnotes

**Authors' Contribution:** Study concept and design: R. S. and M. A. Acquisition of the data: R. S. and M. A. Analysis and interpretation of the data: M. A. and R. S. Drafting of the manuscript: M. A. Critical revision of the manuscript for important intellectual content: R. S. and M. N. Statistical analysis: M. A. and D. R. Study supervision: R. S. and M. N.

**Conflict of Interests:** There is no conflict of interests.

**Data Reproducibility:** Since the information is confidential and contained in the patient's hospital record, the data is not publicly available.

**Ethical Approval:** This study was approved by the

Ethics Committee of the Iran University of Medical Sciences, Tehran, Iran (code: IR.IUMS.REC.1399.594). Link: [ethics.research.ac.ir/ProposalCertificateEn.php?id=154864](https://ethics.research.ac.ir/ProposalCertificateEn.php?id=154864).

**Funding/Support:** This study did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

**Informed Consent:** This was a retrospective and noninterventive study. No experiments were carried out on human subjects. A waiver of written informed consent was obtained since all the data were collected retrospectively and anonymously.

## References

- World Health Organization. *Coronavirus disease 2019 (COVID-19) Situation Report - 69*. Geneva: World Health Organization; 2020, [cited 2022]. Available from: <https://www.who.int/docs/default-source/coronavirus/situation-reports/20200329-sitrep-69-covid-19.pdf>.
- Moghadami M, Hassanzadeh M, Wa K, Hedayati A, Malekolkalami M. Modeling the Corona Virus Outbreak in IRAN. *medRxiv*. 2020; **March**. doi: [10.1101/2020.03.24.20041095](https://doi.org/10.1101/2020.03.24.20041095).
- Benson NM, Ongur D, Hsu J. COVID-19 testing and patients in mental health facilities. *Lancet Psychiatry*. 2020; **7**(6):476-7. doi: [10.1016/S2215-0366\(20\)30198-X](https://doi.org/10.1016/S2215-0366(20)30198-X). [PubMed: [32407671](https://pubmed.ncbi.nlm.nih.gov/32407671/)]. [PubMed Central: [PMC7213967](https://pubmed.ncbi.nlm.nih.gov/PMC7213967/)].
- Yao H, Chen JH, Xu YF. Patients with mental health disorders in the COVID-19 epidemic. *Lancet Psychiatry*. 2020; **7**(4). e21. doi: [10.1016/S2215-0366\(20\)30090-0](https://doi.org/10.1016/S2215-0366(20)30090-0). [PubMed: [32199510](https://pubmed.ncbi.nlm.nih.gov/32199510/)]. [PubMed Central: [PMC7269717](https://pubmed.ncbi.nlm.nih.gov/PMC7269717/)].
- Zhu Y, Chen L, Ji H, Xi M, Fang Y, Li Y. The Risk and Prevention of Novel Coronavirus Pneumonia Infections Among Inpatients in Psychiatric Hospitals. *Neurosci Bull*. 2020; **36**(3):299-302. doi: [10.1007/s12264-020-00476-9](https://doi.org/10.1007/s12264-020-00476-9). [PubMed: [32096116](https://pubmed.ncbi.nlm.nih.gov/32096116/)]. [PubMed Central: [PMC7056754](https://pubmed.ncbi.nlm.nih.gov/PMC7056754/)].
- Varatharaj A, Thomas N, Ellul MA, Davies NWS, Pollak TA, Tenorio EL, et al. Neurological and neuropsychiatric complications of COVID-19 in 153 patients: a UK-wide surveillance study. *Lancet Psychiatry*. 2020; **7**(10):875-82. doi: [10.1016/S2215-0366\(20\)30287-X](https://doi.org/10.1016/S2215-0366(20)30287-X). [PubMed: [32593341](https://pubmed.ncbi.nlm.nih.gov/32593341/)]. [PubMed Central: [PMC7316461](https://pubmed.ncbi.nlm.nih.gov/PMC7316461/)].
- Galea S, Merchant RM, Lurie N. The Mental Health Consequences of COVID-19 and Physical Distancing: The Need for Prevention and Early Intervention. *JAMA Intern Med*. 2020; **180**(6):817-8. doi: [10.1001/jamainternmed.2020.1562](https://doi.org/10.1001/jamainternmed.2020.1562). [PubMed: [32275292](https://pubmed.ncbi.nlm.nih.gov/32275292/)].
- Holmes EA, O'Connor RC, Perry VH, Tracey I, Wessely S, Arseneault L, et al. Multidisciplinary research priorities for the COVID-19 pandemic: a call for action for mental health science. *Lancet Psychiatry*. 2020; **7**(6):547-60. doi: [10.1016/S2215-0366\(20\)30168-1](https://doi.org/10.1016/S2215-0366(20)30168-1). [PubMed: [32304649](https://pubmed.ncbi.nlm.nih.gov/32304649/)]. [PubMed Central: [PMC7159850](https://pubmed.ncbi.nlm.nih.gov/PMC7159850/)].
- Zubair AS, McAlpine LS, Gardin T, Farhadian S, Kuruville DE, Spudich S. Neuropathogenesis and Neurologic Manifestations of the Coronaviruses in the Age of Coronavirus Disease 2019: A Review. *JAMA Neurol*. 2020; **77**(8):1018-27. doi: [10.1001/jamaneurol.2020.2065](https://doi.org/10.1001/jamaneurol.2020.2065). [PubMed: [32469387](https://pubmed.ncbi.nlm.nih.gov/32469387/)]. [PubMed Central: [PMC7484225](https://pubmed.ncbi.nlm.nih.gov/PMC7484225/)].
- Lai J, Ma S, Wang Y, Cai Z, Hu J, Wei N, et al. Factors Associated With Mental Health Outcomes Among Health Care Workers Exposed to Coronavirus Disease 2019. *JAMA Netw Open*. 2020; **3**(3). e203976. doi: [10.1001/jamanetworkopen.2020.3976](https://doi.org/10.1001/jamanetworkopen.2020.3976). [PubMed: [32202646](https://pubmed.ncbi.nlm.nih.gov/32202646/)]. [PubMed Central: [PMC7090843](https://pubmed.ncbi.nlm.nih.gov/PMC7090843/)].
- Xiang YT, Jin Y, Cheung T. Joint International Collaboration to Combat Mental Health Challenges During the Coronavirus Disease 2019 Pandemic. *JAMA Psychiatry*. 2020; **77**(10):989-90. doi: [10.1001/jamapsychiatry.2020.1057](https://doi.org/10.1001/jamapsychiatry.2020.1057). [PubMed: [32275289](https://pubmed.ncbi.nlm.nih.gov/32275289/)].
- Zandifar A, Badrfam R, Yazdani S, Arzaghi SM, Rahimi F, Ghasemi S, et al. Prevalence and severity of depression, anxiety, stress and perceived stress in hospitalized patients with COVID-19. *J Diabetes Metab Disord*. 2020; **19**(2):1431-8. doi: [10.1007/s40200-020-00667-1](https://doi.org/10.1007/s40200-020-00667-1). [PubMed: [33145259](https://pubmed.ncbi.nlm.nih.gov/33145259/)]. [PubMed Central: [PMC7594988](https://pubmed.ncbi.nlm.nih.gov/PMC7594988/)].
- Rogers JP, Chesney E, Oliver D, Pollak TA, McGuire P, Fusar-Poli P, et al. Psychiatric and neuropsychiatric presentations associated with severe coronavirus infections: a systematic review and meta-analysis with comparison to the COVID-19 pandemic. *Lancet Psychiatry*. 2020; **7**(7):611-27. doi: [10.1016/S2215-0366\(20\)30203-0](https://doi.org/10.1016/S2215-0366(20)30203-0). [PubMed: [32437679](https://pubmed.ncbi.nlm.nih.gov/32437679/)]. [PubMed Central: [PMC7234781](https://pubmed.ncbi.nlm.nih.gov/PMC7234781/)].
- Hosseini SH, Elyasi F, Moosazadeh M, Ghafour I. Evaluation of the Frequency of Post-traumatic Stress Disorder in Patients with COVID-19 Admitted to Hospitals in Sari, Iran in 2020. *Iran J Psychiatry Behav Sci*. 2022; **16**(1). e115663. doi: [10.5812/ijpbs.115663](https://doi.org/10.5812/ijpbs.115663).
- Taquet M, Luciano S, Geddes JR, Harrison PJ. Bidirectional associations between COVID-19 and psychiatric disorder: retrospective cohort studies of 62354 COVID-19 cases in the USA. *Lancet Psychiatry*. 2021; **8**(2):130-40. doi: [10.1016/S2215-0366\(20\)30462-4](https://doi.org/10.1016/S2215-0366(20)30462-4).
- Xiang YT, Zhao YJ, Liu ZH, Li XH, Zhao N, Cheung T, et al. The COVID-19 outbreak and psychiatric hospitals in China: managing challenges through mental health service reform. *Int J Biol Sci*. 2020; **16**(10):1741-4. doi: [10.7150/ijbs.45072](https://doi.org/10.7150/ijbs.45072). [PubMed: [32226293](https://pubmed.ncbi.nlm.nih.gov/32226293/)]. [PubMed Central: [PMC7098035](https://pubmed.ncbi.nlm.nih.gov/PMC7098035/)].
- Li L, Li F, Fortunati F, Krystal JH. Association of a Prior Psychiatric Diagnosis With Mortality Among Hospitalized Patients With Coronavirus Disease 2019 (COVID-19) Infection. *JAMA Netw Open*. 2020; **3**(9). e2023282. doi: [10.1001/jamanetworkopen.2020.23282](https://doi.org/10.1001/jamanetworkopen.2020.23282). [PubMed: [32997123](https://pubmed.ncbi.nlm.nih.gov/32997123/)]. [PubMed Central: [PMC7527869](https://pubmed.ncbi.nlm.nih.gov/PMC7527869/)].
- Nemani K, Li C, Olfson M, Blessing EM, Razavian N, Chen J, et al. Association of Psychiatric Disorders With Mortality Among Patients With COVID-19. *JAMA Psychiatry*. 2021; **78**(4):380-6. doi: [10.1001/jamapsychiatry.2020.4442](https://doi.org/10.1001/jamapsychiatry.2020.4442). [PubMed: [33502436](https://pubmed.ncbi.nlm.nih.gov/33502436/)]. [PubMed Central: [PMC7841576](https://pubmed.ncbi.nlm.nih.gov/PMC7841576/)].