Published online 2024 February 6.

**Brief Report** 



# The Role of Emotion Regulation as a Mediator Between Difficulties in Emotion Regulation and Aggression and Bullying Behavior in Motorcyclists with Mild Traumatic Brain Injury

Elham Shafiei <sup>1</sup>, Arian Karimi Rouzbahani <sup>2,3</sup>, Arash Nademi <sup>4</sup> and Molouk Jafarpour <sup>5,\*</sup>

<sup>1</sup>Clinical Research Development Unit, Ayatollah Taleghani Hospital, Ilam University of Medical Sciences, Ilam, Iran

<sup>2</sup>USERN Office, Lorestan University of Medical Sciences, Khorramabad, Iran

<sup>3</sup>Student Research Committee, Lorestan University of Medical Sciences, Khorramabad, Iran

<sup>4</sup> Department of Statistics, Ilam Branch, Islamic Azad University, Ilam, Iran

<sup>5</sup>Department of Midwifery, School of Nursing and Midwifery, Ilam University of Medical sciences, Ilam, Iran

corresponding author: Department of Midwifery, School of Nursing and Midwifery, Ilam University of Medical sciences, Ilam, Iran. Email: jaafarpourm@medilam.ac.ir

Received 2023 May 28; Revised 2023 September 10; Accepted 2023 December 19.

## Abstract

**Background:** Traumatic brain injury (TBI) may be a cause of criminal behavior, affecting various aspects of an individual's mental health.

**Objectives:** The aim of the present study was to investigate the effect of difficulties in emotion regulation on psychiatric symptoms among patients with mild TBI.

**Methods:** This research was a retrospective cohort study that included 120 motorcyclists with mild TBI referred to the psychiatric clinic of Shahid Mostafa Khomeini Hospital, Ilam, during a 23-month period from January 2017 to March 2019. The data were analyzed using structural equation modeling with P < 0.05.

**Results:** The results of AMOS analyses revealed a direct path from maladaptive cognitive emotion regulation strategies and an indirect path from adaptive cognitive emotion regulation to the relationship between difficulties in emotion regulation (DERS), bullying behavior, and aggression in 120 motorcyclists with TBI.

**Conclusions:** Reducing TBI rates, which is a major public health necessity, may have benefits in terms of reducing crime and improving clinical services.

Keywords: Difficulties in Emotion Regulation Scale, Bullying Behavior, Traumatic Brain Injury, Aggression

## 1. Background

Although hospital services and the system for caring for traumatic brain diseases have improved, the impact of TBI-related behavioral dysregulation, such as bullying behavior and aggression, on the occurrence of crimes remains significant. Traumatic brain injury (TBI) is a condition that results in disability and loss of years of life (1-5). While several potential mediators have been identified in previous studies (6), the role of cognitive emotion regulation as a mediator in the relationship between difficulties in emotion regulation (DERS) and current symptoms of bullying behavior and aggression is still relatively unexplored. Previous studies have examined each of these emotion regulation components independently as potential contributors to emotion regulation problems. However, the correlation between DERS and aggressive behavior has not been thoroughly investigated more generally (7). In recent years, research efforts have shifted towards identifying the elements that contribute to emotional tone. Research results have shown that individuals who use positive emotional strategies are less likely to experience depression and exhibit better adaptability. On the other hand, when individuals employ avoidance strategies, they tend to feel depressed and exhibit poor adaptability (8). Improved adaptability is associated with emotional expression. However, when emotions are not expressed, they remain

Copyright © 2024, Shafiei et al. This open-access article is available under the Creative Commons Attribution 4.0 (CC BY 4.0) International License (https://creativecommons.org/licenses/by/4.0/), which allows for unrestricted use, distribution, and reproduction in any medium, provided that the original work is properly cited.

unresolved, which can impact the patient's health and increase sympathetic nervous system activity (9). Given that TBI patients experience significant levels of stress and negative emotions in addition to the challenges they face, it is crucial to understand the emotional components of these patients' experiences and the strategies they employ to regulate their emotions (10). Therefore, uncovering these associations could assist in the management of the large numbers of patients with these disorders presenting to health services.

## 2. Objectives

Due to the limited number of studies that have examined the possible mediating variables, the current study aimed to explore the relationship between DERS, aggregation, and bullying behavior in TBI patients.

## 3. Methods

#### 3.1. Participants and Procedures

This research is a retrospective cohort study using community-based samples. Information on motorcyclists who survived a full year from their first TBI was collected to allow adequate time for an outcome to occur. They were advised to visit a mental clinic at the Shahid Mostafa Khomeini Hospital in Ilam, Iran, after being contacted. A psychiatrist conducted clinical interviews with the patients. They were instructed to bring the 5 questionnaires they had been given to their next appointment. Informed consent, demographic information, and neurological information of the patients were recorded in the questionnaires. Over the 23-month period from January 2017 to March 2019, 120 motorcyclists with mild TBI participated in the current research. The inclusion criteria for this study included brain trauma patients with a GCS score between 13 and 15 years, ages 18 to 50 years, willingness to participate in the study, no history of mental illness (such as bipolar disorder, personality disorder, severe psychosis), and no mental retardation, smoking, drug use, or alcohol consumption.

The exclusion criteria included non-cooperation in completing the questionnaire, moderate and severe traumatic injuries, and a previous history of head injury. The questionnaires were coded without including first and last names and were only marked with acronyms (A, B, C) to protect the participants' privacy and prevent the disclosure of their names.

### 3.2. Questionnaires

#### 3.2.1. The Cognitive Emotion Regulation Questionnaire (CERQ)

The CERQ is a 36-item scale with 9 subscales. For our analysis, we summed the CERQ-M and CERQ-A strategies (11).

#### 3.2.2. Bullying Behavior

The initial Olweus questionnaire is an abbreviated questionnaire comprising 28 items. Its validity and reliability have been confirmed in several studies, with a test-retest reliability estimated using Cronbach's alpha coefficient as 0.83 (12).

#### 3.2.3. Impulsive and Premeditated Aggression Scale

The Impulsive and Premeditated Aggression Scale (IPAS) (13) consists of 30 items for reporting aggressive acts over the past years. It has been used in various populations, including community adult samples (14, 15).

#### 3.2.4. Difficulties in Emotion Regulation Scale

This questionnaire was developed by Gratz and Roemer (2004) and comprises 36 items rated on a 5-point Likert scale. In a study conducted by Shafiei et al. (2016), the tool's reliability was confirmed as 0.89 using Cronbach's alpha (16).

#### 3.3. Data Analysis

Descriptive statistics were reported as mean  $\pm$  standard deviation (SD). Correlation coefficients were used for normally and non-normally distributed variables, respectively. The mediating variables were CERQ-A and CERQ-M sub-scores from the CERQ.

After completing the questionnaires, the data were analyzed using SPSS AMOS software. Structural equation modeling based on the maximum likelihood method was applied to study the components of emotion regulation in patients with mTBI.

The most common statistical indices for evaluating the fit of the structural model, including the absolute fit (chi-square, CMIN), fit indices (root mean square error of approximation (RMSEA), normalized fit index (NFI), comparative fit index (CFI), and Bentler-Bonett normed fit index (NFI)), and the Tucker-Lewis index (TLI), were all calculated. If the chi-square is not statistically significant, it indicates an appropriate fit for the model. However, this indicator tends to be significant in larger samples and is not considered an appropriate index for model fit. Values close to 1 for the TLI, NFI, and CFI indices, values greater than 0.5 for the normalized fit index, and values less than or equal to 0.05 for the RMSEA index all indicate a good fit.

## 4. Results

#### 4.1. Demographics and Psychiatric Symptoms

In the present study, 120 participants were included, with an average age of  $34.00 \pm 11.56$  years. Of the participants, 61.6% were married, 35.5% were single, and 3.3% were divorced (Table 1).

Table 1. Demographic Characteristics of the Patients <sup>a</sup>				
Variables	No. (%)			
Marital status				
Married	74 (61.6)			
Single	42 (35)			
Divorced	4 (3.3)			
Education level				
Illiterate	10 (8.3)			
High school	55 (45.83)			
High school diploma	30 (25)			
Bachelor's degree	20 (16.66)			
Master's degree	3 (2.5)			
PhD	2 (1.6)			

<sup>a</sup> Values are expressed as No. (%).

Table 2 summarizes all the information, including means, SDs, and ranges. The overall mean score for bullying behavior was  $13.3 \pm 0.63$ . In this sample, 32.4% of participants met the cut-off for mild bullying, 41.8% for moderate bullying, and 25.8% for severe bullying behavior. The mean scores for CERQ-M and CERQ-A were  $53.48 \pm 11.0$  and  $70.1 \pm 13.31$ , respectively. The overall mean score for aggression was  $7.24 \pm 5.87$ . In this sample, 23.8% of participants met the cut-off for mild aggression, 39.7% for moderate aggression, and 36.5% for severe aggression.

By fitting the structural equation model and the final model, the fit indices for chi-square (CMIN = 10.762, df = 15, P = 0.096 > 0.05), Tucker-Lewis index (TLI = 0.857), comparative fit index (CFI = 0.943), and Bentler-Bonett normed fit index (NFI = 0.932) were obtained. Additionally, the value of the normalized fit index (PNFI) was 0.725, and the RMSEA was 0.031. These values indicate that the model fits the data well. Table 3 displays the regression coefficients for the obtained model.

Based on the results presented in Table 3, all the factors related to emotion regulation, cognitive adjustment, and emotional adjustment were significant at a 5% error level (P < 0.05). Furthermore, there was a significant correlation between the components of cognitive and emotional regulation and maladaptation at the 1% error level, with a correlation coefficient of 30% (P < 0.01).

Figure 1 illustrates the final structural model for measuring the emotional adjustment variable. In this model, the emotional adjustment variables, along with the components of cognitive emotion regulation that are compatible and incompatible with Bayesian symbols and the dimensions of their measurement with rectangular symbols, are displayed.

## 5. Discussion

Traumatic brain injury is defined as an injury that disrupts the brain's normal functioning, often caused by shaking, impact, or head contact. Due to the intricate interplay between acute (short-term) and long-term changes in quality of life following a brain injury, complications arising from traumatic brain injuries and their effects have garnered significant attention. Recently, there has been an increased focus on the prevalence of psychiatric illnesses as outcomes following traumatic brain injuries (17). This study investigates the mediating role of maladaptive and adaptive emotional regulation in the relationship between DERS, bullying behavior, and aggression among TBI patients.

The study demonstrates a significant correlation between DERS, aggression, and bullying behavior with cognitive mediation. Additionally, structural equation modeling analysis reveals that patients with maladaptive cognitive processes have difficulty in regulating their emotions, contributing to increased emotional turmoil. In general, both maladaptive and adaptive cognitive processes can predict difficulties in emotional control, aligning with previous research findings (18, 19). It appears that in the absence of adaptive regulation, patients may resort to impulsive and maladaptive regulation methods.

Furthermore, the findings indicate an indirect correlation between DERS and adaptive cognitive processes. Patients who do not struggle with adaptive cognitive regulation experience more positive emotions, higher self-esteem, and better psychological adaptation. Positive emotions serve as essential psychological resources, enabling individuals to employ coping strategies against psychological stress, maintain physical health, and confront stressful life events (20).

The role and impact of emotional regulation on various mental disorders have been explored in previous studies (21, 22). These studies have shown a connection between emotional regulation and anger. Individuals with stable emotional regulation tend to experience fewer negative emotions, such as anger and irritability. They manage their emotions before becoming overwhelmed and avoid situations that may trigger negative emotions.

Fable 2. Clinical Characteristics of the Participants				
Characteristics	Mean (SD, Range) %			
Bullying behavior	13.3 (0.63, 3.66)			
Maladaptive cognitive emotion regulation strategies (CERQ-M)	53.48 (11, 57)			
Adaptive cognitive emotion regulation strategies (CERQ-A)	70.1 (13.31, 67)			
Difficulties in Emotion Regulation Scale (DERS)	112.6 (21.8, 56.2)			
Aggregation	7.24 (5.87, 19)			

Table 3. Regression Coefficients for Model Paths

Paths	Non-standard Coefficients	Standard Coefficients	Standard Deviation	Critical Region	P-Value
CERQ-A $\rightarrow$ DERS	-1.151	-0.110	0.104	0.178	0.000
$DERS \rightarrow CERQ-M$	1.054	0.340	0.215	0.103	0.000
CERQ-M $\rightarrow$ C1	1.238	0.410	0.416	0.370	0.000
CERQ-M $\rightarrow$ C2	1.255	0.462	0.229	0.263	0.000
CERQ-M $\rightarrow$ C3	0.912	0.374	0.416	0.350	0.037
CERQ-M $\rightarrow$ C4	1.311	0.611	0.514	0.629	0.002
CERQ-M $\rightarrow$ C5	0.983	0.522	0.281	0.415	0.000
CERQ-A $\rightarrow$ C6	0.541	0.193	0.168	0.167	0.031
CERQ-A $\rightarrow$ C7	7.512	0.842	0.401	0.113	0.000
CERQ-A $\rightarrow$ C8	1.863	0.494	0.112	0.329	0.000
CERQ-A $\rightarrow$ C9	-0.712	-0.063	0.341	-0.563	0.110
DERS $\rightarrow$ D1	1.210	0.663	0.514	0.510	0.000
DERS $\rightarrow$ D2	1.110	0.843	0.203	0.563	0.000
DERS $\rightarrow$ D3	1.168	0.331	0.517	0.143	0.000
$DERS \rightarrow D4$	-0.074	-0.054	0.342	-0.329	0.231
DERS $\rightarrow$ D5	1.103	0.538	0.161	0.313	0.000
DERS $\rightarrow$ D6	-0.181	-0.121	0.109	-0.370	0.013
CERQ-M $ ightarrow$ Bullying behavior	1.311	0.441	0.154	0.629	0.002
CERQ-M $\rightarrow$ Aggregation	0.983	0.512	0.281	0.415	0.000
CERQ-A $\rightarrow$ Bullying behavior	-0.541	-0.123	0.168	0.617	0.031
CERQ-A $\rightarrow$ Aggregation	-7.512	-0.172	0.401	0.311	0.000

It has been revealed that people may face challenges in controlling their emotions for various reasons. This research suggests that the inability to control emotions can be linked to both compatible and incompatible cognitive-emotional adjustment components (23, 24). Consequently, enhancing emotional control in patients with brain injuries can involve identifying, modifying, and replacing components of emotional regulation. Furthermore, addressing the emotional aspects of traumatic brain injuries through psychological therapy can help patients overcome the challenges posed by this condition. This study is the first to examine the elements of both adaptive and incompatible cognitive-emotional regulation in traumatic brain injuries.

A primary limitation of our study was the limited sample size and the lack of similar studies that comprehensively addressed the generalizability of the results. Therefore, to strengthen the findings of this study, further research within the field is strongly encouraged.

Based on the study's results, it is recommended that psychological interventions be developed to teach emotion regulation strategies as part of the therapeutic protocol for TBI patients. This could help prevent the recurrence of traumatic events. Additionally,



Figure 1. Structural model of measuring the variable of emotion regulation based on the elements of emotional regulation.

research should be designed and conducted to assess the effectiveness of these interventions in TBI patients, aiming to determine the precise role of these components.

## Acknowledgments

The authors would like to thank the Shahid Mostafa Khomeini Hospital's Emergency Department staff for their help in gathering data (ethical code: IR.MEDILAM.REC.1396.64).

# Footnotes

**Authors' Contribution:** SE developed and designed the evaluation and collected the clinical data. NA drafted

the manuscript. JM and SE participated in the study conception and design, supervised the study, and revised the manuscript critically for important intellectual contents. KRA participated in data collection and writing the article. SE revised the manuscript critically for important intellectual contents. All the authors read and approved the final manuscript.

**Conflict of Interests:** The authors have no conflict of interest.

**Data Availability:** The dataset presented in the study is available on request from the corresponding author during submission or after publication.

**Ethical Approval:** This study is approved by the Ilam University of Medical Sciences (ethical code:

#### IR.MEDILAM.REC.1396.64).

**Funding/Support:** This study was partially supported by financial assistance from the Research Vice-Chancellor of Ilam University of Medical Sciences (No. 967008/82).

**Informed Consent:** The informed consent was recorded in the questionnaires.

#### References

- Schofield PW, Malacova E, Preen DB, D'Este C, Tate R, Reekie J, et al. Does traumatic brain injury lead to criminality? A whole-population retrospective cohort study using linked data. *PLoS One*. 2015;**10**(7). e0132558. [PubMed ID: 26172545]. [PubMed Central ID: PMC4501545]. https://doi.org/10.1371/journal.pone.0132558.
- Babaei F, Hosseini SH, Haddadi K. [Relationship between quality of life after traumatic brain injury and demographic characteristics in a three-year follow-up]. J Maz Univ Med Sci. 2021;30(192):77–86. Persian.
- Dalton MK, Jarman MP, Manful A, Koehlmoos TP, Cooper Z, Weissman JS, et al. Long-term healthcare expenditures following combat-related traumatic brain injury. *Mil Med*. 2022;**187**(3-4):513-7. [PubMed ID: 34173828]. https://doi.org/10.1093/milmed/usab248.
- Sirois K, Tousignant B, Boucher N, Achim AM, Beauchamp MH, Bedell G, et al. The contribution of social cognition in predicting social participation following moderate and severe TBI in youth. *Neuropsychol Rehabil.* 2019;29(9):1383–98. [PubMed ID: 29254438]. https://doi.org/10.1080/09602011.2017.1413987.
- Hoffer ME. Mild traumatic brain injury: Neurosensory effects. Curr Opin Neurol. 2015;28(1):74–7. [PubMed ID: 25502047]. https://doi.org/ 10.1097/WCO.00000000000164.
- Prus R, Appelhans O, Logash M, Pokotylo P, Nowicki GJ, Slusarska B. A histological and morphometric assessment of the adult and juvenile rat livers after mild traumatic brain injury. *Cells*. 2021;10(5). [PubMed ID: 34066539]. [PubMed Central ID: PMC8148537]. https:// doi.org/10.3390/cells10051121.
- Long K, Felton JW, Lilienfeld SO, Lejuez CW. The role of emotion regulation in the relations between psychopathy factors and impulsive and premeditated aggression. *Personal Disord*. 2014;5(4):390-6. [PubMed ID: 25198433]. [PubMed Central ID: PMC4198451]. https://doi.org/10.1037/per0000085.
- Sawchyn JM, Mateer CA, Suffield JB. Awareness, emotional adjustment, and injury severity in postacute brain injury. J Head Trauma Rehabil. 2005;20(4):301-14. [PubMed ID: 16030438]. https://doi.org/10.1097/00001199-200507000-00003.
- Henry JD, Phillips LH, Crawford JR, Ietswaart M, Summers F. Theory of mind following traumatic brain injury: the role of emotion recognition and executive dysfunction. *Neuropsychologia*. 2006;44(10):1623-8. [PubMed ID: 16643969]. https://doi.org/10.1016/j.neuropsychologia.2006.03.020.
- McDonald S, Rushby J, Li S, de Sousa A, Dimoska A, James C, et al. The influence of attention and arousal on emotion perception in adults with severe traumatic brain injury. *Int J Psychophysiol*. 2011;**82**(1):124–31. [PubMed ID: 21315778]. https://doi.org/10.1016/j.ijpsycho.2011.01.014.

- Abdi S, Taban S, Ghaemian A. Cognitive emotion regulation questionnaire: Validity and reliability of Persian translation of CERQ-36 item. *Procedia Soc Behav Sci.* 2012;32:2–7. https: //doi.org/10.1016/j.sbspro.2012.01.001.
- Rezapour M, Soori H, Khodakarim S. [Testing psychometric properties of the perpetration of bullying and victimization scales with olweus bullying questionnaire in middle schools]. Safety Promot Inj Prev. 2014;1(4):212–21. Persian.
- Stanford MS, Houston RJ, Mathias CW, Villemarette-Pittman NR, Helfritz LE, Conklin SM. Characterizing aggressive behavior. Assessment. 2003;10(2):183-90. [PubMed ID: 12801190]. https: //doi.org/10.1177/1073191103010002009.
- Romans L, Fresan A, Senties H, Sarmiento E, Berlanga C, Robles-Garcia R, et al. Validation of the Impulsive/Premeditated Aggression Scale in Mexican psychiatric patients. *Nord J Psychiatry*. 2015;**69**(5):397-402. [PubMed ID: 25541859]. https://doi.org/10.3109/08039488.2014. 994033.
- Firoozi M, zfeizabadi Z. [Behind the wall of violence: Psychological impairment of premeditated and impulsive aggression in compare to non - aggressive individuals in prison]. J Psychol Sci. 2019;18(81):1029–36. Persian.
- Shafiei E, Nademi A, Fakharian E, Omidi A. [An artificial neural network comparison with logistic regression in predicting post-traumatic mental disorders in mild brain injury patients]. *Thought Behav Clin Psychol.* 2017;12(46):37–46. Persian.
- 17. Nemattavousi M, Soltaninia S. The mediating role of experiential avoidance in the relationship between Alexithymia and emotion regulation in patients with major depression disorder after traumatic brain injury. *Shenakht J Psychol Psychiatry*. 2020;7(2):140–52. https://doi.org/10.52547/shenakht.7.2.140.
- Garnefski N, Kraaij V. Relationships between cognitive emotion regulation strategies and depressive symptoms: A comparative study of five specific samples. *Pers Individ Differ*. 2006;40(8):1659–69. https: //doi.org/10.1016/j.paid.2005.12.009.
- McGee R, Wolfe D, Olson J. Multiple maltreatment, attribution of blame, and adjustment among adolescents. *Dev Psychopathol.* 2001;13(4):827-46. [PubMed ID: 11771910].
- Nolen-Hoeksema S, McBride A, Larson J. Rumination and psychological distress among bereaved partners. J Pers Soc Psychol. 1997;72(4):855-62. [PubMed ID: 9108698]. https://doi.org/10.1037/ /0022-3514.72.4.855.
- 21. Roberton T, Daffern M, Bucks RS. Emotion regulation and aggression. Aggress Violent Behav. 2012;17(1):72–82. https://doi.org/10.1016/j.avb. 2011.09.006.
- Martin RC, Dahlen ER. Cognitive emotion regulation in the prediction of depression, anxiety, stress, and anger. *Pers Individ Differ*. 2005;**39**(7):1249–60. https://doi.org/10.1016/j.paid.2005.06.004.
- Ogden J. Health Psychology: A Textbook. UK: McGraw-Hill Education; 2012. 565 p.
- Tugade MM, Fredrickson BL. Resilient individuals use positive emotions to bounce back from negative emotional experiences. J Pers Soc Psychol. 2004;86(2):320–33. [PubMed ID: 14769087]. [PubMed Central ID: PMC3132556]. https://doi.org/10.1037/0022-3514.86.2.320.