



Effectiveness of an Internet-Based Transdiagnostic Unified Protocol Intervention for the Treatment of Depressive and Anxiety Disorders in Women with Threatened Abortion: A Feasibility Study

Seyyede Mahboobeh Mirtabar¹, Shahnaz Barat², Farzan Kheirkhah³, Amrollah Mostafazadeh⁴, Hoda Shirafkan³, Zeynab Pahlavan⁵, Mahbobeh Faramarzi^{3,*}

¹ Student Research Committee, Health Research Institute, Babol University of Medical Sciences, Babol, Iran

² Infertility and Reproductive Health Research Center, Health Research Institute, Babol University of Medical Sciences, Babol, Iran

³ Social Determinants of Health Research Center, Health Research Institute, Babol University of Medical Sciences, Babol, Iran

⁴ Cellular and Molecular Biology Research Center, Health Research Institute, Babol University of Medical Sciences, Babol, Iran

⁵ Department of Obstetrics and Gynecology, Rouhani Hospital, Babol University of Medical Sciences, Babol, Iran

*Corresponding author: Social Determinants of Health Research Center, Health Research Institute, Babol University of Medical Sciences, Babol, Iran. Email: mahbob330@yahoo.com

Received 2024 April 4; Revised 2024 July 20; Accepted 2024 September 14.

Abstract

Background: Despite the high prevalence of depressive and anxiety disorders among women with threatened abortion (TA), there is scant evidence regarding the efficacy of psychotherapies for this group

Objectives: This research aimed to assess the effectiveness, acceptability, and feasibility of specific internet-based cognitive behavioral therapy (iCBT-specific) and unified protocol (iUP) for women with a TA.

Methods: In this semi-experimental study (from April to October 2023), after the treatment content was prepared, a total of 80 pregnant women with TA, over 18 years of age, having a gestational age of 5 - 12 weeks, from a hospital and two private clinics, who were diagnosed with depressive and anxiety disorders based on the short structured diagnostic interview (Mini-International Neuropsychiatric Interview), were selected. The participants were divided into two groups, undergoing iUP or iCBT-specific psychotherapies. All participants completed the Brief Symptom Inventory (BSI-18) to measure the levels of depressive and anxiety symptoms. Following eight weeks of internet-based therapy, consisting of eight 50-minute sessions, they completed the BSI-18 again, along with the Client Satisfaction Questionnaire (CSQ-8) and the System Usability Scale (SUS). Statistical analysis was performed using SPSS version 22 software, and *t*-test, paired *t*-test, and ANCOVA tests were used.

Results: Both the iUP and iCBT-Specific models demonstrated high acceptability, and the average feasibility of treatment was above average for both models. In terms of acceptability and feasibility of treatment for women with TA, neither model proved to be superior (all $P > 0.05$). Both iUP and iCBT-Specific models were effective in reducing depression and anxiety in these women (effect size for within groups were respectively $\eta^2 = 0.370$, $\eta^2 = 0.536$). Also, in terms of improving depression and anxiety, there was no significant difference in the effectiveness of iUP and iCBT-Specific treatments (respectively $P = 0.146$, $\eta^2 = 0.027$ and $P = 0.221$, $\eta^2 = 0.019$)

Conclusions: The iUP treatment can be recognized as a reliable alternative to the iCBT-Specific method for alleviating depressive and anxiety symptoms in women with TA. Also, our results help gynecologists to use iUP as an alternative treatment when there is no definitive diagnosis of the type of disorder.

Keywords: Internet-Based Intervention, Transdiagnostic Unified Protocol, Anxiety, Depression, Threatened Abortion

1. Background

Threatened abortion (TA) is identified when an individual experiences vaginal bleeding, with or without accompanying abdominal pain during

pregnancy under 20 weeks (1). The clinical diagnosis is based on spotting or vaginal bleeding with a closed cervix in the first trimester of pregnancy (2). It could be a common complication of pregnancy, occurring in

approximately 20% of pregnant women. Nearly half of these cases result in a complete abortion (3). In a study, the number of all types of abortions in 2021 worldwide was estimated at 23 million (4), and TA affects about 7 to 24 percent of the population (5). Factors involved with TA include changing lifestyles and eating habits, increasing severe environmental pollution, increasing job and work stress among women and elderly pregnant women, and reducing the quality of oocytes (6). Evidence indicates that approximately 17% of women with TA encounter various complications in the later stages of pregnancy (7). Other various factors contribute to TA, including chromosomal abnormalities, gynecologic disorders, unhealthy lifestyle choices, and occupational stress (8).

Bleeding associated with TA can induce maternal anxiety, thereby elevating the risk of adverse fetal/maternal outcomes (9). Various studies have associated TA with an elevated risk of complications, such as preterm delivery, placenta previa, hypertension/preeclampsia, fetal growth restriction, and the need for neonatal intensive care unit (NICU) support (10, 11). Abortion is also a distressing experience that can affect a woman's emotional state and potentially lead to psychological disorders, such as depression (12). In a study carried out by Zhu et al., it was found that the incidence of major depression and anxiety symptoms was significantly higher in women experiencing TA compared to those with a stable pregnancy (13). In another study in Singapore, they reported that depression and anxiety were higher in women who were at risk of abortion, and TA may be a risk factor for antenatal depression (14). Steinberg et al. study on psychosocial and pre-abortion factors and mental health that the importance of stigma. Perceived abortion stigma appeared to explain the most additional variance in depression, anxiety, and stress symptoms. The relationship between the stigma of abortion and symptoms of depression, anxiety, and stress before abortion was strong and highly significant (15).

Psychotherapy can indeed serve as an effective intervention in managing the traumatic experience associated with TA. Evidently, providing psychological support is crucial for women undergoing this ordeal (16, 17). While numerous studies highlight the impact of psychotherapy on enhancing the mental well-being of women post-abortion (18, 19) there are limited reports on its impact on patients facing TA. A qualitative study

conducted in South Africa on pregnant women at risk of abortion revealed that counseling, raising awareness among these women, timely psychotherapy, and early diagnosis can contribute to improved childbirth outcomes (20). A study that examined the effectiveness of a happiness counseling program on depression, anxiety, and stress in pregnant women with a history of abortion showed that this non-pharmacological intervention improves the cognitive and emotional state of the patients and allows them to have a more positive attitude towards their abortion (21).

In recent decades, internet-based treatments have been employed to address a range of psychiatric disorders during pregnancy and the postpartum period, including depression and anxiety (22), as well as psychological issues arising postpartum (23). Numerous articles have confirmed the accessibility, feasibility, and acceptability of internet-based treatments for managing various disorders, even during pregnancy (24, 25). The advantages of internet-based cognitive behavioral therapy (iCBT) include easy and round-the-clock access to treatment, evasion of the stigma associated with in-person therapist meetings (26), preservation of privacy (27), increased adherence to therapeutic tasks (28), cost-effectiveness, elimination of geographical limitations (29), and patient anonymity (30). In Kheirkhah et al.'s study, it was found that iCBT is feasible and available for providing treatment to infertile patients. Also, this study confirmed that both iCBT and face-to-face CBT are equally effective in reducing patients' depression and anxiety (25).

The unified protocol (UP) is a transdiagnostic CBT approach that targets common psychopathological trauma related to emotional disorders and has a special emphasis on emotion regulation (31). The objective of integrated transdiagnostic treatment is to impart new adaptive strategies for emotion regulation, enabling patients to better tolerate intense emotions and enhance their quality of life, irrespective of their definitive diagnosis (32).

Despite the high prevalence of mental disorders, such as depression and anxiety, in women facing TA and the reported effectiveness of iCBT and internet-based UP (iUP) psychotherapies in various populations, the efficacy of these two models has not been compared in pregnant women. To the leading of our knowledge, this is the primary study to evaluate the feasibility and acceptability of iCBT-Specific and iUP psychotherapies in

women with TA, encountering depression and anxiety disorders.

2. Objectives

Our objectives were threefold: (1) to assess the acceptability and feasibility and evaluate the effectiveness of iCBT-Specific and iUP psychotherapies in women with TA, experiencing depression or anxiety disorders; (2) To compare the effectiveness of iCBT-Specific and iUP psychotherapies in improving depression and anxiety symptoms in women with TA.

3. Methods

This study was conducted in two stages. In the initial phase, we developed two treatment programs, including a transdiagnostic program and a specific iCBT program, for women with TA diagnosed with anxiety and depression. In the subsequent phase, we evaluated the effectiveness of these programs through a semi-experimental study.

3.1. Phase 1: Design of the Treatment Plan

The content of both iCBT-Specific treatment and iUP programs received approval from the Department of Psychiatry and Psychology at Babol University of Medical Sciences, Iran. Also, both programs were uploaded to a website (www.peacefulmindme.com).

3.1.1. iCBT-Specific Treatment

The standard iCBT-Specific treatment for depression and anxiety was grounded in the methodologies proposed by Beck and Beck (33) and Wright et al. (34), with a particular emphasis on psych education related to TA (including signs of TA and ways to control it, teaching the thought cycle and defining thought-feeling and behavior and rumination, identifying common cognitive distortions and their impact on the pregnancy process and the challenges ahead- Behavioral activation and coping behavior training and self-care).

The specific computer-based therapy was structured into eight sessions, each lasting 50 minutes. Each session was composed of multiple steps, which included videos featuring a mental health specialist. This specialist guided the patient and discussed various self-help strategies. Additionally, videos featuring a fictional character grappling with anxiety and depression were

developed. These videos were designed to help patients recognize their issues, work towards improving their condition, and understand that they are not alone in their struggle. The therapy sessions incorporated photos, texts, and educational PowerPoint presentations as tools to enhance understanding. During the sessions, we engaged the patients by asking questions about the content, aiming to focus their attention on it. Through exercises and assignments, the patients were encouraged to apply these strategies to their own problems. The treatment was guided weekly for 20 minutes over the phone by the first author.

3.1.2. Unified Protocol

The transdiagnostic treatment comprised of eight sessions, each lasting 50 minutes. This approach was adopted from the treatment protocol outlined by Barlow and colleagues, as presented in a book titled "Integrated Protocol for Transdiagnostic Treatment of Emotional Disorders (Therapist's Guide)" (35). Generally, the UP for the transdiagnostic treatment of emotional disorders is an intervention rooted in emotion-focused CBT. It is structured into five modules or "core" components, each of which is grounded in the elements of CBT with proven efficacy. These components are specifically designed to address negative emotions and adverse reactions to these emotions (36). The structure of the meetings is presented in Appendix 1. (skill of observing emotional experiences- Fostering awareness to recognize thought patterns- articulating the reciprocal relationship between cognition and emotion - execution of physical confrontation exercises appropriate for expectant mothers,...). The treatment was administered by the first author, who provided guidance over the phone for a duration of 20 minutes each week.

3.2. Phase 2: Conducting a Pilot Study to Determine the Effectiveness of Treatments

We carried out a semi-experimental investigation at Rohani Hospital (an inpatient hospital) and at two private outpatient obstetrics and gynecology clinics in Babol, a city in northern Iran. The study spanned from April to October 2023, encompassing 80 eligible participants, all of whom were diagnosed with depression and anxiety disorders and were included in the iCBT-Specific and transdiagnostic online UP programs.

3.3. Participants

Pregnant patients, both inpatient and outpatient, were chosen from a public hospital and two private obstetrics and gynecology clinics in Babol. The inclusion criteria were being over 18 years of age, having a gestational age of 5-12 weeks, and a diagnosis of anxiety disorders and depression based on the Mini-International Neuropsychiatric Interview (MINI) (37). Other inclusion criteria were access to a computer and the internet and an education level above elementary school. The exclusion criteria were severe mental disorders (e.g., bipolar disorders and schizophrenia), risk of suicide at the time of evaluation, drug use in the last three months, and experiencing psychotherapy or taking drug for the treatment of anxiety/depression in the last six months. Patients with serious mental disorders were referred to the specialized psychiatric clinics of the hospital for appropriate treatment.

G*Power software was used to determine the sample size taking into account the findings of a study by Shafierizi et al. (38). With an effect size of 0.14, an error level assumption of 0.05, and a power of 80%, the minimum required sample size for each group was estimated to be 40 individuals. Factoring in a 10% attrition rate, a total of 88 individuals were included in the study.

In total, the psychologist interviewed 192 pregnant women at risk of abortion. Out of these women, 104 did not meet the inclusion criteria (94 did not have disorders, six had severe psychiatric disorders, and four were receiving treatment at other medical centers). Consequently, 88 women were admitted into the treatment process. However, four participants from each group were later excluded from the analysis due to discontinuation of the sessions and failure to complete the post-test.

Following the interview, eligible women filled out the informed consent form. Participants from both groups then completed the Brief Symptom Inventory-18 (BSI-18) prior to the commencement of treatment, allowing for the assessment of depression and anxiety symptoms. After completing an eight-week course of 50-minute internet-based treatment sessions, the participants filled out the BSI-18 again, along with the Client Satisfaction Questionnaire (CSQ-8) and the System Usability Scale (SUS).

3.4. Measures

We used the Mini-International Neuropsychiatric Interview (MINI) to select patient TA with depression and anxiety disorder to participate in the study. The three instruments including, BSI-18, SUS, and CSQ-8 was used to assess the outcomes.

3.4.1. The Mini-International Neuropsychiatric Interview

It is a brief structured psychiatric interview designed as a diagnostic tool to identify psychiatric disorders according to DSM-5. Mini-international neuropsychiatric interview (M.I.N.I) can be done in less than 15 minutes. and has similar reliability and validity to similar longer instruments. Clinicians/researchers intending to use the M.I.N.I undergo special training before using the tool to ensure that the quality standard is maintained (37). According to NejatiSafa et al.'s study, the Persian version of M.I.N.I is an applicable, reliable, and valid tool for diagnosing some mental disorders. Test-retest reliability was very good ($\phi = 2$, Cramer's $V = 0.89$, $P < 0.0001$) (39).

3.4.2. Brief Symptom Inventory

The shortened form of BSI-53 contains 18 items from BSI-53. This test includes three subscales: Somatization, depression, and anxiety. Its total score is (maximum = 72). Higher scores indicate more severe psychiatric symptoms. The scoring of questions is based on a five-point Likert Scale (0 = not at all, 1 = slightly, 2 = moderately, 3 = slightly, 4 = severely). The BSI-18's psychometric properties have been utilized in numerous countries and languages (40). The Persian version of BSI-18 exhibits good validity and reliability. The test-retest reliability of Persian BSI-18 was reported as 0.81 (41).

3.4.3. Client Satisfaction Questionnaire

The CSQ-8 was developed by Lasern et al. (42). Client Satisfaction Questionnaire is the short form of a 31-item instrument that measures general satisfaction. This tool includes 8 Likert-type items with scores from 1 to 4 (1 indicates the lowest degree of satisfaction and 4 the highest).

3.4.4. System Usability Scale

The SUS, developed by Brooke, is a simple self-report scale to assess perceived ease of use (43). It consists of 10 items with a 5-point Likert Scale, from strongly agree to strongly disagree. To calculate the SUS score, the score of odd-numbered -is subtracted by 1, and the score of even-numbered -is calculated by subtracting the score from 5. By multiplying the total scores by 2.5, the overall is obtained. The total score can range from 0 (very poor usability) to 100 (excellent usability). We used the validated Persian SUS (44). The computed Cronbach's alpha for the Persian version of SUS was 0.79.

3.5. Statistical Analysis

SPSS software (SPSS 21, SPSS Inc., Chicago, IL, USA) was used for data analysis. Descriptive statistics were also used to check the frequency distribution and mean and standard errors of the mean. A paired *t*-test was also used within a group of psychotherapies. Analysis of covariance (ANCOVA) was used to compare changes in the outcomes in two groups of specific iCBTs versus iUP participation. *T*-tests were used to compare the satisfaction and feasibility of the treatment between iCBTs and iUP. A *P*-value < 0.05 was considered to be statistically significant.

4. Results

The demographic characteristics of the studied population are shown in Table 1. There was no significant difference between the iCBT-Specific and transdiagnostic iUP groups in terms of age (32.22 ± 6.28 vs. 30.47 ± 5.77 *P* = 0.199) and other demographic variables. The average age of the participants in this study is 31.35 ± 6.06 years old. Approximately 64% of the participants had an education level below a high school diploma, while 36% had a university education. The number of pregnancies did not significantly differ between the two groups (*P* = 0.357). The average gestational age was eight weeks, and there was no significant difference between (8.30 ± 1.97 vs. 9.02 ± 2.32 *P* = 0.137).

Table 2 provides a comparison of the acceptability and feasibility of the two treatment methods. The data indicated that both the iUP and iCBT-Specific models had high average acceptability scores (25.25 and 25.35 out of a total score of 32, respectively). Furthermore, the average feasibility of treatment was above average for both models (58.93 and 53.56 out of a total score of 100, respectively). The results of the *t*-test showed that there

is no significant difference between the two treatment models in terms of acceptability and feasibility (*P* > 0.05).

Table 3 provides a comparison of intra- and inter-group treatment outcomes for the two treatment methods. The results of the paired *t*-test and ANCOVA revealed that the iUP treatment significantly reduced depression (95% CI: 2.42 to 5.32) by 3.87, with an effect size of 0.370. It also reduced anxiety (95% CI: 3.32 to 5.47) by 4.40, with an effect size of 0.536, thereby significantly reducing TA in women. Similarly, the iCBT-Specific treatment significantly reduced depression by 2.42 (95% CI: 1.22 to 3.62) with an effect size of 0.370 and anxiety (95% CI: 1.91 to 4.28) by 3.10 with an effect size of 0.536 in women with TA.

In terms of the somatization sub-component, the iUP treatment was able to reduce somatization symptoms by 1.87 (95% CI: 0.83 to 2.91), while the iCBT-Specific treatment reduced them by 3.07 (95% CI: 1.96 to 4.18). As for the sum of sub-components, or total BSI, the iUP treatment significantly reduced psychological distress in women with TA by 8.39 (95% CI: 7.46 to 12.83), and the iCBT-Specific treatment reduced it by 8.60 (95% CI: 5.76 to 11.43). The ANCOVA comparison of the two treatment groups revealed no significant difference in the effectiveness of iUP and iCBT-Specific treatments in terms of improving depression (*P* = 0.146) or anxiety (*P* = 0.221).

5. Discussion

This study investigated the effectiveness, acceptability, and feasibility of two treatments, including a transdiagnostic and a diagnosis-specific method, aimed at alleviating depression and anxiety in women with TA. Both the iUP and iCBT-Specific models demonstrated high acceptability. Furthermore, the feasibility of both treatment models was found to be above average. However, when comparing the two models in terms of acceptability and feasibility among women with TA, neither model proved to be superior.

Given the lack of studies comparing the acceptability and feasibility of iUP and iCBT-Specific treatments, reference has been made to a study that reported on one of these treatments. In the research conducted by Shariat Panahi et al., emotional iCBT was utilized to treat anxiety in pregnant women and their spouses. The results of this study echoed our findings, indicating the high acceptability and feasibility of this treatment

Table 1. Characteristics of the Study Population ^a

Variables	Specific-iCBTs	iUnified Protocol	Overall (n = 80)	P-Value
Age	32.22 ± 6.28	30.47 ± 5.77	31.35 ± 6.06	0.199
Education;				0.329
Primary school	28 (70)	23 (57.5)	51 (63.75)	
≥ High School	12 (30)	17 (42.5)	29 (36.25)	
Gravid				0.357
1	13 (32.5)	17 (42.5)	30 (37.4)	
2	15 (37.5)	12 (30)	27 (33.8)	
≥ 3	12 (29)	11 (27.5)	23 (28.8)	
Gestational age weeks	8.30 ± 1.97	9.02 ± 2.32	8.66 ± 2.17	0.137

Abbreviations: Specific-iCBTs: cognitive behavioral therapy; iUP, unified protocol.

^a Values are expressed as mean ± SD or No. (%).

Table 2. Acceptability and Feasibility of the iUnified Protocol vs Specific-iCBTs ^a

Acceptability	Specific-iCBTs Mean (SD)	iUnified Protocol	P-Value
1. Quality of service	3.07 (0.26)	3 (00)	0.083
2. Kind of service wanted	3 (0.32)	3 (00)	1
3. Needs met	3.07 (0.34)	3.12 (0.33)	0.516
4. Would recommend to friend	3.07 (0.26)	3 (00)	0.083
5. Satisfaction with help received	3.25 (0.43)	3.15 (0.36)	0.269
6. Dealt with problems	3.42 (0.50)	3.52 (0.50)	0.377
7. Overall satisfaction	3.20 (0.40)	3.22 (0.42)	0.788
8. Would return to program	3.25 (0.43)	3.22 (0.42)	0.796
Total score (range 8 - 32)	25.35 (1.91)	25.25 (1.17)	0.779
Feasibility			0.063
Total score (range 1 - 100)	53.56 (11.69)	58.93 (13.72)	

Abbreviations: Specific-iCBTs, cognitive behavioral therapy; iUP, unified protocol.

^a Range of scores: Acceptability, 1- 4, total 8 - 32; feasibility 1 - 100%.

among both pregnant women and their spouses (mean CSQ-8 = 24.2, SD = 4.9) ($P = 0.008$) (24). Additionally, our findings align with those of another study implementing iCBT. This study also reported mean CSQ-8 scores of 25.06 with a standard deviation (SD) of 4.18 (25). Conversely, a different study investigating the use of online CBT for treating coronary artery disease reported a lower level of treatment acceptance compared to our study (45).

Quantitative and qualitative data from another study suggest that Web-based therapy is generally useful, although it has important advantages and disadvantages. In web-based treatments, participants enjoyed access, anonymity, and weekly reminders, as well as an introduction to the principles of CBT and mindfulness. However, this treatment was not sufficiently user-friendly, long-lasting, and compatible with smartphones. Parts of the content were not always

relevant or appropriate. Participants felt that it could be improved by having it in the form of a smartphone app and by making the content more concise and including different parenting styles (46).

In our study, the average feasibility scores were 53.56 (SD = 11.69) for the iCBT-Specific group and 58.93 (SD = 13.72) for the transdiagnostic iUP group. These scores are lower than those reported in another study, which had an average feasibility score of 67.07 (SD = 17.23) (25). It appears that in our study, due to the threat of abortion, the participants prioritized concerns related to their physical health and pregnancy over their mental distress. However, given the unique circumstances of these pregnancies (i.e., TA) and the necessity for complete rest, it seems that such online treatments could offer greater benefits for the patients. To substantiate this point, we can cite the study by Hantsoo et al., examining pregnant women, non-pregnant

Table 3. Comparison of the Effectiveness of Unified Protocol vs. Specific-iCBTs on the Improvement of Depressive and Anxiety Symptoms in Women with Threatened Abortion

Measure Group	Intervention, Mean (SD)	Pre-post Test	Within-Group Mean Diff (CI95%)	Eta Effect Size for Time	Between-Group P-Value	Between-Group Eta Effect Size
Depression				0.370	0.146	0.027
iUnified protocol	6.45 (5.35)	2.57 (2.92)	3.87 (2.42-5.32)			
Specific iCBTs	5.52 (4.32)	3.10 (2.77)	2.42 (1.22-3.62)			
Anxiety				0.536	0.221	0.019
iUnified protocol	7.85 (4.17)	3.45 (3.28)	4.40 (3.32-5.47)			
Specific iCBTs	6.87 (3.99)	3.77 (3.22)	3.10 (1.91-4.28)			
Somatization				0.357	0.254	0.017
iUnified protocol	5.05 (3.35)	3.17 (2.38)	1.87 (0.83-2.91)			
Specific iCBTs	5.90 (3.56)	2.82 (2.43)	3.07 (1.96-4.18)			
Total BSI				0.548	0.874	0.0003
iUnified protocol	19.35 (10.81)	9.20 (7.38)	8.39 (7.46-12.83)			
Specific iCBTs	18.30 (9.39)	9.70 (7.26)	8.60 (5.76-11.43)			

Abbreviations: Specific-iCBTs, cognitive behavioral therapy; iUP, unified protocol.

women, and men. Their results indicated that approximately one-third of the pregnant women selected some form of computer-based therapy as their preferred choice (47).

Several assumptions have been made to account for the high treatment satisfaction observed in our study. Firstly, the treatments provided in this study were delivered online with the support of a therapist. Prior research has demonstrated that internet-based treatments with therapist support tend to yield higher satisfaction levels compared to those without therapist support (48). Secondly, the task-oriented nature of the treatments in this study likely contributes to their high acceptability and effectiveness. There is substantial evidence suggesting that homework assignments are crucial to the success of CBT across a range of psychiatric conditions. Specifically, in the context of depression treatment with CBT, the adaptation of homework has been linked to significant clinical improvements and has been shown to predict a reduction in depressive symptoms (49).

Another observation from this study was that both iUP and iCBT-Specific programs were effective in reducing depression and anxiety in women facing TA, with effect sizes of 0.027 and 0.019, respectively. Furthermore, neither iUP nor iCBT-Specific treatment demonstrated superiority over the other in terms of alleviating depression and anxiety symptoms. A study conducted in Iran by Khairkhan et al., which focused on

infertile women and was published on the PeacefulMind website, found that iCBT was not less effective than traditional face-to-face CBT in terms of enhancing all outcome variables (25). Numerous systematic reviews and meta-analyses have drawn comparisons between the effectiveness of traditional face-to-face CBT and online CBT (both therapist-guided and self-guided) programs. They found that all three modalities demonstrated comparable effectiveness (27, 50).

Another study was conducted to investigate whether a UP remains equivalent to single disorder protocols (SDPs) in the treatment of anxiety disorders at 12-month follow-up. The results showed that the conditions of UP and SDP remained the same according to the degree of severity of the doctor's initial diagnosis in the 12-month follow-up. Furthermore, there were no significant differences between conditions in secondary outcomes at 12-month follow-up. The effect estimates of UP versus SDPs from baseline to 12-month follow-up was 0.13 (95% CI, -0.37 to 0.62) (51).

Additionally, a study conducted in Iran in 2022, titled "The Effectiveness of Online Integrated Transdiagnostic Treatment on Psychological Distress, Mindfulness, Hyper-Excitement, and Mental Capacity of Pregnant Women", found that the online integrated transdiagnostic treatment was effective in reducing psychological distress and negative meta-emotion. It also helped promote positive meta-emotion and psychological empowerment among the subjects.

However, it did not have a significant impact on mindfulness (52).

While all of the aforementioned studies, including our own, used different measures of anxiety and depression, it is important to note that there is no clear correlation between the number of treatment sessions, the interval between sessions, and the change in symptom scores. A meta-analysis examining the dose-response relationship in standard psychotherapy revealed that patients with less severe symptoms tended to respond to treatment more rapidly (53).

This study has certain limitations. Being a semi-experimental study, it necessitates further randomized controlled trials to validate the effectiveness and feasibility of online treatments. The limited sample size and absence of follow-up studies also constrain the scope of our findings. Also, the therapist's involvement in this study could have potentially influenced the results, suggesting the need for future studies to explore the efficacy of internet-only treatments without therapist support.

This study suggests potential applications. It indicated the possibility of substituting traditional psychological treatments with transdiagnostic treatments beyond diagnosis, thereby addressing a broad spectrum of mental disorders with high comorbidity rates. Furthermore, it underscores the advantages of online treatments, particularly for high-risk pregnant women.

5.1. Conclusions

The findings of our research suggest that online transdiagnostic treatment is comparably effective to specific diagnostic treatments, with satisfactory acceptability and accessibility. Given the significant benefits of UP treatment, our results support the use of iUP by psychiatrists and gynecologists as a reliable alternative to iCBT-specific psychotherapies. This could potentially enhance the treatment of depression and anxiety in women at risk of miscarriage.

Acknowledgements

The authors thank all the pregnant women who participated in this study.

Supplementary Material

Supplementary material(s) is available [here](#) [To read supplementary materials, please refer to the journal website and open PDF/HTML].

Footnotes

Authors' Contribution: M. F. and S. M. M.: Designed and conducted the project and writing the initial draft; S. B. and Z. P.: Reviewed the paper; H. S.: Analyzed the data; A. M. and F. K.: Contributed to the study conception, design, and supervision. All authors read and approved the final manuscript.

Clinical Trial Registration Code: None.

Conflict of Interests Statement: The authors declare that there is no conflict of interest.

Data Availability: The dataset presented in the study is available on request from the corresponding author during submission or after publication.

Ethical Approval: This study was approved by the Ethics Committee of Babol University of Medical Sciences, Babol, Iran with code [IR.MUBABOL.REC.1401.158](#).

Funding/Support: This study was financially supported by Babol University of Medical Sciences with number 724134501.

Informed Consent: All patients signed the informed consent form for this study.

References

1. Lee HJ, Park TC, Kim JH, Norwitz E, Lee B. The Influence of Oral Dydrogesterone and Vaginal Progesterone on Threatened Abortion: A Systematic Review and Meta-Analysis. *Biomed Res Int*. 2017;2017:3616875. [PubMed ID: [29392134](#)]. [PubMed Central ID: [PMC5748117](#)]. <https://doi.org/10.1155/2017/3616875>.
2. Garg M. Maternal and Fetal Complications And Outcome of Pregnancy In Patients Presenting With Threatened Abortion. *Int J Life Sci, Biotechnol Pharma Res*. 2024;13(1).
3. Yassaee F, Shekarriz-Foumani R, Afsari S, Fallahian M. The effect of progesterone suppositories on threatened abortion: a randomized clinical trial. *J Reprod Infertil*. 2014;15(3):147-51. [PubMed ID: [25202672](#)]. [PubMed Central ID: [PMC4138421](#)].
4. Hantoushzadeh S, Gargar OK, Jafarabady K, Rezaei MM, Asadi F, Eshraghi N, et al. Diagnostic value of neutrophil-to-lymphocyte and platelet-to-lymphocyte ratio to predict recurrent pregnancy loss and abortion; a systematic review and meta-analysis. *Immun Inflamm Dis*. 2024;12(3). e1210. [PubMed ID: [38506423](#)]. [PubMed Central ID: [PMC10953208](#)]. <https://doi.org/10.1002/iid3.1210>.
5. Valladares-Garrido MJ, Failoc-Rojas VE, Ichiro-Peralta C, Astudillo-Rueda D, Silva-Diaz H. Toxoplasma gondii Infection and Threatened Abortion in Women from Northern Peru. *Infect Dis Obstet Gynecol*.

- 2022;2022:1163655. [PubMed ID: 35978967]. [PubMed Central ID: PMC9377906]. <https://doi.org/10.1155/2022/1163655>.
6. Yuan S, Gao F, Xin Z, Guo H, Shi S, Shi L, et al. Comparison of the efficacy and safety of phloroglucinol and magnesium sulfate in the treatment of threatened abortion: A meta-analysis of randomized controlled trials. *Med (Baltimore)*. 2019;98(24):e16026. [PubMed ID: 31192955]. [PubMed Central ID: PMC6587576]. <https://doi.org/10.1097/MD.00000000000016026>.
 7. Barooti E, Rashidi Fakari F, Darvish S, Tavakoly N. The Comparison of CAI25 Levels in the Normal Pregnancy and Threatened Abortion. *J Obstetrics, Gynecol Cancer Res*. 2022;7(3):226-9. <https://doi.org/10.30699/jogcr.7.3.226>.
 8. Bu Z, Hu L, Su Y, Guo Y, Zhai J, Sun YP. Factors related to early spontaneous miscarriage during IVF/ICSI treatment: an analysis of 21,485 clinical pregnancies. *Reprod Biomed Online*. 2020;40(2):201-6. [PubMed ID: 31883882]. <https://doi.org/10.1016/j.rbmo.2019.11.001>.
 9. Sukur YE, Goc G, Kose O, Acmaz G, Ozmen B, Atabekoglu CS, et al. The effects of subchorionic hematoma on pregnancy outcome in patients with threatened abortion. *J Turk Ger Gynecol Assoc*. 2014;15(4):239-42. [PubMed ID: 25584033]. [PubMed Central ID: PMC4285213]. <https://doi.org/10.5152/jtgga.2014.14170>.
 10. Petriglia G, Palaia I, Musella A, Marchetti C, Antonilli M, Brunelli R, et al. Threatened abortion and late-pregnancy complications: a case-control study and review of literature. *Minerva Ginecol*. 2015;67(6):491-7. [PubMed ID: 25668506].
 11. Abd El-Raheem AE, Mohamed A, Elboghady A. Obstetric Outcomes in Women with Threatened Abortion. *Al-Azhar Int Med J*. 2022;0(0):0. <https://doi.org/10.21608/aimj.2022.106760.1665>.
 12. Taksande V, Gomase K. A Case Report On: Incomplete Abortion with Chronic Hypertension. *J Pharmaceutical Res Int*. 2021;664-9. <https://doi.org/10.9734/jpri/2021/v33i59A34317>.
 13. Zhu CS, Tan TC, Chen HY, Malhotra R, Allen JC, Ostbye T. Threatened miscarriage and depressive and anxiety symptoms among women and partners in early pregnancy. *J Affect Disord*. 2018;237:1-9. [PubMed ID: 29754019]. <https://doi.org/10.1016/j.jad.2018.04.012>.
 14. Astepe BS, Bosgelmez S. Antenatal Depression and Anxiety Among Women with Threatened Abortion: A Case-Control Study. *Gynecol Obstetrics Reproductive Med*. 2020;26(2):75-82. <https://doi.org/10.21613/gorm.2019.867>.
 15. Steinberg JR, Tschann JM, Furgerson D, Harper CC. Psychosocial factors and pre-abortion psychological health: The significance of stigma. *Soc Sci Med*. 2016;150:67-75. [PubMed ID: 26735332]. [PubMed Central ID: PMC4737478]. <https://doi.org/10.1016/j.socscimed.2015.12.007>.
 16. Wang N, Creedy DK, Zhang M, Lu H, Elder E, Allen J, et al. Designing a Needs-Oriented Psychological Intervention for Chinese Women Undergoing an Abortion. *Int J Environ Res Public Health*. 2022;20(1). [PubMed ID: 36613103]. [PubMed Central ID: PMC9819149]. <https://doi.org/10.3390/ijerph20010782>.
 17. Lewandowska M, Scott R, Meiksin R, Reiter J, Salaria N, Lohr PA, et al. How can patient experience of abortion care be improved? Evidence from the SACHA study. *Women's Health*. 2024;20:17455057241242700. <https://doi.org/10.1177/17455057241242675>.
 18. Barat S, Yazdani S, Faramarzi M, Khafri S, Darvish M, Rad MN, et al. The Effect of Brief Supportive Psychotherapy on Prevention of Psychiatric Morbidity in Women with Miscarriage: A Randomized Controlled Trial about the First 24-hours of Hospitalization. *Oman Med J*. 2020;35(3): e130. [PubMed ID: 32550017]. [PubMed Central ID: PMC7294535]. <https://doi.org/10.5001/omj.2020.48>.
 19. Wenzel A. Cognitive behavioral therapy for pregnancy loss. *Psychotherapy (Chic)*. 2017;54(4):400-5. [PubMed ID: 29251960]. <https://doi.org/10.1037/pst0000132>.
 20. Radithhalo E, Maputle S, Mothiba T. Vaginal Bleeding as a Pregnancy Problem during First Trimester at Polokwane Municipality, Limpopo Province. *J Hum Ecol*. 2017;51(1-2):73-9. <https://doi.org/10.1080/09709274.2015.11906896>.
 21. Elsharkawy NB, Mohamed SM, Awad MH, Ouda MMA. Effect of Happiness Counseling on Depression, Anxiety, and Stress in Women with Recurrent Miscarriage. *Int J Womens Health*. 2021;13:287-95. [PubMed ID: 33688267]. [PubMed Central ID: PMC7936698]. <https://doi.org/10.2147/IJWH.S283946>.
 22. Bright KS, Mughal MK, Wajid A, Lane-Smith M, Murray L, Roy N, et al. Internet-based interpersonal psychotherapy for stress, anxiety, and depression in prenatal women: study protocol for a pilot randomized controlled trial. *Trials*. 2019;20(1):814. <https://doi.org/10.1186/s13063-019-3897-z>.
 23. Dennis CL, Grigoriadis S, Zupancic J, Kiss A, Ravitz P. Telephone-based nurse-delivered interpersonal psychotherapy for postpartum depression: nationwide randomised controlled trial. *Br J Psychiatry*. 2020;216(4):189-96. [PubMed ID: 32029010]. <https://doi.org/10.1192/bjp.2019.275>.
 24. Shariatpanahi M, Faramarzi M, Barat S, Farghadani A, Shirafkan H. Acceptability, feasibility, and effectiveness of internet-based emotion-focused cognitive behavior therapy with husband participation for Iranian pregnant women with anxiety disorder. *Int J Systemic Ther*. 2023;34(3):167-84. <https://doi.org/10.1080/2692398X.2023.2174773>.
 25. Kheirkhah F, Faramarzi M, Shafierizi S, Chehrazhi M, Basirat Z. Preliminary examination of acceptability, feasibility, and effectiveness of internet-based cognitive behavioral therapy for treatment of depression and anxiety in infertile women. *Heliyon*. 2023;9(5): e15760. [PubMed ID: 37180939]. [PubMed Central ID: PMC10173620]. <https://doi.org/10.1016/j.heliyon.2023.e15760>.
 26. Rollman BL, Herbeck Belnap B, Abebe KZ, Spring MB, Rotondi AJ, Rothenberger SD, et al. Effectiveness of Online Collaborative Care for Treating Mood and Anxiety Disorders in Primary Care: A Randomized Clinical Trial. *JAMA Psychiatry*. 2018;75(1):56-64. [PubMed ID: 29117275]. [PubMed Central ID: PMC5833533]. <https://doi.org/10.1001/jamapsychiatry.2017.3379>.
 27. Andrews G, Basu A, Cuijpers P, Craske MG, McEvoy P, English CL, et al. Computer therapy for the anxiety and depression disorders is effective, acceptable and practical health care: An updated meta-analysis. *J Anxiety Disord*. 2018;55:70-8. [PubMed ID: 29422409]. <https://doi.org/10.1016/j.janxdis.2018.01.001>.
 28. Loughnan SA, Sie A, Hobbs MJ, Joubert AE, Smith J, Haskelberg H, et al. A randomized controlled trial of 'MUMentum Pregnancy': Internet-delivered cognitive behavioral therapy program for antenatal anxiety and depression. *J Affect Disord*. 2019;243:381-90. [PubMed ID: 30266030]. <https://doi.org/10.1016/j.jad.2018.09.057>.
 29. Hedman E, Ljotsson B, Kaldø V, Hesser H, El Alaoui S, Kraepelien M, et al. Effectiveness of Internet-based cognitive behaviour therapy for depression in routine psychiatric care. *J Affect Disord*. 2014;155:49-58. [PubMed ID: 24238951]. <https://doi.org/10.1016/j.jad.2013.10.023>.
 30. Rost T, Stein J, Lobner M, Kersting A, Luck-Sikorski C, Riedel-Heller SG. User Acceptance of Computerized Cognitive Behavioral Therapy for Depression: Systematic Review. *J Med Internet Res*. 2017;19(9): e309. [PubMed ID: 28903893]. [PubMed Central ID: PMC5617907]. <https://doi.org/10.2196/jmir.7662>.

31. Timulak L, Richards D, Bhandal-Griffin L, Healy P, Azevedo J, Connon G, et al. Effectiveness of the internet-based Unified Protocol transdiagnostic intervention for the treatment of depression, anxiety and related disorders in a primary care setting: a randomized controlled trial. *Trials*. 2022;**23**(1):721. [PubMed ID: 36045387]. [PubMed Central ID: PMC9429701]. <https://doi.org/10.1186/s13063-022-06551-y>.
32. Barlow DH, Farchione TJ, Fairholme CP, Ellard KK, Boisseau CL, Allen LB, et al. *Protocolo unificado para el tratamiento transdiagnóstico de los trastornos emocionales*. Madrid, Spain: Alianza Editorial; 2015.
33. Beck JS, Beck AT. *Cognitive Behavior Therapy, Third Edition: Basics and Beyond*. New York, US: Guilford Publications; 2020.
34. Wright JH, Brown GK, Thase ME, Basco MR, Gabbard GO. *Learning Cognitive-Behavior Therapy: An Illustrated Guide*. Washington, D.C: American Psychiatric Publishing; 2017.
35. Ehrenreich-May J, Kennedy SM, Sherman JA, Bilek EL, Buzzella BA, Bennett SM, et al. *Unified Protocols for Transdiagnostic Treatment of Emotional Disorders in Children and Adolescents: Therapist Guide*. Oxford, England: Oxford University Press; 2017. <https://doi.org/10.1093/med-psych/9780199340989.001.0001>.
36. Barlow DH, Farchione TJ, Bullis JR, Gallagher MW, Murray-Latin H, Sauer-Zavala S, et al. The Unified Protocol for Transdiagnostic Treatment of Emotional Disorders Compared With Diagnosis-Specific Protocols for Anxiety Disorders: A Randomized Clinical Trial. *JAMA Psychiatry*. 2017;**74**(9):875-84. [PubMed ID: 28768327]. [PubMed Central ID: PMC5710228]. <https://doi.org/10.1001/jamapsychiatry.2017.2164>.
37. Sheehan DV, Lecrubier Y, Sheehan KH, Amorim P, Janavs J, Weiller E, et al. The Mini-International Neuropsychiatric Interview (M.I.N.I.): the development and validation of a structured diagnostic psychiatric interview for DSM-IV and ICD-10. *J Clin Psychiatry*. 1998;**59** Suppl 20:22-33. quiz 34-57. [PubMed ID: 9881538].
38. Shafierizi S, Faramarzi M, Nasiri-Amiri F, Chehrazi M, Basirat Z, Khairkhah F, et al. Therapist-guided internet-based cognitive behavioral therapy versus face-to-face CBT for depression/anxiety symptoms in infertile women with adjustment disorders: A randomized controlled trial. *Psychother Res*. 2023;**33**(6):803-19. [PubMed ID: 36628473]. <https://doi.org/10.1080/10503307.2022.2158763>.
39. Nejatiasafa AA, Sharafi E, Motamed M, Mohammadjafari A, Etesam F, Shahmansouri N, et al. Persian Translation and Cultural Adaptation of Mini International Neuropsychiatric Interview and its Psychometric Properties. *Arch Iran Med*. 2022;**25**(5):294-9. [PubMed ID: 35943004]. <https://doi.org/10.34172/aim.2022.48>.
40. Grassi L, Caruso R, Mitchell AJ, Sabato S, Nanni MG. Screening for emotional disorders in patients with cancer using the Brief Symptom Inventory (BSI) and the BSI-18 versus a standardized psychiatric interview (the World Health Organization Composite International Diagnostic Interview). *Cancer*. 2018;**124**(11):2415-26. [PubMed ID: 29660109]. <https://doi.org/10.1002/cncr.31340>.
41. Akhavan Abiri F, Shairi MR. [Short Forms of Symptom Checklist (SCL): Investigation of validity & Reliability]. *Clin Psychol Personality*. 2020;**18**(1):137-62. FA. <https://doi.org/10.22070/cpap.2020.2929>.
42. Larsen DL, Attkisson CC, Hargreaves WA, Nguyen TD. Assessment of client/patient satisfaction: development of a general scale. *Eval Program Plann*. 1979;**2**(3):197-207. [PubMed ID: 10245370]. [https://doi.org/10.1016/0149-7189\(79\)90094-6](https://doi.org/10.1016/0149-7189(79)90094-6).
43. Brooke J. SUS: A quick and dirty usability scale. *Usability Evaluation Industry*. 1996;**189**(3):189-94.
44. Dianat I, Ghanbari Z, AsghariJafarabadi M. Psychometric properties of the persian language version of the system usability scale. *Health Promot Perspect*. 2014;**4**(1):82-9. [PubMed ID: 25097841]. [PubMed Central ID: PMC4122034]. <https://doi.org/10.5681/hpp.2014.011>.
45. Bendig E, Bauereiß N, Buntrock C, Habibović M, Ebert DD, Baumeister H. Lessons learned from an attempted randomized-controlled feasibility trial on "WIDeCAD" - An internet-based depression treatment for people living with coronary artery disease (CAD). *Internet Interventions*. 2021;**24**:100375. <https://doi.org/10.1016/j.invent.2021.100375>.
46. Ashford MT, Olander EK, Rowe H, Fisher JR, Ayers S. Feasibility and Acceptability of a Web-Based Treatment with Telephone Support for Postpartum Women With Anxiety: Randomized Controlled Trial. *JMIR Ment Health*. 2018;**5**(2). e19. [PubMed ID: 29678804]. [PubMed Central ID: PMC5938691]. <https://doi.org/10.2196/mental.9106>.
47. Hantsoo L, Podcasy J, Sammel M, Epperson CN, Kim DR. Pregnancy and the Acceptability of Computer-Based Versus Traditional Mental Health Treatments. *J Womens Health (Larchmt)*. 2017;**26**(10):1106-13. [PubMed ID: 28426287]. [PubMed Central ID: PMC5651968]. <https://doi.org/10.1089/jwh.2016.6255>.
48. Mehta S, Peynenburg VA, Hadjistavropoulos HD. Internet-delivered cognitive behaviour therapy for chronic health conditions: a systematic review and meta-analysis. *J Behav Med*. 2019;**42**(2):169-87. [PubMed ID: 30387008]. <https://doi.org/10.1007/s10865-018-9984-x>.
49. Tang W, Kreindler D. Supporting Homework Compliance in Cognitive Behavioural Therapy: Essential Features of Mobile Apps. *JMIR Ment Health*. 2017;**4**(2). e20. [PubMed ID: 28596145]. [PubMed Central ID: PMC5481663]. <https://doi.org/10.2196/mental.5283>.
50. Olthuis JV, Watt MC, Bailey K, Hayden JA, Stewart SH. Therapist-supported Internet cognitive behavioural therapy for anxiety disorders in adults. *Cochrane Database Syst Rev*. 2015;(3). CD011565. [PubMed ID: 25742186]. <https://doi.org/10.1002/14651858.CD011565>.
51. Eustis EH, Gallagher MW, Tirpak JW, Nauphal M, Farchione TJ, Barlow DH. The Unified Protocol compared with diagnosis-specific protocols for anxiety disorders: 12-month follow-up from a randomized clinical trial. *General Hospital Psychiatry*. 2020;**67**:58-61. <https://doi.org/10.1016/j.genhosppsych.2020.08.012>.
52. Goudarzi M, Gholamrezaei S, Amraei K. [The effectiveness of online unified transdiagnostic treatment on psychological distress, mindfulness, meta-emotion and psychological empowerment of pregnant women]. *Shenakht J Psychol Psychiatry*. 2021;**8**(1):88-105. FA. <https://doi.org/10.32598/shenakht.8.1.88>.
53. Robinson L, Delgadillo J, Kellett S. The dose-response effect in routinely delivered psychological therapies: A systematic review. *Psychother Res*. 2020;**30**(1):79-96. [PubMed ID: 30661486]. <https://doi.org/10.1080/10503307.2019.1566676>.