












Prevalence of Non-suicidal Self-Injury in the General Population in Iran: A Systematic Review and Meta-Analysis

Sakineh Javadian Koutanaei ¹, Romina Hamzehpour ¹, Azadeh Ramezani ¹, Mahjoobeh Betyar ¹, Angela Hamidia ¹, Sussan Moudi ¹, Alireza Azizi ¹, Fatemeh Alizadeh ¹, Armon Massoodi ^{1,*}

¹ Social Determinants of Health Research Center, Health Research Institute, Babol University of Medical Sciences, Babol, Iran

*Corresponding Author: Social Determinants of Health Research Center, Health Research Institute, Babol University of Medical Sciences, Babol, Iran. Email: armonmassodi2@gmail.com

Received: 17 December, 2024; Revised: 20 February, 2025; Accepted: 16 March, 2025

Abstract

Context: Non-suicidal self-injury (NSSI) is an intentional behavior without suicidal intent, recently recognized as an independent diagnostic entity in the diagnostic and statistical manual of mental disorders, fifth edition (DSM-5). Various studies indicate that NSSI is prevalent worldwide, but there are no reports on its prevalence among the Iranian population.

Evidence Acquisition: A systematic review of the literature was conducted using databases such as Scopus, PubMed/Medline, Web of Science, IMBIS, Google Scholar, Cochrane, and PsycInfo. Domestic databases, including Iran Medex, Mogiran, and SID, were also utilized. All available data until the end of June 2024 were reviewed. Data extraction was performed by two researchers, and study quality was assessed using the Newcastle-Ottawa Scale (NOS). Studies on NSSI in the general population of Iran that reported prevalence rates and accurate sample sizes were included. The heterogeneity of the studies was evaluated using the Cochran test and I^2 statistics. Additionally, a meta-regression analysis was conducted based on the year of study.

Results: A total of 2,180 articles were reviewed, with 17 articles ultimately selected and included in the study. The random-effects model estimated the pooled prevalence of NSSI in the general population at 16.51% (95% CI, 13.59 - 19.43). The prevalence of NSSI in women and men was 19.27% (95% CI, 13.31 - 25.24) and 14.74% (95% CI, 10.53 - 18.94), respectively. A correlation was found between the number of years since the study was conducted and the prevalence of NSSI (Reg Coef = 0.01, 95% CI: 0.003 to 0.020, $P = 0.011$); newer studies reported higher prevalence rates.

Conclusions: The present study revealed a high prevalence of NSSI in the general population, particularly in certain provinces and among women, indicating the need for specific prevention and treatment programs.

Keywords: Iran, Non-suicidal Self-Injury, Prevalence, Self-Harm Behavior

1. Context

The diagnostic and statistical manual of mental disorders, fifth edition (DSM-5), defines non-suicidal self-injury (NSSI) as intentional, socially unacceptable behavior in which an individual harms their body to relieve emotional distress (1, 2). This behavior includes actions such as cutting or burning the skin, often undertaken to alleviate emotions or attract attention. The motivations for this behavior typically include reducing mental pressure, controlling emotional distress, or escaping life problems (1). However, NSSI can become addictive and dangerous, potentially leading to suicide (3). Additionally, self-harm may result in

incorrect learning of stress management methods, increased feelings of guilt, depression, and ultimately, exacerbation of primary mental illnesses (4). Several factors, such as depression, stress, low self-esteem, family problems, peer influence, and sexual or physical abuse, are primary contributors to self-harm (5). Treatment for this behavior requires various approaches, including parent training, brief skills training, and behavioral therapies (6). Research indicates that NSSI has become more prevalent in recent decades, with a higher prevalence than suicide in some clinical and non-clinical groups (7). Studies have shown that NSSI is prevalent in more than 4% to 6% of the general adult population and between 5.7% and 8.4%

among soldiers (8, 9). The lifetime prevalence of NSSI is estimated at 5.9%, with 2.7% of individuals experiencing injuries more than five times. Furthermore, 0.9% reported NSSI in the past year (10). In Korean youth, NSSI prevalence ranges from 11% to 39%, exceeding actual suicide rates (11). Among medical students in Rawalpindi, NSSI prevalence was reported at 28.4%, with higher rates in males (54.5%) compared to females (45.4%) (12). In South Asia, self-harm behavior prevalence ranges from 3.2% to 44.8% in non-clinical populations and from 5% to 16.4% in clinical populations, indicating a significant burden (13). During the COVID-19 pandemic, the prevalence of NSSI in non-clinical samples reached 22.5%, with a higher prevalence of 32.40% in adolescents compared to adults (14, 15). A study in Iran reported that the prevalence of self-harm among Iranian youth varies between 4.3% and 40.5%, underscoring the importance of this behavior in mental health (16).

2. Objectives

Given the high costs associated with NSSI for the health system and its incompatibility with societal values and cultural standards, a more thorough investigation of NSSI is necessary. Although extensive research has been conducted in this field, there remains insufficient information about the prevalence and characteristics of NSSI in the Iranian population. Therefore, we conducted this systematic review and meta-analysis to determine the prevalence of NSSI in the Iranian population, aiming to provide appropriate and timely interventions in the future and to help manage this problem more effectively.

3. Evidence Acquisition

In the present study, a systematic review and meta-analysis regarding the prevalence of NSSI in the general population in Iran were conducted. The preferred reporting items for systematic reviews and meta-analyses (PRISMA) checklist was utilized for this study (17)

3.1. Search Strategy

The following [MeSH] terms were used across various databases: Prevalence, epidemiology, Iran, self-injurious behavior, and self-mutilation. International electronic databases included Scopus, PubMed/Medline, Web of Science, IMBIS, Cochrane, and PsycINFO. National databases used were Iran Medex, Mogiran, and SID. Searches encompassed all available data until the end of June 2024. A gray literature search was also conducted using Google Scholar. Keywords for both domestic and

international databases included self-destruction, self-mutilation, self-injury, intentional poisoning, self-punishment, self-hitting, self-immolation, self-harm, nail-biting, scratching, and self-cutting (Appendix 1 in Supplementary File). References in retrieved articles were also searched. EndNote X7 software was used to manage the collected information, and duplicate articles were automatically removed. Articles were examined separately by two researchers.

3.2. Eligibility Criteria

We included in this study all observational (cross-sectional) studies that address NSSI in the general population linked to Iran, without imposing a publication time limit. The study population consisted of both men and women. All included studies were required to report NSSI rates in the general population. Several types of studies were excluded, including case reports, case series, reviews, and meta-analyses. Additionally, studies that did not report sample sizes were excluded from this meta-analysis. It should be noted that because suicide attempts are considered distinct from NSSI (18, 19), we also excluded studies that categorized suicide attempts as NSSI to ensure a pure prevalence of NSSI. The term "general population" refers to random sampling from a non-clinical and non-self-harming group within the normal community. Participants did not have any of the following risk factors: Psychiatric diseases (such as depression), hospitalization in service organizations (such as welfare), stressful conditions (such as living on the streets), or family problems (such as divorce).

3.3. Quality Appraisal

Article quality was assessed using the Newcastle-Ottawa Scale (NOS), which includes three parts: Selection (4 questions), comparability (1 question), and outcome (3 questions). The final score is categorized as good quality (three or four stars in selection, one or two stars in comparability, and two or three stars in outcome), fair quality (2 stars in selection, one or two stars in comparability, and two or three stars in outcome/exposure), and poor quality (0 or 1 stars in selection, 0 stars in comparability, or 0 or 1 stars in outcome/exposure) (20). Appendix 2 in Supplementary File presents the qualitative assessment results.

3.4. Screening of Studies

Two individuals conducted the initial search for studies. A team of two researchers independently screened studies, extracted results, and evaluated the

quality control of articles. In cases of disagreement, the team leader provided the final decision.

3.5. Data Extraction

All final articles were selected according to a pre-prepared checklist, which included author names, publication years, study periods, sample sizes, study locations, gender, average age, NSSI types, tools used, and prevalence of NSSI.

3.6. Statistical Analysis

Statistical analysis was conducted using STATA 14.0 (Stata Corporation, College Station, TX) software. The random-effects model was used to estimate the overall prevalence of NSSI, accounting for heterogeneity among the included studies. This model assumes that true effect sizes vary across studies, enabling more accurate estimation despite differences in study design and population characteristics. Heterogeneity was evaluated using Cochran's test and Higgins I^2 test. Forest plots illustrated the effect size of each study and the pooled estimates. The term 'General' refers to populations including both women and men, with overall NSSI prevalence reported for the entire population. Studies focusing solely on a specific gender were analyzed separately for NSSI prevalence by gender. Meta-regression and subgroup analysis were conducted, with significant results defined as those with a p-value below 0.05.

3.7. Publication Bias

Publication bias was evaluated using funnel plots and Egger's weighted regression (21). A P-value > 0.05 indicated no publication bias.

4. Results

4.1. Study Selection

A total of 3504 articles were identified through searches of all national and international databases. After removing duplicate articles, 2180 articles proceeded to the title and abstract screening stage. Subsequently, 64 full-text articles were evaluated, and 17 articles were ultimately selected for analysis. These 17

articles were also reviewed for related studies based on their references. Figure 1 illustrates the study selection process.

4.2. Study Characteristics

Among the 17 articles spanning from 2004 to 2024, 16 studies reported the prevalence of NSSI in the general population (both sexes), 7 studies reported prevalence in women, and 6 studies reported prevalence in men. Table 1 presents the descriptive information.

4.3. Quality Appraisal

Article quality assessment results are shown in Appendix 2 in Supplementary File. According to the review using the relevant scale, all studies were rated as having good quality.

4.4. Heterogeneity

The chi-squared test and I^2 Index indicated significant heterogeneity among the studies. In the general population, the heterogeneity of NSSI prevalence was $I^2 = 98.9\%$ ($P < 0.001$); in men, it was $I^2 = 95.6\%$ ($P < 0.001$); and in women, it was $I^2 = 98.4\%$ ($P < 0.001$). Consequently, all analyses were conducted using the random-effects model.

4.5. Meta-Analysis Results

The articles were first sorted by publication year, followed by determining the prevalence of NSSI in both genders and provinces. A meta-regression was also performed based on the study year.

4.6. Prevalence of Non-suicidal Self-Injury in the General Population and by Gender

Sixteen studies reported the prevalence of NSSI in the general population out of a total of 17 articles. Based on the random-effects model, the estimated pooled prevalence of NSSI was 16.51% (95% CI, 13.59 - 19.43) (Figure 2). Seven studies reported the estimated pooled prevalence of NSSI in women as 19.27% (95% CI, 13.31 - 25.24) (Figure 3), and six studies reported the estimated pooled prevalence of NSSI in men as 14.74% (95% CI, 10.53 - 18.94) (Figure 4).




Figure 1. Preferred reporting items for systematic reviews and meta-analyses (PRISMA) 2020 flow diagram for new systematic reviews, including database and other source searches (17)

4.7. Meta-Analysis of Subgroups According to Provinces

According to the prevalence of NSSI by province, Tehran province had six studies with an estimated pooled prevalence of 25.54% (95% CI, 14.48 - 36.80), followed by West Azerbaijan province with five studies at 7.10% (95% CI, 4.41 - 9.79), Gilan province with two studies at 11.83% (95% CI, 9.90 - 13.76), and Mazandaran, Fars, and Kurdistan provinces each had one study (Table 2) (Appendix 3 in Supplementary File).

4.8. Meta-regression According to the Years of the Study

The estimated pooled prevalence of NSSI in the general population was related to the years in which the studies were conducted (Reg Coef = 0.01, 95% CI: 0.003 to 0.020, $P = 0.011$). This indicates that as the years progressed from the earlier years (2004) to the more recent years (2024), the estimated pooled prevalence of NSSI in the general population significantly increased (Figure 5).

4.9. Publication Bias

Finally, funnel plots were drawn to assess publication bias for the prevalence of NSSI in the general population. This bias was not confirmed by the Egger test (bias for total: -0.73, 95% CI = -12.82 to 11.34; $P = 0.897$) (Appendix 4 in Supplementary File).

5. Discussion

In this study, the estimated pooled prevalence of NSSI in the general population of Iran was calculated as 16.51%. This prevalence was higher in women (19.27%) than in men (14.74%), highlighting the importance of gender differences in this behavior. Various studies worldwide have reported a high prevalence of NSSI, ranging from 17% to 38% (38-42). As suggested by Ghaedi Heidari et al. (43), this gender difference may be attributed to distinct coping mechanisms employed by men and women in response to stress and psychological difficulties. Women may be more likely to express emotional distress through self-harming behaviors, whereas men may employ different, potentially less visible, coping strategies. However, the study by Izadi-Mazidi et al. in Tehran did not find significant gender differences in the prevalence of NSSI, indicating that in some contexts, gender may not be the sole determining factor for self-harm behavior (27). This discrepancy emphasizes the need for region-specific research that accounts for local socio-cultural conditions, which may mediate or mitigate gender differences in self-injury rates.

Furthermore, a 2023 review study corroborates our findings, demonstrating that the global prevalence of NSSI varies widely, ranging from 11.5% to 33.8%, influenced by sample types and study designs. This review also observed a rising trend in NSSI, particularly

Table 1. Basic Information of Included Studies in the Systematic Review and Meta-Analysis

Authors	Duration of Data Collection	Province (City)	Sex	Instrument for Self-Injury Findings	Type of Self-Injury	Total Sample Size
Abbasian et al. (22)	2018 - 2019	Markazy (Saveh)	Female	Standard instrument	NSSI	571
Abdollahi et al. (23)	2018 - 2018	Gilan (Rasht)	All sexes	Researcher checklist	Self-cutting	617
Babaeifard et al. (24)	2024 - 2024	Tehran (Tehran)	All sexes	Standard instrument	NSSI	1014
Ghanizadeh and Shekoohi (25)	2008 - 2008	Fars (Shiraz)	All sexes	Researcher checklist	Nail biting	460
Gholamrezaei et al. (26)	2017 - 2017	Tehran (Tehran)	All sexes	Standard instrument	NSSI	554
Izadi-Mazidi et al. (27)	2018 - 2018	Tehran (Tehran)	All sexes	Standard instrument	NSSI	642
Marin et al. (28)	2017 - 2018	Western Azerbaijan (Tabriz)	All sexes	Researcher checklist	NSSI	6229
Mohammadpoorasl et al. (29)	2005 - 2005	Western Azerbaijan (Tabriz)	All sexes	Researcher checklist	NSSI	1772
Mohammadpoorasl et al. (30)	2010 - 2010	Western Azerbaijan (Tabriz)	All sexes	Researcher checklist	NSSI	4903
Mozafari et al. (31)	2018 - 2019	Kurdistan (Sanandaj)	All sexes	Standard instrument	NSSI	1334
Nemati et al. (16)	2017 - 2017	Western Azerbaijan (Tabriz)	All sexes	Researcher checklist	NSSI	3966
Nobakht and Dale (32)	2016 - 2016	Mazandaran (Babol)	All sexes	Standard instrument	NSSI	200
Mohammadpoorasl et al. (33)	2005 - 2005	Western Azerbaijan (Tabriz)	All sexes	Researcher checklist	NSSI	1768
Rezaei et al. (34)	2020 - 2021	Tehran (Tehran)	All sexes	Standard instrument	NSSI	655
Sabet Dizkuhi and Kafie Masooleh (35)	2019 - 2020	Gilan (Rasht)	All sexes	Standard instrument	NSSI	508
Taheri et al. (36)	2019 - 2020	Tehran (Tehran)	All sexes	Standard instrument	NSSI	223
Bathae and Kamali Fard (37)	2004 - 2004	Tehran (Tehran-Rayy)	All sexes	Researcher checklist	Nail biting	18000

Abbreviation: NSSI, non-suicidal self-injury.

in developing countries, which aligns with the findings of our study. While NSSI is prevalent in both developed and developing countries, the socio-cultural and economic factors contributing to this behavior may differ across regions (44). In developed Western countries, the focus is often on individual mental health conditions, whereas in developing nations like Iran, a complex interplay of socio-economic stressors, cultural taboos, and limited access to mental health care may contribute to the growing prevalence (45).

The increasing trend of NSSI in adolescents, as reported by Sahay and Nilanjana and seen in other studies, underscores the importance of addressing the psychological and social factors driving this behavior, particularly in younger populations (46). The lack of adequate mental health services, high levels of economic stress, and societal pressures could exacerbate these issues, especially in countries like Iran, where social stigma around mental health may deter individuals from seeking help. Therefore, the findings of

our study call for targeted preventive and therapeutic interventions that are culturally sensitive and socioeconomically appropriate.

It should be noted that the socio-cultural and economic landscape of Iran plays a pivotal role in the prevalence and nature of NSSI. The societal emphasis on conformity and the stigma associated with mental health issues often prevent individuals from openly addressing psychological distress. Women, in particular, may be more susceptible to NSSI as a means of coping with the societal pressure to fulfill traditional roles, which may heighten emotional and psychological burdens. Economic factors, such as unemployment, financial insecurity, and the rising cost of living, further compound mental health challenges, particularly among vulnerable populations. In regions like Tehran, where urban stressors are more pronounced, the prevalence of severe self-harm methods, such as cutting, is higher, reflecting the intensifying pressures faced by individuals in urban settings. In contrast, rural areas,




Figure 2. Forest plot of the prevalence of self-injury in the general population of Iran (16, 23-37)




Figure 3. Forest plot of the prevalence of self-injury in the general population of Iran (female) (16, 22, 27, 28, 31, 32, 35)

like Gilan, may report lower rates of severe self-harm, which could be attributed to a combination of less exposure to mental health stigma, fewer healthcare resources, and different coping mechanisms prevalent in more rural, community-oriented environments. Moreover, the limited availability of mental health services, coupled with the growing economic and social challenges in Iran, exacerbates the situation,

contributing to the underreporting and lack of early interventions for NSSI. Therefore, expanding access to mental health care, reducing stigma, and addressing the socio-economic conditions influencing self-harm behaviors are crucial steps toward improving mental health outcomes in Iran.

In line with the present study, the study by Akbari et al. in 2024 also showed that the prevalence of NSSI is




Figure 4. Forest plot of the prevalence of self-injury in the general population of Iran (male) (16, 27, 28, 31, 32, 35)

Table 2. Results of the Meta-Analysis and Heterogeneity of Self-Injury Estimated Pooled Prevalence in the General Population of Iran Based on Province

Provinces	No. of Study	Effect Estimates (Estimated Pooled Prevalence)	95 % CI	I ²	P-Value
Tehran	6	25.64	14.48 - 36.80	99.2	< 0.001
Western Azerbaijan	5	7.10	4.41 - 9.79	98.1	< 0.001
Gilan	2	11.83	9.90 - 13.76	3.9	0.078
Mazandaran	1	38.50	31.75 - 45.24	-	-
Fars	1	14.00	10.82 - 17.17	-	-
Kurdistan	1	8.39	6.90 - 9.88	-	-
Total	16	16.51	13.59 - 19.43	98.9	< 0.001

higher in women than in men (47). In another study conducted in Iran, the prevalence of NSSI in male and female adolescents was reported as 26.8% and 17.9%, respectively (48). Evidence shows that this behavior is more common in girls. These differences could be related to psychological, cultural, and social factors (49-52). In a study conducted by Abbasian et al. in Iran in 2021, the prevalence of NSSI in women was 17.51%, higher than in men (22). A prevalence rate of 27.6% was reported in European countries. The most prevalent NSSI incidences were reported in Estonia, France, Germany, and Israel, while the lowest prevalence rates were reported in Hungary, Ireland, and Italy (53). It was more in women than in men, which is consistent with the findings of this study. These differences in NSSI prevalence between genders may be influenced by various psychological and social factors. Women are

generally more likely to internalize emotions and distress, which can lead to higher rates of self-harming behaviors. Additionally, gender roles and societal pressures may contribute to this disparity, with women often facing more emotional stress due to expectations related to caregiving and emotional expression. Social factors, such as gender inequality and higher exposure to interpersonal violence, can also increase the risk of NSSI in women (40). Further research is needed to explore how these factors interact to contribute to the gender differences observed in NSSI prevalence.

As observed in this study, regional variations in the prevalence of NSSI across Iran are significant. Tehran had the highest prevalence rate at 25.64%, while provinces like Western Azerbaijan and Gilan reported lower figures. These differences are likely influenced by various cultural, economic, and social factors that merit




Figure 5. Meta-regression of self-injury prevalence in the general population based on year of study

further investigation. For example, a study by Babaeifard et al. reported an exceptionally high NSSI prevalence of 41.91% in Tehran's general population (24). In contrast, studies by Mohammadpoorasl et al. and Mozafari et al. found much lower prevalence rates of under 10% in Western Azerbaijan and Kurdistan (31, 33). These discrepancies may stem from regional differences in healthcare access, social support systems, cultural attitudes towards mental health, and other socio-economic factors. Further research is essential to understand these regional differences more comprehensively and to develop targeted prevention and intervention strategies.

Moreover, the type of self-harm methods used may also contribute to these regional variations. Although data on self-harm methods were available from only three studies, certain patterns emerged. Specifically, Ghanizadeh and Shekoohi (25) and Bathaee and Kamali Fard (37), both conducted in Tehran and Fars, reported higher NSSI prevalence compared to Abdollahi et al. (23), which focused on self-cutting in Gilan. These findings suggest that more severe self-harm methods, such as cutting, are associated with higher prevalence rates of NSSI. Geographic factors, including access to mental health care, healthcare infrastructure, and cultural attitudes, may further explain the observed differences

in both self-harm methods and NSSI prevalence. Additionally, socio-economic conditions in each province could influence not only the occurrence of NSSI but also the choice of self-harm methods. For instance, urban areas like Tehran may experience more severe forms of self-harm due to higher stress levels, greater availability of means, and distinct cultural norms compared to rural areas such as Gilan. More research is needed to explore the relationship between geographical conditions, self-harm methods, and NSSI prevalence, which will facilitate the development of more effective, region-specific interventions.

5.1. Conclusions

Self-injury is considered a serious mental health issue in Iran, requiring special attention. The high prevalence of this behavior, especially in some provinces and among women, indicates the necessity of designing and implementing preventive and therapeutic programs that focus specifically on these groups. The increase in NSSI prevalence in new studies also indicates that more research should be done to identify the reasons for its increasing trend. This should be done to design timely interventions.

5.2. Future Research Directions

Given the high prevalence of NSSI in the Iranian population, particularly among women and in certain provinces, it is crucial that future research explores the psychological, social, and cultural factors contributing to this behavior. Longitudinal studies are needed to assess the long-term impacts of interventions aimed at reducing NSSI. Additionally, further investigation into the increasing prevalence of NSSI, especially in light of recent societal changes, will help inform the development of targeted prevention and treatment programs.

5.3. Strengths and Weaknesses

This study used meta-analysis to combine different data, which increased the accuracy of the results. Additionally, the geographical diversity of this study provides a more comprehensive viewpoint of this behavior in the country. Furthermore, women and men have been analyzed separately for differences in the prevalence of NSSI. This information may help future treatment and prevention plans to be more effective. However, the number of studies in some provinces was small and may not be a proper representative of the prevalence rate of NSSI in those regions. For example, some provinces had only one or two studies, and some provinces had none. The NSSI prevalence was correlated with the year of study, but changes in social and psychological conditions during these years were not fully explored. Therefore, considering these weaknesses, it is suggested to collect more data from less studied or unstudied provinces and analyze social and psychological changes in more detail in future research. By using these measures, we can better understand the factors affecting NSSI prevalence and improve therapeutic interventions.

Acknowledgements

The study's research proposal received approval from the Research Ethics Committee at Babol University of Medical Sciences (IR.MUBABOL.HRI.REC.1403.187).

Supplementary Material

Supplementary material(s) is available [here](#) [To read supplementary materials, please refer to the journal website and open PDF/HTML].

Footnotes

Authors' Contribution: S. J. K., R. H., and A. M. conceived and designed the evaluation and drafted the manuscript. A. R., A. A., and M. B. participated in designing the evaluation, performed parts of the statistical analysis and helped to draft the manuscript. A. H., S. M., and F. A. A. re-evaluated the clinical data, revised the manuscript and performed the statistical analysis and revised the manuscript. S. M. and A. M. collected the clinical data, interpreted them and revised the manuscript. S. J. K., F. A. A., and A. M. re-analyzed the clinical and statistical data and revised the manuscript. All authors read and approved the final manuscript.

Conflict of Interests Statement: The authors declare no conflict of interest.

Data Availability: The dataset presented in the study is available on request from the corresponding author during submission or after publication. The data are not publicly available due to confidentiality.

Funding/Support: The present study received no funding/support.

References

1. Xiao Q, Song X, Huang L, Hou D, Huang X. Global prevalence and characteristics of non-suicidal self-injury between 2010 and 2021 among a non-clinical sample of adolescents: A meta-analysis. *Front Psychiatry*. 2022;**13**:912441. [PubMed ID: 36032224]. [PubMed Central ID: PMC9399519]. <https://doi.org/10.3389/fpsy.2022.912441>.
2. Owens R, Filoromo SJ, Landgraf LA, Lynn CD, Smetana MR. Deviance as an historical artefact: a scoping review of psychological studies of body modification. *Humanities Soc Sci Commun*. 2023;**10**(1). <https://doi.org/10.1057/s41599-023-01511-6>.
3. Nada-Raja S, Skegg K, Langley J, Morrison D, Sowerby P. Self-harmful behaviors in a population-based sample of young adults. *Suicide Life Threat Behav*. 2004;**34**(2):177-86. [PubMed ID: 15191274]. <https://doi.org/10.1521/suli.34.2.177.32781>.
4. Dahlby L. *Mediators of Childhood Maltreatment and Suicidal Ideation versus Attempts: A Systematic Review and Research Proposal [Dissertation]*. Edinburgh, UK: University of Edinburgh; 2022.
5. Lee WK. Psychological characteristics of self-harming behavior in Korean adolescents. *Asian J Psychiatr*. 2016;**23**:119-24. [PubMed ID: 27969068]. <https://doi.org/10.1016/j.ajp.2016.07.013>.
6. Glenn CR, Esposito EC, Porter AC, Robinson DJ. Evidence Base Update of Psychosocial Treatments for Self-Injurious Thoughts and Behaviors in Youth. *J Clin Child Adolesc Psychol*. 2019;**48**(3):357-92. [PubMed ID: 31046461]. [PubMed Central ID: PMC6534465]. <https://doi.org/10.1080/15374416.2019.1591281>.
7. Baetens I, Claes L, Muehlenkamp J, Grietens H, Onghena P. Non-suicidal and suicidal self-injurious behavior among Flemish adolescents: A web-survey. *Arch Suicide Res*. 2011;**15**(1):56-67. [PubMed ID: 21294000]. <https://doi.org/10.1080/13811118.2011.540467>.
8. Klonsky ED, Glenn CR. Resisting Urges to Self-Injure. *Behav Cogn Psychother*. 2008;**36**(2):211-20. [PubMed ID: 29527120]. [PubMed Central ID: PMC5841247]. <https://doi.org/10.1017/S1352465808004128>.
9. Soltaninejad A, Fathi Ashtiani A, Ahmadi K, Azad Fallah P, Anisi J, Rahmati Najjar Kalayi F, et al. The Relationship Between Religious

- Orientation and Suicidal Behaviors in Soldiers. *Islamic Lifestyle Centered Health*. 2012;**1**(2):22-7. <https://doi.org/10.5812/ilch.9019>.
10. Klonsky ED. Non-suicidal self-injury in United States adults: prevalence, sociodemographics, topography and functions. *Psychol Med*. 2011;**41**(9):1981-6. [PubMed ID: 21208494]. <https://doi.org/10.1017/S0033291710002497>.
 11. Yoo HJ. Agony of the Society, Non-Suicidal Self-Injury. *Soa Chongsinyon Chongsin Uihak*. 2023;**34**(3):151. [PubMed ID: 37426828]. [PubMed Central ID: PMC10326350]. <https://doi.org/10.5765/jkacap.230042>.
 12. Mehmood M, Awan NU, Fiaz T, Raza SA. Prevalence of non-suicidal self-injury in medical students of Rawalpindi; its socio-demographics, methods, and functions. *J Pak Med Assoc*. 2023;**73**(12):2370-4. [PubMed ID: 38083914]. <https://doi.org/10.47391/jpm.7544>.
 13. Haregu T, Chen Q, Arafat SMY, Cherian A, Armstrong G. Prevalence, correlates and common methods of non-suicidal self-injury in South Asia: a systematic review. *BMJ Open*. 2023;**13**(11). e074776. [PubMed ID: 37993150]. [PubMed Central ID: PMC10668266]. <https://doi.org/10.1136/bmjopen-2023-074776>.
 14. Deng H, Zhang X, Zhang Y, Yan J, Zhuang Y, Liu H, et al. The pooled prevalence and influential factors of non-suicidal self-injury in non-clinical samples during the COVID-19 outbreak: A meta-analysis. *J Affect Disord*. 2023;**343**:109-18. [PubMed ID: 37802326]. <https://doi.org/10.1016/j.jad.2023.09.036>.
 15. Ghorbani A, Kamali Z, Ebrahimi P, Abodollahi Z, Minaei M, Movahedi A. The Effects of Quarantine and Corona Virus on Dietary Habits, Physical Activity, and Anthropometric Indices. *J Nutrition, Fasting Health*. 2024;**12**(1):42-50. <https://doi.org/10.22038/jnfh.2023.74524.1462>.
 16. Nemati H, Sahebighag MH, Mahmoodi M, Ghiasi A, Ebrahimi H, Barzanjeh Atri S, et al. Non-Suicidal Self-Injury and Its Relationship with Family Psychological Function and Perceived Social Support among Iranian High School Students. *J Res Health Sci*. 2020;**20**(1). e00469. [PubMed ID: 32814690]. [PubMed Central ID: PMC7585761]. <https://doi.org/10.34172/jrhs.2020.04>.
 17. Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. Updating guidance for reporting systematic reviews: development of the PRISMA 2020 statement. *J Clin Epidemiol*. 2021;**134**:103-12. [PubMed ID: 33577987]. <https://doi.org/10.1016/j.jclinepi.2021.02.003>.
 18. Klonsky ED, Victor SE, Saffer BY. Nonsuicidal self-injury: what we know, and what we need to know. *Can J Psychiatry*. 2014;**59**(11):565-8. [PubMed ID: 25565471]. [PubMed Central ID: PMC4244874]. <https://doi.org/10.1177/070674371405901101>.
 19. Farkas BF, Takacs ZK, Kollarovics N, Balazs J. The prevalence of self-injury in adolescence: a systematic review and meta-analysis. *Eur Child Adolesc Psychiatry*. 2024;**33**(10):3439-58. [PubMed ID: 37486387]. [PubMed Central ID: PMC11564408]. <https://doi.org/10.1007/s00787-023-02264-y>.
 20. Peterson J, Welch V, Losos M, Tugwell P. The Newcastle-Ottawa scale (NOS) for assessing the quality of nonrandomised studies in meta-analyses. *Ottawa Hosp Res Institute*. 2011;**2**(1):1-12.
 21. Begg CB, Berlin JA. Publication bias and dissemination of clinical research. *J Natl Cancer Inst*. 1989;**81**(2):107-15. [PubMed ID: 2642556]. <https://doi.org/10.1093/jnci/81.2.107>.
 22. Abbasian M, Pourshahbaz A, Taremian F, Poursharifi H. The Role of Psychological Factors in Non-suicidal Self-injury of Female Adolescents. *Iran J Psychiatry Behav Sci*. 2021;**15**(1). <https://doi.org/10.5812/ijpbs.101562>.
 23. Abdollahi E, Kousha M, Bozorgchenani A, Bahmani M, Rafiei E, Eslamdoust-Siahestalkhi F. Prevalence of Self-Harm Behaviors and Deliberate Self-Cutting in High School Students in Northern Iran and Its Relationship with Anxiety, Depression, and Stress. *J Holistic Nurs Midwifery*. 2022;**32**(3):169-77. <https://doi.org/10.32598/jhnm.32.3.2193>.
 24. Babaeifard M, Akbari M, Mohammadkhani S, Hasani J, Shahbazian R, Selby EA. Early maladaptive schemas, distress tolerance and self-injury in Iranian adolescents: serial mediation model of transdiagnostic factors. *BJPsych Open*. 2024;**10**(3). e116. [PubMed ID: 38770605]. [PubMed Central ID: PMC11363084]. <https://doi.org/10.1192/bjo.2024.708>.
 25. Ghanizadeh A, Shekoohi H. Prevalence of nail biting and its association with mental health in a community sample of children. *BMC Res Notes*. 2011;**4**:116. [PubMed ID: 21481256]. [PubMed Central ID: PMC3082216]. <https://doi.org/10.1186/1756-0500-4-116>.
 26. Gholamrezaei M, Heath N, Panaghi L. Non-suicidal self-injury in a sample of university students in Tehran, Iran: prevalence, characteristics and risk factors. *Int J Culture Mental Health*. 2016;**10**(2):136-49. <https://doi.org/10.1080/17542863.2016.1265999>.
 27. Izadi-Mazidi M, Yaghubi H, Mohammadkhani P, Hassanabadi H. Assessing the Functions of Non-Suicidal Self-Injury: Factor Analysis of Functional Assessment of Self-Mutilation among Adolescents. *Iran J Psychiatry*. 2019;**14**(3):184-91. [PubMed ID: 31598120]. [PubMed Central ID: PMC6778602].
 28. Marin S, Hajizadeh M, Sahebighag MH, Nemati H, Ataeiasl M, Anbarlouei M, et al. Epidemiology and Determinants of Self-Injury Among High School Students in Iran: a Longitudinal Study. *Psychiatr Q*. 2020;**91**(4):1407-13. [PubMed ID: 32418140]. <https://doi.org/10.1007/s11266-020-09764-z>.
 29. Mohammadpoorasl A, Fakhari A, Vahidi R, Rostami F, Talebi M. Predicting the incidence of self-injury in Iranian high school students. *Res J Biol Sci*. 2007;**2**(4).
 30. Mohammadpoorasl A, Nedjat S, Fakhari A, Yazdani K, Rahimi Foroushani A, Fotouhi A. Substance abuse in high school students in association with socio-demographic variables in northwest of Iran. *Iran J Public Health*. 1970;**41**(12).
 31. Mozafari N, Bagherian F, Mohammad Zadeh A, Heidari M. Prevalence and functions of self-harming behaviors in adolescents in Sanandaj. *Shenakht J Psychol Psychiatry*. 2021;**8**(4):110-23. <https://doi.org/10.32598/shenakht.8.4.110>.
 32. Nobakht HN, Dale KY. The prevalence of deliberate self-harm and its relationships to trauma and dissociation among Iranian young adults. *J Trauma Dissociation*. 2017;**18**(4):610-23. [PubMed ID: 27736465]. <https://doi.org/10.1080/15299732.2016.1246397>.
 33. Mohammadpoorasl A, Vahidi R, Fakhari A, Rostami F, Dastghiri S. Substance abuse in Iranian high school students. *Addict Behav*. 2007;**32**(3):622-7. [PubMed ID: 16815638]. <https://doi.org/10.1016/j.addbeh.2006.05.008>.
 34. Rezaei O, Athar ME, Ebrahimi A, Jazi EA, Karimi S, Ataie S, et al. Psychometric properties of the Persian version of the inventory of statements about self-injury (ISAS). *Borderline Personal Disord Emot Dysregul*. 2021;**8**(1):27. [PubMed ID: 34772468]. [PubMed Central ID: PMC8588687]. <https://doi.org/10.1186/s40479-021-00168-4>.
 35. Sabet Dizkuhi K, Kafie Masooleh SM. Self-harm Behaviors among University Students. *J Mod Psychol*. 2021;**1**(3):9-24. <https://doi.org/10.22034/jmp.2021.285993.1016>.
 36. Taheri E, Taremian F, Dolatshahe B, Sepehrnia N. [Comparative study of believing in deserving of pain and punishment and access to self-harm-related mental associations in student with no suicidal self-injury and normal students]. *Nurse Physician Within War*. 2021;**9**(32):88-99. FA. <https://doi.org/10.29252/npwjw.9.32.88>.
 37. Bathaee FS, Kamali Fard K. [Investigating the prevalence of physical and mental disorders among first grade male middle school students in Ray and south of Tehran]. *Med Spiritual Cultivation*. 2004;**4**(52):73-68. FA.

38. Nock MK, Favazza AR. Nonsuicidal self-injury: Definition and classification. In: Nock MK, editor. *Understanding nonsuicidal self-injury: Origins, assessment, and treatment*. Washington, DC: American Psychological Association; 2009. p. 9-18. <https://doi.org/10.1037/11875-001>.
39. Selby EA, Bender TW, Gordon KH, Nock MK, Joiner TE. Non-suicidal self-injury (NSSI) disorder: a preliminary study. *Personal Disord*. 2012;**3**(2):167-75. [PubMed ID: 22452757]. <https://doi.org/10.1037/a0024405>.
40. Brunner R, Kaess M, Parzer P, Fischer G, Carli V, Hoven CW, et al. Life-time prevalence and psychosocial correlates of adolescent direct self-injurious behavior: a comparative study of findings in 11 European countries. *J Child Psychol Psychiatry*. 2014;**55**(4):337-48. [PubMed ID: 24215434]. <https://doi.org/10.1111/jcpp.12166>.
41. Swannell SV, Martin GE, Page A, Hasking P, St John NJ. Prevalence of nonsuicidal self-injury in nonclinical samples: systematic review, meta-analysis and meta-regression. *Suicide Life Threat Behav*. 2014;**44**(3):273-303. [PubMed ID: 24422986]. <https://doi.org/10.1111/sltb.12070>.
42. Muehlenkamp JJ, Gutierrez PM. An investigation of differences between self-injurious behavior and suicide attempts in a sample of adolescents. *Suicide Life Threat Behav*. 2004;**34**(1):12-23. [PubMed ID: 15106884]. <https://doi.org/10.1521/suli.34.1.12.27769>.
43. Ghaedi Heidari F, Bahrami M, Kheirabadi G, Maghsoudi J. Factors Associated with Non-Suicidal Self-Injury (NSSI) in Iran: A Narrative Systematic Review. *J Pediatric Perspectives*. 2020;**8**(1):10785-99. <https://doi.org/10.22038/ijp.2019.42967.3599>.
44. De Luca L, Pastore M, Palladino BE, Reime B, Warth P, Menesini E. The development of Non-Suicidal Self-Injury (NSSI) during adolescence: A systematic review and Bayesian meta-analysis. *J Affect Disord*. 2023;**339**:648-59. [PubMed ID: 37479039]. <https://doi.org/10.1016/j.jad.2023.07.091>.
45. Andrei LE, Efrim-Budisteanu M, Mihailescu I, Buica AM, Moise M, Rad F. Non-Suicidal Self-Injury (NSSI) Patterns in Adolescents from a Romanian Child Psychiatry Inpatient Clinic. *Children (Basel)*. 2024;**11**(3). [PubMed ID: 38539332]. [PubMed Central ID: PMC10969364]. <https://doi.org/10.3390/children11030297>.
46. Sahay S, Nilanjana M. Non suicidal self-injury among youths: A probe into the 'what's' and 'whys' of self-harm. *Int J Psychol Res*. 2024;**6**(1):14-6. <https://doi.org/10.33545/26648903.2024.v6.1a.44>.
47. Akbari M, Seydavi M, Firoozabadi MA, Babaeifard M. Distress tolerance and lifetime frequency of non-suicidal self-injury (NSSI): A systematic review and meta-analysis. *Clin Psychol Psychother*. 2024;**31**(1). e2957. [PubMed ID: 38343352]. <https://doi.org/10.1002/cpp.2957>.
48. Ayyubi E, Nazarzadeh M, Bidel Z. [Prevalence of bullying and deliberated self-harm behaviors among high school students]. *J Principles Mental Health*. 2013;**15**(1). FA.
49. Zetterqvist M. *Non-Suicidal Self-Injury in Swedish Adolescents Prevalence, Characteristics, Functions and Associations with Childhood Adversities*. Linköping, Sweden: Linköpings Universitet; 2014.
50. Barrocas AL, Giletta M, Hankin BL, Prinstein MJ, Abela JR. Nonsuicidal self-injury in adolescence: longitudinal course, trajectories, and intrapersonal predictors. *J Abnorm Child Psychol*. 2015;**43**(2):369-80. [PubMed ID: 24965674]. <https://doi.org/10.1007/s10802-014-9895-4>.
51. Muehlenkamp JJ, Williams KL, Gutierrez PM, Claes L. Rates of non-suicidal self-injury in high school students across five years. *Arch Suicide Res*. 2009;**13**(4):317-29. [PubMed ID: 19813109]. <https://doi.org/10.1080/1381110903266368>.
52. Bresin K, Schoenleber M. Gender differences in the prevalence of nonsuicidal self-injury: A meta-analysis. *Clin Psychol Rev*. 2015;**38**:55-64. [PubMed ID: 25795294]. <https://doi.org/10.1016/j.cpr.2015.02.009>.
53. Plener PL, Libal G, Keller F, Fegert JM, Muehlenkamp JJ. An international comparison of adolescent non-suicidal self-injury (NSSI) and suicide attempts: Germany and the USA. *Psychol Med*. 2009;**39**(9):1549-58. [PubMed ID: 19171079]. <https://doi.org/10.1017/S0033291708005114>.