

Fragile X Mental Retardation Protein is Reduced in Cerebella of Subjects with Autism

Seyyed Hossein Fatemi, MD, PhD^{**}, Timothy D Folsom, BS, MS^{*}

Autism and fragile X syndrome have multiple behavioral phenotypes in common, however there has been little research into the potential role of fragile X mental retardation protein (FMRP) in autism. We sought to measure protein levels of FMRP in the lateral cerebella of subjects with autism and matched controls. We found significantly reduced levels of FMRP ($p < 0.0063$) in subjects with autism. The loss of FMRP expression in subjects with autism may help explain behavioral deficits, presence of seizure, and reduction in GABA receptors in subjects with autism.

Declaration of interest: None

Iranian Journal of Psychiatry and Behavioral Sciences (IJPBS), Volume 4, Number 2, Autumn and Winter 2010: 56-57.

Introduction

Fragile X mental retardation protein (FMRP) regulates post-transcriptional events including splicing, nuclear export, and translation of multiple RNAs (1). Anywhere from 25-47% of patients with Fragile X syndrome (FXS), in whom FMRP expression is lost due to transcriptional silencing, display autistic behavior (2,3) and it has been hypothesized that these behavioral deficits may be the result of this decreased FMRP expression (3). A number of structural abnormalities of the cerebellum have been identified in subjects with autism, which may be responsible for motor system dysfunction observed in this disorder (4). We therefore decided to examine protein levels of FMRP in cerebella of adult subjects with autism vs. matched controls. All experimental procedures were approved by the Institutional Review Board of the University of Minnesota, School of Medicine.

Materials and Methods

Postmortem blocks of lateral cerebellum (N=6 control; N=4 autism) were obtained

from the Autism Research Foundation and various brain banks. The tissue samples were prepared (5,6) and were subjected to SDS-PAGE and western blotting as described previously (5,6). The blots were then incubated with anti-FMRP (Abcam) or anti- β -actin antibodies (Sigma Aldrich) followed by an appropriate secondary antibody. We measured FMRP protein using western blotting technique and normalized results against β -actin.

Results

There was a significant 80% reduction in FMRP protein in cerebella of subjects with autism ($p < 0.0063$) when compared with controls (mean value of 0.323 ± 0.132 for controls vs. 0.0658 ± 0.0506 for subjects with autism) (Figure 1). There was no statistically significant difference in age and postmortem interval (PMI) between the two groups (data not shown).

Discussion

We observed a reduction in FMRP in cerebella of subjects with autism, none of whom were comorbid for FXS. The reduction in FMRP, and consequent dysregulation of downstream genes, may help to explain defects that are common between autism and FXS: social anxiety, mental retardation, seizures, and reduced size of cerebellum (3,7). Reduction in FMRP may also impact the

Authors' affiliations : * Department of Psychiatry, Division of Neuroscience Research, University of Minnesota Medical School
** Departments of Pharmacology and Neuroscience, University of Minnesota Medical School

•**Corresponding author :** Department of Psychiatry, Division of Neuroscience Research, University of Minnesota Medical School, 420 Delaware St SE, MMC 392, Minneapolis, MN 55455 USA.
Tel : 612 626 3633
Fax : 612 624 8935
E-mail: fatem002@umn.edu

GABAergic signaling system as a murine model of FXS also displays reduced expression of a number of GABA_A receptor subunits (GABR α 1, GABR α 3, GABR α 4, GABR β 1, GABR β 2, GABR γ 1, GABR γ 2, and GABR δ) (8). Moreover, our laboratory has demonstrated significant reductions in a number of GABA_A (GABR α 1 and GABR β 2) and GABA_B (GABBR1 and GABBR2) receptor subunits in cerebella from the same set of tissues (5,6). Further studies, examining additional brain areas including superior frontal cortex, and hippocampus need to be undertaken to determine whether reduction of FMRP in brains from subjects with autism is a global phenomenon.




Figure 1. A. Representative samples of FMRP and β -actin from cerebella from control (C) and autistic (A) subjects. B. Mean FMRP/ β -actin ratios for autistic (filled histogram bars) and control subjects are shown for cerebellum. (Error bars expressed as standard deviation of the mean.) *, $p < 0.05$.

Acknowledgement

Human tissue was obtained from the NICHD Brain and Tissue Bank for Developmental Disorders at the University of Maryland, Baltimore, MD; the Harvard Brain Tissue Resource Center, which is supported in part by PHS grant number R24 MH068855; the Brain Endowment Bank, which is funded in part by the National Parkinson Foundation,

Inc., Miami, Florida; and the Autism Tissue Program is greatly appreciated. Grant support by National Institute of Child Health and Human Development (#5R01HD052074-01A2 and 3R01HD052074-03S1) to SHF is gratefully acknowledged.

References

1. De Rubeis S, Bagni C. Fragile X mental retardation protein control of neuronal mRNA metabolism: Insights into mRNA stability. *Mol Cell Neurosci* 2010; 43(1): 43-50.
2. Cohen I. Behavioral profiles of autistic and nonautistic fragile X Males. *Dev Brain Dyn* 1995; 8: 252-269.
3. Hatton DD, Sideris J, Skinner M, Mankowski J, Bailey DB Jr, Roberts J, et al. Autistic behavior in children with fragile X syndrome: prevalence, stability, and the impact of FMRP. *Am J Med Genet A*. 2006; 140A(17): 1804-13.
4. Nayate A, Bradshaw JL, Rinehart NJ. Autism and Asperger's disorder: are they movement disorders involving the cerebellum and/or basal ganglia? *Brain Res Bull* 2005; 67(4): 327-34.
5. Fatemi SH, Reutiman TJ, Folsom TD, Thuras PD. GABA(A) receptor down regulation in brains of subjects with autism. *J Autism Dev Disord* 2009; 39(2): 223-30.
6. Fatemi SH, Folsom TD, Reutiman TJ, Thuras PD. Expression of GABA (B) receptors is altered in brains of subjects with autism. *Cerebellum* 2009; 8(1): 64-9.
7. Gothelf D, Furfaro JA, Hoeft F, Eckert MA, Hall SS, O'Hara R, et al. Neuroanatomy of fragile X syndrome is associated with aberrant behavior and the fragile X mental retardation protein (FMRP). *Ann Neurol* 2008; 63(1):40-51.
8. D'Hulst C, De Geest N, Reeve SP, Van Dam D, De Deyn PP, Hassan BA, et al. Decreased expression of the GABAA receptor in fragile X syndrome. *Brain Res* 2006; 1121(1): 238-45.