

Factors Associated With Suicidal Attempts in Iran: A Systematic Review

Mitra Hakim Shooshtari,^{1,*} Seyyed Kazem Malakouti,¹ Leili Panaghi,² Shohreh Mohseni,¹ Naghmeh Mansouri,¹ and Afarin Rahimi Movaghar³

¹Department of Psychiatry, Mental Health Research Center, Tehran Institute of Psychiatry, Faculty of Behavioral Sciences and Mental Health, Iran University of Medical Sciences, Tehran, IR Iran

²Department of Psychology, Family Research Institute, Shahid Beheshti University, Tehran, IR Iran

³Iranian Research Center for HIV/AIDS, Tehran University of Medical Sciences, Tehran, IR Iran

*Corresponding author: Mitra Hakim Shooshtari, Department of Psychiatry, Mental Health Research Center, Tehran Institute of Psychiatry, Faculty of Behavioral Sciences and Mental Health, Iran University of Medical Sciences, Tehran, IR Iran. Tel/Fax: +98-2166506899, E-mail: mitra.hakim@gmail.com

Received 2014 July 27; Revised 2015 February 28; Accepted 2015 October 10.

Abstract

Context: Suicide prevention is a health service priority. Some surveys have assessed suicidal behaviors and potential risk factors.

Objectives: The current paper aimed to gather information about etiology of suicide attempts in Iran.

Data Sources: Pubmed, ISI web of science, PsychInfo, IranPsych, IranMedex, IranDoc as well as gray literature were searched.

Study Selection: By electronic and gray literature search, 128 articles were enrolled in this paper. Pubmed, ISI web of science, PsychInfo, IranPsych, IranMedex, IranDoc were searched for electronic search. After reading the abstracts, 84 studies were excluded and full texts of 44 articles were reviewed critically.

Data Extraction: Pubmed, ISI web of science, PsychInfo, IranPsych, IranMedex, IranDoc as well as gray literature were searched to find any study about etiologic factors of suicide attempt in Iran.

Results: Depressive disorder was the most common diagnosis in suicide attempters that is 45% of the evaluated cases had depression. One study that had used Minnesota multiphasic personality inventory (MMPI) found that Histrionics in females and Schizophrenia and Paranoia in males were significantly influential. Family conflicts with 50.7% and conflict with parents with 44% were two effective psychosocial factors in suicidal attempts. In around one fourth (28.7%) of the cases, conflict with spouse was the main etiologic factor.

Conclusions: According to the methodological limitations, outcomes should be generalized cautiously. Further studies will help to plan preventive strategies for suicidal attempts; therefore, continued researches should be conducted to fill the data gaps.

Keywords: Etiology, Iran Suicidal Attempt, Suicidal Behavior, Suicidal Idea, Suicide

1. Context

Suicide is one of the mental health priorities in world health organization (WHO). According to the WHO report (2003), about 1,000,000 people die because of suicide each year. Findings of suicide attempts in general population demonstrate that the rate may vary from 0.2 to 2.6 and increase up to 10% during lifetime (1, 2). The one month-prevalence of 25% was reported in a study carried out in Morocco (3). A large population study systematically assessed suicide incidence with various psychiatric disorders. In Denmark 21,169 suicides over a 17-year period were compared with gender-age-time-matched people controls (4). According to this study, suicide risk significantly increased for persons with a hospitalized psychiatric disorder. This risk varies significantly by diagnosis, gender and age of subjects. The influence of various disorders usually decreases with increasing age; however, there are some exceptions. Schizophrenia affects people aged ≤ 35 years in the strongest form. Suicide risk strongly increases by recurrent depression in all ages. In young people, borderline personality disorder has a strong effect. For the persons over 60 years old, reaction to stress

and adjustment disorder increases the risk for suicide.

A meta-analysis about the prevalence of depression in Iranian adolescents showed depression rates of 43.5%, 15.8% and 13%, employing BDI, symptom checklist-90 (SCL-90) and CDI, respectively (5). The WHO world mental health (WMH) surveys assessed suicidal behaviors and potential risk factors among 108,705 adults from 21 countries. In this research the WHO composite international diagnostic interview (CIDI) was applied. Risk factors for suicidal behaviors in both developed and developing countries were as follows:

Younger age, female, lower income and education, being single, unemployment, parents with psychiatric disorder, childhood adversities, and presence of 12-month DSM-IV psychiatric disorders (6).

A research was conducted in Korea to assess socio-demographic status in persons with suicidal ideation. Suicidal ideation was most prevalent among middle-aged females who were blue-collar workers (7). The middle-aged men with absent spouse showed significantly higher suicidal ideation.

Since various studies were carried out on suicide in different population in Iran, and each of them evaluated just some effective factors in attempting suicide (8-10), implementing a systematic review study to draw a vast picture of suicide and identifying informational gaps would be helpful for future studies.

2. Objectives

Attempted suicide behavior is a multi-factorial phenomenon from etiologic perspective. The current systematic review aimed to assess various effective factors in suicide attempts. These factors were classified and evaluated in three subgroups: psychiatric illnesses, personality problems and psycho-social variables. Also, demographic information of suicide attempter, the history of their mental-physical illnesses and level of perceived psychiatric services were appraised in this systematic review.

Cross-sectional and case-control studies which had evaluated etiology of suicide attempts were included in this review. The investigation covered all of the suicide attempters aged more than 15 years referred to the emergency wards.

3. Data Sources

3.1. Electronic Search (Desk Research)

Studies that focused on etiologic factors, published in peer-reviewed journals, and examining both social factors and suicide or related phenomenon (death wishes, suicide ideation and deliberate self-harm/non-fatal suicidal behavior) were considered eligible for inclusion.

Studies were identified through electronic searches performed through the Pubmed, PsychINFO, IranPsych, SID, IranMedex, IranDoc, Cinahil, WHO Int and ISI databases.

No search terms relating to social factors were applied. All publication years were considered. The following keywords were used in the search:

Suicide, suicidal idea, suicidal attempt, suicidal behavior and etiology, Iran.

3.2. Gray literature

About 5,000 theses from all universities of medical sciences all over the country (40 universities) and psychology colleges under coverage of ministry of science, research and technology were searched. Most of them were accessible from the library of universities. Also, all final reports of researches documented in Iran Psych Database were searched. None of them was about etiological factors in suicidal attempts.

4. Study Selection

4.1. Review Method

4.1.1. Study Identification and Selection

At first, the abstracts of the attained articles from elec-

tronic search and gray materials were separately reviewed by two investigators for the relevancy with the subject. In cases with no agreement, final decisions were made after discussion. Full texts of the researches selected based on their relevance to the research criteria were collected. Then, they were assessed and related forms were completed. Totally, 128 articles were included for abstract review.

4.2. Quality Assessment

The full texts of selected articles were appraised by two independent reviewers using critical appraisal form.

5. Data Extraction

5.1. Data Collection and Integration

In most parts of the literatures, there is only descriptive information about suicidal attempts and relationship between etiologic factors and demographic variables are not analyzed.

Since in each study, different etiologic factors were investigated, time trend estimation in etiology of suicidal attempts was not possible. Eighty-four papers were excluded from the study due to non-relevant data. A flow-chart was drawn to show the search and selection process identification (Figure 1).




Figure 1. Flowchart of the Search and Selection Process Identification

The full texts of included articles (44 literatures) were reviewed and the designed tables for data extraction were completed for etiologic factors. Because of the highly heterogeneous factors and variety of data collection scales, the results were not integrated and narrative report is provided. To compare different studies more easily, also to get a better perception of etiologic factors; the collected data were categorized into three subgroups of psychiatric illnesses, personality problems and psycho-social variables. It is obvious that interpretation and generalization of these divergent results should be made cautiously.

5.2. Literature Description

Totally, 128 articles were enrolled in the study. After reading the abstracts, 84 studies were excluded and full texts of 44 articles were reviewed critically and 23 articles addressing different etiologic factors were considered. Among them, 21 articles were excluded because of their irrelevancy to psychological and psycho-social factors. Four of them had evaluated association between suicide commitment and lunar months. One article had assessed mental disorders in patients who had committed suicide with fatal and serious methods.

One research had evaluated suicide attempts in patients with addiction; another one was specified just to children aged 6 - 13 years old. In 14 studies, young people had been evaluated and considering different assessed factors in each study, they were not compared. From 44 related studies, 23 articles met the criteria of systematic review; eight studies were about psychiatric illnesses and six of them about personality problems. In nine studies psycho-social variables were attained.

Since two studies had assessed both mental disorders and psycho-social variables and one article had assessed both personality problems and psycho-social variables, they are mentioned in the tables twice.

Three studies had been done using hospital archive files and the others performed by cross-sectional design about people referred to emergency departments. Of the total 6062 persons attempting suicide, 5389 were from emergency departments and 673 cases were related to hospital files.

To evaluate psychiatric disorders, diagnostic and statistical manual of mental disorders, fourth edition (DSM-IV) criteria had been used in three studies and one study was based on DSM-III criteria. Two studies had applied SCL-90 and mood disorders questionnaire (MDQ) to measure psychiatric symptoms.

Assessment of personality problems in suicide attempters had been done by MMPI, the Millon clinical multi-axial inventory-II (MCMI-II) and the Eysenck personality inventory.

The studies on psycho-social factors had used new questionnaires and none of them referred to validity and reliability of the scales. None of these studies had used valid questionnaires. They did not use any open ended questions.

6. Results

Among the 44 full text papers, 23 articles met the criteria of systematic review; eight studies were on psychiatric illnesses and six of them about personality problems. Nine studies had attained psycho-social variables. Of the total 5,220 persons attempting suicide in these papers 5,247 were from emergency departments and 673 cases from hospital files.

Young adults of 20 - 24 years had the highest suicide attempt rates. In one study the rate was 64.5% of the suicide attempters. Females attempted suicide two times more than the males. One study showed that significant interpersonal problems had an important role in suicide attempts among 15 to 25-year-old single persons, while significant family problems were important in married subjects less than 20 years. Educated persons with college certification had less suicide attempts. In order to get the conclusion easier, effective factors in suicidal attempts were ranked based on median and based on that three charts were drawn.

6.1. Psychiatric Diagnosis

Depressive disorder was the most common diagnosis in suicide attempters; 45% of the evaluated cases had depression. The rate of bipolar and anxiety disorders were 20% and 17.8% respectively. According to Figure 2, personality disorder with 5.8% had the lowest role among the assessed psychiatric disorders in attempting suicide.




Figure 2. Psychiatric Disorder in Suicide Attempters

6.2. Personality Problems

A study that used MMPI found that histrionics in females and schizophrenia and paranoia in males were significantly more than other problems. Depression and hypochondria in male and female suicide attempters were significantly more than other problems. Based on the study that applied the Eysenck personality inventory, tendency to commit suicide was associated with neuroticism and psychoticism. This result was confirmed in other studies. Table 1 shows the findings of all studies about personality problems.

Table 1. Personality Problems in Suicide Attempters

References	Year	Sample Size	Study Type	Scale	Personality Problems
(11)	2003	30 college students	Case-control	MCMI-II	Anxiety, somatization, mania, depression dysthymia, alcohol dependence, drug dependence , schizophrenia, paranoid, avoidant personality, antisocial, borderline, schizotypal in suicide attempters is more than others.
(12)	1995	667 Subjects, Hamadan		MMPI	Personality disorders: 4.19% neurotic disorder: 3.16% psychosis: 5.13% health : 6.47% NOS: 2.3% severe depression: 4.34%
(13)	1995 - 96	40 Subjects Semnan	Retrospective	MMPI	Histrionic in females and schizophrenia & paranoia in males were significantly more than others. Depression and hypochondria in male and females suicide attempters was significantly more than others.
(14)	1978	440 college Iranian and Turkish student		Eyzenck	Tendency to suicide was associated with neuroticism and psychoticism
(15)	1997	56 Subjects, Ilam		Eyzenck	Extraversion in both genders was lower and psychoticism and neuroticism were significantly higher (except psychoticism in females) lying in males was more than control group.

Abbreviations: NOS, not otherwise; OCD, obsessive compulsive disorder.




Figure 3. Psychosocial Problems in Suicide Attempters




Figure 4. History of Mental Disorders in Suicide Attempters

6.3. Specified Psychosocial Problems

Family conflicts with 50.7% and conflict with parents with 44.1% were two effective psychosocial factors in suicidal attempts. In around one fourth (28.7%) of the cases, conflict with spouse was the main etiologic factor (Figure 3). Psychosocial factors involved in suicide attempts are

summarized in Tables 2 - 4.

Some information such as personal and family history of suicide attempts, history of mental and physical illnesses, and the level of received psychiatric services could be observed in Figure 4 and Table 5.

Table 2. [Part I] Psychosocial Factors Involved in Suicide Attempt

References	Sample Size	Year	Study Type	Demographic Factors	Scale	Family Problems	Conflict With Spouse	Conflict With Parents	Addiction	Romance Problems	Financial Problems	Death of Loving persons	Illness	Educational Problems	Legal Problems	Other Factors
(16)	600 subjects, house wives and employed women, (toxic city, Isfahan)	2005	Descriptive, convenient			House wives, 71% employed, 63%; no significant difference			House wives 4.8%; employed, 6.4%; No significant difference			%4	%4.3			
(17)	173 subjects referred to Rasht EMS	2001	Descriptive, convenient	In age between 18 to 24, 53.7% illiterate, 53.2% elementary school, 4.6% high school, 3.4%; high education 9.21%; Female, 67.8%; male, 32.2%; Single, 53.2%; married, 42.2%; significant interpersonal problems between 15 to 25 years and singles, significant family problems in married subjects	Researcher made	%76.1				%25	%1.19	%4	%4.3	%3.2	%1.1	
(18)	390 subjects referred to Bandar Abbas EMS	2001	Descriptive, convenient	Under 20 year: 6.33%; 20 to 30 years, 3.51%; married and house wives, maximum 39.7%		%83			20% history of substance abuse							
(19)	539 subjects referred to Zanjan EMS	1996	Descriptive, convenient	20% unsuccessful marriage, financial problems in males; unsuccessful education in females			40%, 20% unsuccessful marriage	%30								
(20)	100 subjects referred to EMS in Mashhad		Descriptive, convenient	Illiterate, 9%; elementary, 30%; guidance school, 30%; high school, 29%; college 2%; single, 43%; married, 42%		%59	%29			49% medium, 4% poor, 11% good						
(21)	62 subjects aged 11-25 years referred to Shiraz EMS	1996	Descriptive, convenient	%64.5 Female attempter		%32								%31		Not being successful, 24%; have no marriage permission, 11%
(19)	519 subjects referred to Zanjan EMS	96-1995	Descriptive, convenient	illiterate, 18.5%; elementary school, 26%; high school, 22.3%; college 4.3%; married, 53.7%		%50.7	%55			%4.8	%9 (Vocational) %4		%3.4	%4.6		

Abbreviation: EMS, emergency medical services.

Table 3. [Part 2] Psychosocial Factors Involved in Suicide Attempt

References	Sample Size	Year	Study Type	Demographic Factors	Scale	Fam-ily Prob-lems	Conflict With Spouse	Conflict With Parents	Addiction	Romance Problems	Financial Prob-lems	Death of Loving persons	Illness	Educa-tion Prob-lems	Legal Prob-lems	Other Factors
(22)	58 subjects from Chaharmahal and Bakhtiari province	2000	Descriptive files	20 to 24 years, 29.4%; 25 to 29 years, 27.6%; illiterate, 22.4%; elementary, 41.2%; guidance school, 20.7%; High school, 12.2%, married, 53.4%	Researcher made questionnaire	41% physical punishment via spouse, 28%	32.1%			35.7% expensive-ness, 25% (about 50%), 37% salary problems	26.7% being inpatient 25%	37.5%	32.1%			Bombing 42.8%, house-keeping respon-sibilities, 28.6% (42.8%), lovely person separation, 25% (31.3%)
(23)	52 subjects, 13 to 19 years referred to AlvazEMS	2000 - 2001	File	Female 55.8% under 18 years interpersonal, 58%; familial 75%; 18 - 24 years financial problems, 20.4%; interpersonal, 23.4%, familial, 76.6%; vocational, 14%; 25 - 34 years, financial, 23.5%; vocational and interpersonal, 17.6% familial, 82.4%; over 35 years, 20.8% death of the beloved persons, familial, 66.7		59.7%	28.4%	58.2%		43% (18%) friends, 82% romance	(21%) 33% vocational					
(24)	173 subjects	2001	File													
(25)	500 subjects referred to Emam Reza hospital Mashhad	2003 - 2004	File	Under 20 years, 48%; 21 - 40 years, 43.4%; female 67.8%, married, 50.4%												Good, 52.2%, intermediate, 11.6%, poor 36.2% unemployed, 73%, employed, 27%

Abbreviation: EMS, emergency medical services.

Table 4. [Part 3] Psychosocial Factors Involved in Suicide Attempt

References	Sample Size	Year	Study Type	Demographic Factors	Scale	Family Problems	Conflict With Spouse	Conflict With Parents	Addiction Problems	Romance Problems	Financial Problems	Death of Loving persons	Education Problems	Legal Problems	Other Factors
(26)	15 subjects, adolescents		Analytic	Suicide ideation questionnaire, hopelessness scale Self-concept test, Children attribution questionnaire											Problem solving skill was effective to decrease suicide ideation thoughts (significant), level of hopelessness was significantly decreased in case group, there was no significant difference in self concept
(27)	26 subjects from Ilam	1997	Analytic	10 female, 16 male with mean age of 16.8 years											Problem solved through training significantly decreased depression severity level and hopelessness, and also improved coping skills
(28)	40 subjects referred to Loghman hospital in Tehran	1999	Analytic												Memory recall speed in depressed suicide attempters was less lower than normal people, retrieving of depression experiences were faster than positive memories, recalling memory experience was highly holistic
(29)	300 subjects from Tabriz	2002	%10		%50	%22.7					Unemployment, 15.7%; financial bankruptcy, 2.7%; poverty, 1.7%	%1.3	%2.7		Loss of social base, 4.7%; spouse illicit relations, 0.7%; aging, 0.7%; lack of aim of life, 3.3%
(30)	946 subjects from Karaj	2003			%27.2	%34.9					%10.3	%5.2	%10.7		
(12)	667 subjects from Hamadan	1995			%30.9	17.4% forced marriage, 3.6%		%1.2	%3.1	%7.6		%1.8	%8.4		

Table 5. History of Physical and Mental Illnesses in Suicide Attempters and the History of Suicide in Attempters and Their Families

References	Year	Sample Size	Visit to a Psychiatrist	History of Tranquilizer Consumption	History of Mental Outpatient	History of Mental Illness	History of Suicide	History of Suicide in Family Members	Having Physical Illness
(31)	2002	50 subjects, two hospitals of Birjand, convenient method	%36		%6	%80			
(32)	1998	673 subjects, teaching hospitals of Urmia					%7.3		
(33)	2000	100 subjects, Shiraz convenient		%38		%59			
(29)	2002	300 subjects, Tabriz convenient					%17.7	%3.7	%6
(30)	2003	946 subjects, Karaj catchment area	58.5% (mental health services)			Neurotic disorders one year before commitment	%28	%15.4	
(34)	1992	subjects, 225 Qazvin						%24	
(16)	2005	600 housewives and employed (toxic urgency, Isfahan) descriptive				Housewives: 38% and Employed 32% (No significant difference)			
(18)	2001	390 subjects, Bandar Abbas EMS					%11	%24	

Abbreviation: EMS, emergency medical services.

7. Conclusions

The current systematic review evaluated the etiologic factors of suicide attempts. The evaluated studies in the research not only had used different scales, but also each of them had assessed a different group of effective factors and just three articles had assessed a variety of etiologic factors. Thus, recognizing the role of each factor in suicide attempts was complicated. It appears that if similar scales were used, the heterogeneity of etiologic factors would be declined. In short, this systematic review showed that young adulthood, female gender and lower education are associated with increased risk of suicidal attempts in the Iranian population. It was concordant with a research in 17 countries on 84,850 subjects, which evaluated effective factors on suicide attempts. Female gender, being single, lower age and education are important factors in suicidal attempts (35). According to the current systematic review, major depression, bipolar

disorders and anxiety disorders were the most frequent psychiatric disorders associated with suicidal attempts in the Iranian population. The Nock research (35) showed that mood disorders and low impulse control were the most important diagnostic risk factors in high and low income countries respectively. If Iran is accounted a low-income developing country, impulse control disorder may play a role in suicide attempts. On the other hand, the results of the current systematic review showed that conflict in interpersonal relationship significantly influenced half of the suicidal attempts. These conflicts may arise from poor impulse control. Next precise researches can clarify such a pathway. These answers can facilitate designing interventions for prevention strategies to decrease suicide in the country.

Although there were numerous limitations in concluding these studies, yet, the most prevalent effective factors

in each subgroup could be described. The lack of advanced statistical models such as logistic regression and odds ratio complicated the identification of main effective factors in suicide attempts.

Another limitation was lack of a control group in most of these samples, which led to difficult generalization of the results. Therefore, considering the methodological limitations, outcomes should be generalized cautiously. Specific suicidal attempts, such as self-poisoning, self-cutting and self-immolation or self-burning are not selected for search. It is suggested to use valid and common scales in future studies.

In the end, it is reminded that there is a lack of systematic assessment of the reliability of suicide statistics about the epidemiology throughout the world (36).

According to the methodological limitations, the outcomes should be generalized cautiously. Such studies will help to plan preventive strategies for suicidal attempts; therefore, continued researches should be conducted to fill data gaps.

Acknowledgments

Authors wish to thank Seyed Reza Majdzadeh, Arash Rashidian, Arash Etemadi, Hojat Salmasian, Saharnaz Nejat, Soroush Mortaz Hejri, Vandad Sharifi and Seyyed Vahid Shariat, who assisted the research team with their consultation. Authors also thank Ms. Zahra Bayat for her assistance to access the full texts of the studies and data transfer.

Footnotes

Authors' Contribution: Study design and manuscript drafting: Mitra Hakim Shooshtari; study design, statistical analysis and manuscript drafting: Seyyed Kazem Malakouti; statistical analysis and manuscript revising: Leili Panaghi; clinical data collection, interpretation and revising the manuscript: Shohreh Mohseni; re-analyzing the clinical and statistical data: Naghmeh Mansouri; writing first draft of manuscript: Afarin Rahimi Movaghar. All authors read and approved the final manuscript.

Funding/Support: This article is a product of contraction number 132/8545 of Tehran University of Medical Sciences, contract number 06/775 world health organization, and contract number 26,212 Tehran psychiatric institute.

Declaration of Interest: None.

References

1. Renberg ES, Hjelmeland H, Koposov R. Building models for the relationship between attitudes toward suicide and suicidal behavior: based on data from general population surveys in Sweden, Norway, and Russia. *Suicide Life Threat Behav.* 2008;**38**(6):661-75. doi: 10.1521/suli.2008.38.6.661. [PubMed: 19152297]
2. Shooshtary MH, Malakouti SK, Bolhari J, Nojomi M, Posht-mashhadi M, Amin SA, et al. Community study of suicidal behaviors and risk factors among Iranian adults. *Arch Suicide Res.* 2008;**12**(2):141-7. doi: 10.1080/1381110701857475. [PubMed:

- 18340596]
3. Agoub M, Moussaoui D, Kadri N. Assessment of suicidality in a Moroccan metropolitan area. *J Affect Disord.* 2006;**90**(2-3):223-6. doi: 10.1016/j.jad.2005.09.014. [PubMed: 16352345]
4. Qin P. The impact of psychiatric illness on suicide: differences by diagnosis of disorders and by sex and age of subjects. *J Psychiatr Res.* 2011;**45**(11):1445-52. doi: 10.1016/j.jpsychires.2011.06.002. [PubMed: 21722920]
5. Sajjadi H, Mohaqeqi Kamal SH, Rafiey H, Vameghi M, Forouzan AS, Rezaei M. A systematic review of the prevalence and risk factors of depression among Iranian adolescents. *Glob J Health Sci.* 2013;**5**(3):16-27. doi: 10.5539/gjhs.v5n3p16. [PubMed: 23618471]
6. Borges G, Nock MK, Haro Abad JM, Hwang I, Sampson NA, Alonso J, et al. Twelve-month prevalence of and risk factors for suicide attempts in the World Health Organization World Mental Health Surveys. *J Clin Psychiatry.* 2010;**71**(12):1617-28. doi: 10.4088/JCP.08m04967blu. [PubMed: 20816034]
7. Moon SS, Park SM. Risk factors for suicidal ideation in Korean middle-aged adults: the role of socio-demographic status. *Int J Soc Psychiatry.* 2012;**58**(6):657-63. doi: 10.1177/0020764011433626. [PubMed: 22231659]
8. Ghafarian Shirazi HR, Hosseini M, Zoladl M, Malekzadeh M, Momennejad M, Noorian K, et al. Suicide in the Islamic Republic of Iran: an integrated analysis from 1981 to 2007/Suicides en République islamique d'Iran: une analyse intégrée de 1981 à 2007. *East Mediterr Health J.* 2012;**18**(6):607-13. [PubMed: 22888617]
9. Heidari Pahlavian A. The study of psycho-social factors and epidemiological characteristics of the people who attempted suicide in Hamadan. *Iran J Psychiatry Clin Psychol.* 1997;**31**(3):1-2.
10. Asghari F, Sadeghi A, Aslani K, Saadat S, Khodayari H. The Survey of Relationship between Perceived Stress Coping Strategies and Suicide Ideation among Students at University of Guilan, Iran. *Int J Sci Technol Educ Res.* 2013;**1**(11):11-8.
11. Hoseiniani A, Moradi AR, Yaryary F, editors. Personality disorders and suicide attempts in the Iranian students; *Proceedings of the 3rd Congress of Students Mental Health*; 2006; University of Science and Technology, Counseling office.
12. Heidari Pahlavian A. The study of psychosocial factors and epidemiological characteristics of the people who attempted suicide in Hamadan. *Quarterly J Andeeshah Raftar.* 1997;**10**(9):11-31.
13. Seyfi A, Rafiee M. Epidemiology of suicide attempters to the hospitals of Semnan University of Medical Sciences from 2006 to 2008.
14. Irfani S. Personality correlates of suicidal tendency among Iranian and Turkish students. *J Psychol.* 1978;**99**(2d Half):151-3. [PubMed: 671376]
15. Mohammadian F. *Study relationship between personal characteristics, stress and fighting ways to suicide in Ilam [Dissertation in Persian]*. Iran, psychiatry institution: Tehran University of Medical Sciences; 2000.
16. Naraghi A, Eizadi N, Akouchakian SH, Masoomi GH, Naderolasli M, Ahmadloo H. Comparison of some risk factors for suicide attempt between housewife's and employed women. *Sci J Forensic Med.* 2015;**20**(2):47-53.
17. Khalkhali SMR, Rahbar M, Farde MR, Jamadi A. Survey of life events prior to suicide attempt[in persian]. *J Guilan Uni Med Sci.* 2001;**10**(39-40):96-106.
18. Yousefi H, Sobhani G, Asadi Nogave F. Determining probable factors of attempted suicide in patients referred to emergency section of the Shahid Mohammadi hospital. *Hormozgan Med J.* 2002;**2**(2)
19. Ghasemi MH, Taremian F, Hashemi M, Hashemipour N. *Prevalence of suicide in Zanjan*. Zanjan: Zanjan University of Medical Sciences; 1998.
20. Hossini SH, Toroski M, Asadi R, Rajabzadeh R, Alavinia SM, Khakshor Trend of attempted suicide and its related factors in Bojnurd city 2006 to 2011. *J North Khorasan Uni Med Sci.* 2012;**4**(4):552.
21. Bahadori Khalili R, Ebrahimi A, Jahanshahi K, Ramezani AA, editors. Study of reasons of suicide in women referred to emergency room in Shiraz [in Persian]; *Second Congress of High Risk Behaviors*; 2007.
22. Zahedi MR, Borjian MT, Shakeri M. *Epidemiology and etiology of*

- suicide in 2001[Unpublished Work]. Chahar Mahal Va Bakhtiari: 2001.
23. Hosseinpour M, Ghafari SM, Mehrbizadeh M. A study on the incentives of suicide attempts among adolescents referred to Golestan Hospital of Ahwaz in 2001-2002[in Persian]. *Sci Med J Ahvaz Uni Med Sci.* 2004;**4**(1):24-30.
 24. Fardmasoud R, Jamadi A, Rahbar M, Khalkhali MR. Life events in suicide attempters. *J Gilan Uni Med Sci.* 2002;**10**(3):96-106.
 25. Attaran H, Shariat M. Epidemiologic study of death due to acute toxicity in 1996, mashhad. *Forensic Med J.* 1996;**3**(10):34-50.
 26. Shafiabadi MH. The effect of group consultation in decreasing tendency to suicide attempt, . *Institute for Cognitive Sciences Studies.* 2002.
 27. Bapiri O. Study of group training of problem solving with psychological characteristics of adolescents with suicide attempt. *Sci j Ilam Uni Med Sci.* 2011;**18**(1)
 28. Kaviani H, Rahimi P, Naghavi HR. Iranian depressed patients attempting suicide showed impaired memory and problem-solving. *Arch Iran Med.* 2004;**7**(2):113-7.
 29. Anvarabnavi M. *The study of risk factors in suicide attempts by drug poisoning in 2003 [Unpublished Work].* 2003.
 30. Nojomi M, Malakouti SK, Bolhari J, Hakimshooshtari M, Fleischmann A, Bertolote JM. Epidemiology of suicide attempters resorting to emergency departments in Karaj, Iran, 2003. *Eur J Emerg Med.* 2008;**15**(4):221-3. doi: 10.1097/MEJ.0b013e3282f4d12a. [PubMed:19078819]
 31. Mehran N, Bulhara J, Asgarnejad FA, Mire M. The socio-psychological characteristics of suicide attempters in contrast with normal people in Birjand. *J Birjand Uni Med Sci.* 2004;**11**(2):3-13.
 32. Marhamati R. *The study of demographic and psychosocial variables in suicide attempters of hospitals of Uromia University of Medical Sciences [Unpublished Work].*
 33. Ashkani H, Dehbozorgi GR, Emamgholi PN. Assessment of the epidemiological factors associated with suicide among the patients admitted in the psychiatric emergency ward[in Persian]. *Med J Tabriz Uni Med Sci Health Serv.* 2003;**(56)**:16-21.
 34. Shaikhholeslami H, Flahzadeh M. Suicides referrals to the emergency clinic of Qazvin[in Persian]. *J Qazvin Uni Med Sci Health Serv.* 1997;**3**:24-30.
 35. Nock MK, Borges G, Bromet EJ, Alonso J, Angermeyer M, Beautrais A, et al. Cross-national prevalence and risk factors for suicidal ideation, plans and attempts. *Br J Psychiatry.* 2008;**192**(2):98-105. doi: 10.1192/bjp.bp.107.040113. [PubMed: 18245022]
 36. Tollefsen IM, Hem E, Ekeberg O. The reliability of suicide statistics: a systematic review. *BMC Psychiatry.* 2012;**12**:9. doi: 10.1186/1471-244X-12-9. [PubMed: 22333684]