

Protective Effect of Crocin on Malathion-induced Cardiotoxicity in Rats: A Biochemical, Histopathological and Proteomics Study

Soudabeh Balarastaghi^a, Rezvan Yazdian-Robati^b, Faezeh Vahdati Hasani^a, Hossein Hosseinzadeh^{c, a}, Khalil Abnous^{c, d}, Mohsen Imenshahidi^a, Leila Mohammadzadeh^e, Ruth Birner-Gruenberger^f and Bibi Marjan Razavi^{g, a*}

^aDepartment of Pharmacodynamics and Toxicology, School of Pharmacy, Mashhad University of Medical Sciences, Mashhad, Iran. ^bMolecular and Cell Biology Research Center, Faculty of Medicine, Mazandaran University of Medical Sciences, Sari, Iran. ^cPharmaceutical Research Center, Pharmaceutical Technology Institute, Mashhad University of Medical Sciences, Mashhad, Iran. ^dDepartment of Medicinal Chemistry, School of Pharmacy, Mashhad University of Medical Sciences, Mashhad, Iran. ^eFood and Drug Control Laboratory, Food and Drug Vice Presidency, Mashhad University of Medical Sciences, Mashhad, Iran. ^fInstitute of Pathology, Research Unit Functional Proteomics and Metabolic Pathways, Medical University of Graz, Graz, Austria. ^gTargeted Delivery Research Center, Pharmaceutical Technology Institute, Mashhad University of Medical Sciences, Mashhad, Iran.

Abstract

In this study, the protective effect of crocin on malathion (MTN) induced cardiotoxicity in rats in subacute exposure was evaluated. Rats were divided into 6 groups; control (normal saline); MTN (100 mg/kg); MTN + crocin (10, 20 and 40 mg/kg) and MTN + vitamin E 200 IU/kg. Treatments were continued for two weeks. Creatine phosphokinase MB (CK-MB), malondialdehyde (MDA) and glutathione (GSH) levels were evaluated in heart tissue at the end of treatments. The effect of crocin and MTN on histopathological changes in rat cardiac tissue was also investigated. The alteration of protein profile in the heart of the animals exposed to MTN was evaluated by proteomic approach through two-dimensional gel electrophoresis followed by matrix-assisted laser desorption/ionization-time of flight (MALDI-TOF) software. MTN induced histopathological damages and elevated the level of cardiac marker CK-MB ($P < 0.01$). The level of MDA increased and the level of GSH reduced ($P < 0.001$). MDA levels were reduced in all crocin plus MTN groups ($P < 0.001$) and vitamin E plus MTN ($P < 0.001$) groups as compared to MTN groups. However, in the crocin (10 mg/kg) + MTN group, the content of GSH compared to MTN treated rats increased ($P < 0.001$). Protein abundance analysis identified proteins implicated in cardiac necrosis, tricarboxylic acid cycle, cellular energy homeostasis, arrhythmias, heart development, heart failure and cardiovascular homeostasis to be affected by MTN. In summary, MTN may induce damage in the heart tissue of rats following subacute exposure and crocin, as an antioxidant, showed protective effects against MTN cardiotoxicity.

Keywords: Malathion; Crocin; Cardiotoxicity; Proteomics; Oxidative stress.

* Corresponding author:
E-mail: razavimr@mums.ac.ir

Introduction

Organophosphate pesticides (OP), being cholinesterase inhibitors, are extensively used to control domestic and farming pests globally, especially developing countries (1). The common use of OP pesticides by community agricultural and health programs has led to severe environmental pollution that constitutes a potential health hazard by chronic or acute poisoning of humans (2). The residual amounts of these pesticides were detected in various agents like soil, water, fruits, vegetables, and other food products (3).

The most important mechanism of OP poisoning is acetylcholine accumulation as a result of AChE inhibition (4). The induction of oxidative stress in animals and humans has been observed following exposure to OPs, which leads to the free radicals generation and then adaptive changes in the antioxidant defense system. The increased formation of reactive oxygen species (ROS) has been approved in the pathology of cardiovascular disease, dementia, cancer and metabolic disorders (5, 6). It has been proved that oxidative stress is considered one of the most important toxicity mechanisms induced by OPs in long-term exposures (7). Malathion (MTN) [O, O-dimethyl-S-(1,2-dicarcethoxyethyl)phosphorodithioate] is one of the most common OP pesticides which has been found in household and agricultural products and is widely used for pest control (8). The widespread usage of this pesticide makes it particularly vital to detect any related health risks. Several acute and chronic MTN poisoning among pesticide workers and children following accidental exposure have been reported. It is well documented that environmental contamination with MTN causes adverse health effects in humans, animals and wildlife (9).

Malathion is metabolized to malaoxon in the body, and this compound is 61 times more toxic than malathion (10). Its harmful effects have been demonstrated in various organisms, including honeybees and aquatic organisms, causing broad-ranging deformities (11). Therefore it is assumed that MTN poses a major risk for many organisms, including humans (12). Several different mechanisms

of toxicity of MTN have been proposed, including inhibition of acetylcholinesterase activity in target tissues, several organ system dysfunctions including liver, pancreas, and the reproductive system, as well as changes in lipid metabolism, such as the elevation of low-density lipoprotein (LDL) and triglyceride levels in the blood (2,13-15). Besides, oxidative stress has been suggested as an alternative molecular mechanism for MTN toxicity: MTN increased MDA levels, which is an important marker of oxidative stress and the end product of lipid peroxidation, and it decreased the level of reduced GSH in the rat cortex and hippocampus (16). Crocin (crocetin digentiobiose ester), a unique water-soluble carotenoid, is one of the pharmacologically active ingredients of saffron (*Crocus sativus*) (17, 18). Several animal and clinical studies have revealed beneficial effects of saffron extracts and crocin, such as anti-tumor, anti-nociceptive, aphrodisiac, antidepressant, anxiolytic, anti-Alzheimer, anti-diabetic maculopathy, *etc.* through antioxidant and anti-inflammatory mechanisms (19-27).

The cardioprotective effects of saffron and its components such as crocetin and crocin have been reported in different studies due to the modulation of endogenous antioxidant enzymatic activities (17). Previously our group has reported the protective effects of crocin on diazinon and acrylamide toxicities (28, 29). According to several studies, crocin, the main coloring pigment of *Crocus sativus* (*C. sativus*), is safe. Hosseinzadeh *et al.* (2010) evaluated biochemical, hematological and pathological effects of crocin (up to 3 g, p.o. and i.p. as well as 15–180 mg/kg, i.p.) in mice and rats. The results showed that crocin at pharmacological doses did not exhibit any toxicity in different organs (30).

Since many literature data reported that OPs elicit different severe acute and long-term adverse effects and as far as we know, the toxic effects of subacute exposure of malathion on the cardiac system are not clear (31-34). Thus, this study was designed to determine the possible toxic effects of MTN on a specific biochemical cardiac enzyme (CK-MB), the MDA and glutathione levels, and morphological changes of the rat heart. Moreover, the possible protective effects of

crocin on MTN toxicity were investigated. A proteomics method using two-dimensional gel electrophoresis and MALDI-TOF/TOF was used to identify differentially expressed proteins in rat hearts caused by the subacute administration of MTN. This is the first proteomics study that investigates protein changes caused by administration of malathion in the heart tissue to the best of our knowledge.

Experimental

Chemicals

MTN was prepared from Ariashimi co., Tehran, Iran, and vitamin E was purchased from OSVE Pharmaceutical Co., Tehran, Iran. 2-thiobarbituric acid (TBA), potassium chloride, n-butanol, phosphoric acid and MDA were obtained from Merck, Germany. Reduced GSH were purchased from Sigma Aldrich, USA. Stigmas of *C. sativus* L. from Novin Saffron (collected from Ghaen, Khorasan province, Iran) was obtained and analyzed in accordance to the ISO/TS 3632-2. Crocin was extracted and purified as defined by Hadizadeh and colleagues (35).

Animals and treatment

Adult male Wistar rats (weight 220–250 g) were supplied by the animal center of the School of Pharmacy of Mashhad University of Medical Sciences. Rats were maintained at a temperature of 23 ± 1 °C and on a 12 h light/dark cycle with free access to water and food during the experiments. Animal experiments were done according to the Mashhad University of Medical Sciences Ethical Committee Acts (941030). Rats were divided into 6 groups of 6 rats each: control group (normal saline); MTN treated group (100 mg/kg); MTN + crocin 10 mg/kg/day treated group; MTN + crocin 20 mg/kg/day treated group; MTN + crocin 40 mg/kg/day treated group; MTN + vitamin E 200 IU/kg treated group. MTN and crocin were administered daily via intraperitoneal injection (IP) for two weeks. Normal saline was given in the same way to control groups. Vitamin E was administrated intraperitoneally three days a week (36).

Biochemical evaluation

After two weeks, rats were sacrificed and blood samples were collected. Serum was

prepared for biochemical tests. Heart tissues were removed, washed in normal saline and flash-frozen at -80 °C.

Cholinesterase activity assay

Acetylcholinesterase activity in serum was measured by the Ellman method. The rate of hydrolysis of acetylthiocholine iodide was measured at 405 nm using the release of the thiol compound which produces a color-forming compound upon reaction with 5, 5-dithiobis-(2-nitrobenzoic acid) (DTNB). The absorbance was recorded at 0.5 min time intervals for 2 min. Cholinesterase activity was calculated as follows: Cholinesterase activity (mU/ml serum, at 25 °C) = change in absorbance in 30 s \times 23400 (37).

Measurement of MDA in the heart tissue

MDA levels were determined according to the method of Fernandez *et al.* (38). This method is based on the spectrophotometric measurement of the color developed by the reaction of MDA to thiobarbituric acid (TBA). Briefly, heart tissues were homogenized for 2 min at 4 °C in 1.15% potassium chloride (KCl) solution to provide a 10% W/V homogenate. Three milliliter of phosphoric acid (H₃PO₄) (1% W/V) and 1 of TBA (0.6% W/V) were then added to 0.5 of the homogenate and the mixture was incubated for 45 min in a boiling water bath. After cooling with cold water, n-butanol (4 mL) was added and then the mixture stirred for 1 min. After vortex, the mixture was centrifuged for 20 min at 3000 \times g. The upper layer was transferred to the new tube and absorption of this layer was measured by the spectrometry device at 532 nm. The amount of MDA is reported as nmol/g tissue.

Measurement of GSH in the heart tissue

GSH content was measured based on the method of Moron *et al.* (39). Hearts were homogenized in ice-cold phosphate-buffered saline (PBS, pH 7.4) to obtain a 10% (w/v) homogenate. The reduced glutathione content was measured using (5, 5'-dithiobis-(2-nitrobenzoic acid) (DTNB), which produced a yellow-colored 5-thio-2-nitrobenzoic acid (TNB). Then 10% trichloroacetic acid (TCA) and the equal amount of sample were mixed and centrifuged at 3000 \times g for 5 min. 0.5 of 0.04% DTNB reagent and 2PBS (0.1 M,

pH 8.0) were added to 0.5 of supernatants (0.1 M, pH 8.0). And the absorption of the yellow-colored TNB was measured by the spectrometry apparatus at 412 nm. Tissue GSH content is reported as nmol/g tissue

Measurement of creatine phosphokinase-MB (CK-MB)

The CK-MB activity in serum was measured using commercial colorimetric kits (Biosystem, Spain) by the auto-analyzer (Tokyo-Boeki Prestige).

Histopathological evaluation

For histopathological evaluation, all animals' hearts were excised at the end of the experiment, fixed in 10% v/v neutral buffered formalin and placed in paraffin wax. Tissue parts were cut in 6 μ m slices and stained with hematoxylin and eosin. Histopathological assessments of the prepared tissues were performed using Olympus light microscope (BH-2, Japan), and the photos were taken with a digital camera attached to the microscope (Olympus, C-7070).

Two-dimensional gel electrophoresis sample preparation

Protein extraction

Frozen heart tissues (0.2 g) from each group (control and MTN 100 mg/kg treated groups) were homogenized in 1 mL cold lysis buffer containing 2 M thiourea, 6 M urea, 2% (w/v) CHAPS, 2% (w/v) DTT, 20% (w/v) ampholyte (pH 3–10, Bio-Rad) and Halt™ Protease Inhibitor Cocktail using Polytron Homogenizer (IKA®T10, Germany). Using the Bio-Rad Protein Assay Kit and based on the Protocol factory, the concentration of supernatants was measured as described before (40).

Two-dimensional gel electrophoresis

For isoelectric focusing (IEF), was dissolved in 300 μ L of the rehydration buffer (6 M urea, 0.3% w/v DTT, 2 M thiourea, 4% w/v CHAPS and 20% (w/v) ampholyte (pH 3–10) containing 300 μ g of total protein, loaded on to each 17 cm 3–10 nonlinear IPG strips from Bio-Rad, USA, and actively rehydrated for 12 h at 50 V (37). IEF was performed using Protean IEF Cell (Bio-Rad, USA) at 20 °C

and the following program: voltage started at 250 V, then linearly increased to 8000 V and kept constant at this voltage for 11 h. The current limit was set at 50 μ A/gel. Then, the IPG strips were equilibrated in 6 mL of reduction buffer containing 37.5 mM Tris-HCl (pH 8.8), 6 M urea, 20% glycerol, 2% SDS, and 2% DTT for 15 min. For alkylation, the strips were equilibrated in 6 mL of solution containing 37.5 mM Tris-HCl (pH8.8), 6M urea, 20% glycerol, 2% SDS, and 2.5% w/v iodoacetamide for 15 min. After that, the strips were placed on top of 12% SDS-PAGE gels for the second dimension with the separation done at 120 V for 5 h, and then at 150 V until the tracking dye reached the end of the gel (41). After electrophoresis, gels were stained with colloidal Coomassie for 12 h (42).

Analysis of images in-gel digestion and mass spectrometry

After the staining stage (3 biological replicates/group), the gels were scanned by Image scanner III (Epson, Japan). Then, to find the different proteins expressed differently, the software Image Master Platinum 6.0 software was used (GE Healthcare, USA). Protein spots with more than a 1.5 fold change and significantly different intensities ($P < 0.05$) in control and MTN treated groups were considered as differentially expressed and selected for excision and mass spectrometric identification (43). Protein spots on the 2-D gels were excised from gels, transferred into a 1.5 mL microtube containing 1% acetic acid and sent to the Centre of Excellence in Mass Spectrometry at the University of York, UK, for in-gel digestion and identification by MALDI-TOF/TOF mass spectrometry. Gel pieces were reduced with DTT and treated with iodoacetamide for S-carbamidomethylation. They were then washed twice with 50% (v/v) aqueous acetonitrile (ACN) containing 25 mM ammonium bicarbonate and once with 100% ACN. After removing ACN, gel slices were dried in a vacuum concentrator for 20 min and rehydrated with 10 μ L of 0.02 μ g/ μ L sequencing-grade modified porcine trypsin (Promega). After 5 min, enough 25 mM ammonium bicarbonate solution) was added to cover the gel pieces. Digests were incubated at 37 °C overnight. The next day, extracted

peptide mixtures (1 μ L) were applied directly onto the ground steel MALDI target plate, followed immediately by mixing with 1 μ L of a freshly-prepared 5 mg/mL of 4-hydroxy- α -cyanocinnamic acid (Sigma) solution in 50% aqueous (v/v) ACN, 0.1% trifluoroacetic acid (v/v). The BrukerUltraflex III in reflectron mode, equipped with an Nd: YAG smart beam laser, was applied for obtaining positive-ion MALDI mass spectra. MS spectra were acquired over a mass range of m/z 800–5000 and externally calibrated against an adjacent spot containing six standard peptides (Angiotensin I, 1296.685; des-Arg1-Bradykinin, 904.681; Glu1-Fibrinopeptide B, 1750.677; ACTH (18–39 clip), 2465.198; ACTH (1–17clip), 2093.086; ACTH (7–38 clip), 3657.929). For MS/MS fragmentation, the ten strongest peaks of interest with an S/N greater than 30 were chosen. Fragmentation was done in LIFT mode without the introduction of a collision gas. Spectral processing and peak list generation for both the MS and MS/MS spectra were performed by Brukerflex Analysis software (version 3.3).

Tandem mass spectral data were submitted to database search using a locally-running copy of the Mascot software (Matrix Science Ltd., version 2.5) through the Bruker BioTools interface (version 3.2). Search criteria included Enzyme Trypsin; Fixed modifications, Carbamidomethyl (C); Variable modifications, Oxidation (M); Peptide tolerance, 100 ppm; MS/MS tolerance, 0.5 Da; Instrument, MALDI-TOF/TOF (42).

Classification of proteins

The PANTHER online database (<http://www.pantherdb.org>) was utilized for functional classification of significantly altered proteins. PANTHER is a library of protein families and Subfamilies that demonstrate the biological functions, protein classes, and pathways (44).

Statistical analysis

Statistical analysis of MDA, GSH, and CK-MB as was done by One-way Analysis of Variance (ANOVA) followed by Tukey post-hoc test using Graph Pad Prism version 6.00 for Windows, GraphPad Software. Protein intensities, expressed as a percent of total gel staining, were analyzed using Student's

t-test (GraphPad Prism version 6.00). All data were reported as mean \pm standard error of the mean (SEM). *P*-values less than 0.05 were considered statistically significant.

Results

Biochemical evaluation

A significant reduction was observed in serum cholinesterase activity in the MTN group compared to the control group (499.3 ± 23.2 vs. 249 ± 3.005) ($P < 0.001$), but there was no significant difference between MTN plus crocin and MTN groups (Figure 1).

A significant increase was detected in MDA level in the MTN treated group in comparison with the control group (271.2 ± 11.51 vs. 35 ± 1.59) ($P < 0.001$) (Figure 2). MDA levels were significantly reduced in crocin 10 mg/kg plus MTN (29.44 ± 2.11), crocin 20 mg/kg plus MTN (38.82 ± 3.5), crocin 40 mg/kg plus MTN (52.5 ± 9.56) ($P < 0.001$) and vitamin E plus MTN (43.5 ± 2.02) ($P < 0.001$). In MTN treated rats, cardiac content of GSH was significantly decreased compared to control (722 ± 11.07 vs. 34 ± 4.63) ($P < 0.001$, Figure 3). However, in crocin 10 mg/kg + MTN (269 ± 41.92), crocin 20 mg/kg + MTN (407 ± 7.24) and crocin 40 mg/kg + MTN (337 ± 25.2) the content of GSH compared to MTN treated rats significantly increased ($P < 0.001$). Serum CK-MB activity showed high cardiotoxicity in MTN treated group (1156.85 ± 122.8 vs. $657.15 \pm 43 \pm 9.6$) ($P < 0.01$, Figure 4). Co-administration of 40 mg/kg of crocin reduced serum CK-MB (647.4 ± 60.92) activity as compared with the MTN group ($P < 0.01$).

Histopathological analysis

Histopathological changes of cardiac tissues are shown in Figure 5. Histological findings were normal in the control group (Figure 5A). Histological changes were observed in MTN treated group (Figure 5B) as mild focal inflammation, myocytolysis and granular degeneration. The cardiac lesions were significantly improved in the MTN plus crocin (10, 20 and 40 mg/kg) or vitamin E compared to the MTN-treated group (Figure 5).

Proteomics

Differentially expressed proteins regulated by MTN exposure (100 mg/kg) and control

in the rat heart tissue were identified and analyzed by 2-DE and MALDI TOF/TOF. Three rats from each group were randomly selected and subjected to 2-DE analysis. Representative examples of the 2-DE gel images of the proteins of the heart from MTN exposed rats are reported in Figures 6 and 7.

MTN-therapy induced changes of the protein and expression pattern in the heart proteome. Statistical analysis of differential protein expression revealed that seven total protein spots were differentially regulated between control and MTN-treated groups. Analysis of the total protein expression profile showed that 6 proteins were down-regulated while only one of them increased in the MTN-treated livers compared to the control samples. Table 1 provides more information about the differentially expressed proteins

exposure to MTN, including SwissProt accession number, fold-change, MASCOT protein score, matched peptide sequenced, expectation-value and a brief description the biological function or molecular pathways.

Pathway analysis

Using the PANTHER database (<http://panther.appliedbiosystems.com/>), the molecular function, biological process and signaling pathway of each identified protein was investigated. Differentially expressed proteins were classified in 9 pathways including; localization 3%, cellular process 23%, reproduction 1%, cellular component organization 6%, biological regulation 6%, developmental process 1%, metabolic process 53% and response to stimulus 6% (Figure 8).




Figure 1. Effect of malathion and crocin treatment (2 weeks) on AChE activity. Data are presented as mean \pm SEM. ^{###} $P < 0.001$ compared to control. Tukey Kramer test, $n = 6$, MTN (malathion).




Figure 2. Effect of malathion and crocin treatment (2 weeks) on MDA Level in the heart rat tissues. Data are shown as mean \pm SEM, *** P < 0.001 compared to malathion group, ### P < 0.001 compared to control group. Tukey–Kramer test, n = 6.




Figure 3. Effects of malathion and crocin treatment (2 weeks) on GSH level in the heart rat tissues. Data are shown as mean \pm SEM, *** P < 0.001 compared to malathion group, ### P < 0.001 compared to control group. Tukey–Kramer test, n = 6.




Figure 4. Effects of malathion and crocin treatment (2 weeks) on CK-MB activity. Data are shown as mean \pm SEM, ** P < 0.01 and * P < 0.05 compared to malathion group, ## P < 0.01 compared to control group. Tukey–Kramer test, n = 6.




Figure 5. (A) Control rats: normal cardiac muscle cells, hematoxylin and eosin, $\times 400$. (B) Malathion-treated rats: mild focal inflammation, myocytolysis and granular degeneration, hematoxylin and eosin, $\times 400$. (C) Crocin 10 mg/kg+ malathion treated rats: normal cardiac muscle cells, hematoxylin and eosin, $\times 400$. (D) Crocin 20 mg/kg+ malathion treated rats: myocytolysis, hematoxylin and eosin, $\times 400$. (E) Crocin 40 mg/kg + malathion treated rats: mild focal inflammation, hematoxylin and eosin, $\times 400$. (F) Vitamin E+ malathion treated rats: myocytolysis, hematoxylin and eosin, $\times 400$.




Figure 6. Two-dimensional gel electrophoresis of malathion treated rat heart (1 = Tropomyosin, 2 = ATP synthase, 3 = Desmin, 4 = L- lactate dehydrogenase, 5 = Malate dehydrogenase, 6 = Aconitate hydratase).




Figure 7. Two-dimensional gel electrophoresis of untreated rat heart.




Figure 8. Functional classification of the differentially expressed proteins in the heart regulated by MTN exposure according to biological processes using the PANTHER classification system. Numbers indicate the percentage of protein against the total number of proteins.

Discussion

MTN is one of the most broadly used organophosphate compounds applied in agriculture as a pesticide, as an ectoparasiticide in veterinary practice, control of public health programs, and food preparation and processing areas (45, 46). In the present study, MTN (100 mg/kg) exhibited cardiac toxicity in subacute exposure (14 days) by inducing morphological damages and elevating the cardiac marker CK-MB level in serum. It has been shown that chronic exposure to OPs introduces oxidative stress damage and free radical production in different organs, such as the cardiovascular system (28, 47 and 48). Oxidative stress in cardiac tissues appeared to be responsible for these morphological and biochemical changes since co-administration with crocin in all doses of vitamin E was protective against MTN-induced cardiac toxicity by decreasing oxidative stress. Several studies reported that subacute exposure to OPs, such as MTN, inhibited cholinesterase enzyme activity and caused oxidative stress. Moreover, MTN was shown to produce free radicals in different organs, such as the cardiovascular system (29, 49-51). Our data revealed that the cardiac level of MDA was significantly enhanced after exposure to MTN. MDA as the common product of lipid peroxidation is widely used to assess the presence of free radicals and lipid peroxidation in different toxicity and disease studies, including in cardiovascular toxicity (52-54). Another biochemical marker we also measured in this work was GSH. GSH is an antioxidant marker, significantly reduced in MTN-treated group compared to the control group. In the present study, vitamin E or α -tocopherol, the biologically most active antioxidants, were utilized as positive controls. Our histopathological findings indicated that sub-chronic exposure to MTN could induce congestion, infiltration of inflammatory cells and multifocal necrosis in cardiac tissue. Crocin as a potent antioxidant agent showed cardioprotective effects (29, 55). Co-administration of Crocin and MTN as compared to MTN alone decreased the cardiac MDA level and the serum CK-MB concentration and increased the GSH content in the heart tissue. Crocin also protected the

heart from histological changes induced by MTN, probably via its antioxidative and anti-inflammatory properties (56, 57). The protective effect of crocin may also be related to its inhibitory effect on lipid peroxidation leading to stabilization of membranes thereby blocking the release of cardiac enzymes (56). A proteomics approach was employed in the present study to identify if protein expression was changed in response to subacute exposure to MTN (Table 1). Seven proteins were found to be differentially regulated following MTN exposure compared to the control group. Our results confirmed a good agreement between well-known biomarkers of human cardiac disease and cellular response to MTN. Besides, using the proteomic approach, we revealed that mitochondrial proteins showing differential expression in response to MTN. ATP synthase, aconitate hydratase, malate dehydrogenase, and L-lactate dehydrogenase (LDH) as mitochondrial enzymes changed significantly in the MTN-exposed group. Involvement of mitochondrial injuries following exposure to OPs has been reported by many researchers (27, 58). Multiple lines of evidence have demonstrated that OPs could alter mitochondrial respiration and respiratory chain enzyme activity (59, 60). Mitochondria have a vital role in ATP production and the ROS, so dysfunction of mitochondria in the OPs toxicity involved in multisystem disorders in different organs including liver, heart and neurons (61, 62). Indeed, changes in the levels of these mitochondrial enzymes may be the most relevant signal of heart toxicity of MTN (36). LDH, as an important glycolytic enzyme, is a potential biomarker in toxicology and clinical chemistry if detected extracellularly as a sign of cell, tissue and organ damage (63). Previously, it has been demonstrated that the LDH activity altered after exposure to toxic compounds or oxygen stress (64-67). Venkatesan *et al.* have also reported that malathion treatment enhanced LDH release in a dose-dependent manner and induced neurotoxicity in N2a neuroblastoma cells (68). Elevated LDH also entirely consistent with that observed by Kalender *et al.* (69). Increased LDH in the malathion-treated group in the present study may indicate toxic effects of MTN in the heart. Multiple studies have

Table 1. Differentially expressed protein species in the heart of rats exposed to malathion as identified by MALDI-TOF/TOF and MASCOT software.

Protein name	Swiss Prot accession number	Fold-change (MTN/control)	Protein score	%sequence coverage	Matched peptides	Expectation-value	Matched peptide sequenced	Biological function or Molecular pathways (PANTHER classification)
Calreticulin	P18418	0.60	92	6	2(2)	0.0012	K.EQLDGDGDAWTNR.W	Metabolic process Cellular component organization or biogenesis, cellular process, developmental process, metabolic process, multicellular organismal process
Tropomyosin	P58775	0.24	754	29%	9(9)	1e-10	K.KATDAEADVASLNR.R	
L-lactate dehydrogenase	P42123	1.57	513	19%	7(7)	8e-11	K.IVADKDYSVTANSK.I	Metabolic process
ATP synthase	P15999	0.56	1004	25%	10(10)	5.6e-11	R.EVAFAQFGSDLDAATQQLSR.G	ATP synthesis
Malate dehydrogenase	O88989	0.26	549	22%	7(7)	7.1e-12	K.ELTEKETAFEFLLSSA.-	Metabolic process
Aconitate hydratase, mitochondrial	Q9ER34	0.6	847	15%	10(10)	2.5e-10	K.FKLEAPDADELPR.S	Metabolic process ,TCA cycle
Desmin	P48675	0.56	751	21%	9(9)	3e-11	R.FASEASGYQDNIAR.L	developmental process cellular component organization or biogenesis

Discussion

MTN is one of the most broadly used organophosphate compounds applied in agriculture as a pesticide, as an ectoparasiticide in veterinary practice, control of public health programs, and food preparation and processing areas (45, 46). In the present study, MTN (100 mg/kg) exhibited cardiac toxicity in subacute exposure (14 days) by inducing morphological damages and elevating the cardiac marker CK-MB level in serum. It has been shown that chronic exposure to OPs introduces oxidative stress damage and free radical production in different organs, such as the cardiovascular system (28, 47 and 48). Oxidative stress in cardiac tissues appeared to be responsible for these morphological and biochemical changes since co-administration with crocin in all doses of vitamin E was protective against MTN-induced cardiac toxicity by decreasing oxidative stress. Several studies reported that subacute exposure to OPs, such as MTN, inhibited cholinesterase enzyme activity and caused oxidative stress. Moreover, MTN was shown to produce free radicals in different organs, such as the cardiovascular system (29, 49-51). Our data revealed that the cardiac level of MDA was significantly enhanced after exposure to MTN. MDA as the common product of lipid peroxidation is widely used to assess the presence of free radicals and lipid peroxidation in different toxicity and disease studies, including in cardiovascular toxicity (52-54). Another biochemical marker we also measured in this work was GSH. GSH is an antioxidant marker, significantly reduced in MTN-treated group compared to the control group. In the present study, vitamin E or α -tocopherol, the biologically most active antioxidants, were utilized as positive controls. Our histopathological findings indicated that sub-chronic exposure to MTN could induce congestion, infiltration of inflammatory cells and multifocal necrosis in cardiac tissue. Crocin as a potent antioxidant agent showed cardioprotective effects (29, 55). Co-administration of Crocin and MTN as compared to MTN alone decreased the cardiac MDA level and the serum CK-MB concentration and increased the GSH content in the heart tissue. Crocin also protected the

heart from histological changes induced by MTN, probably via its antioxidative and anti-inflammatory properties (56, 57). The protective effect of crocin may also be related to its inhibitory effect on lipid peroxidation leading to stabilization of membranes thereby blocking the release of cardiac enzymes (56). A proteomics approach was employed in the present study to identify if protein expression was changed in response to subacute exposure to MTN (Table 1). Seven proteins were found to be differentially regulated following MTN exposure compared to the control group. Our results confirmed a good agreement between well-known biomarkers of human cardiac disease and cellular response to MTN. Besides, using the proteomic approach, we revealed that mitochondrial proteins showing differential expression in response to MTN. ATP synthase, aconitate hydratase, malate dehydrogenase, and L-lactate dehydrogenase (LDH) as mitochondrial enzymes changed significantly in the MTN-exposed group. Involvement of mitochondrial injuries following exposure to OPs has been reported by many researchers (27, 58). Multiple lines of evidence have demonstrated that OPs could alter mitochondrial respiration and respiratory chain enzyme activity (59, 60). Mitochondria have a vital role in ATP production and the ROS, so dysfunction of mitochondria in the OPs toxicity involved in multisystem disorders in different organs including liver, heart and neurons (61, 62). Indeed, changes in the levels of these mitochondrial enzymes may be the most relevant signal of heart toxicity of MTN (36). LDH, as an important glycolytic enzyme, is a potential biomarker in toxicology and clinical chemistry if detected extracellularly as a sign of cell, tissue and organ damage (63). Previously, it has been demonstrated that the LDH activity altered after exposure to toxic compounds or oxygen stress (64-67). Venkatesan *et al.* have also reported that malathion treatment enhanced LDH release in a dose-dependent manner and induced neurotoxicity in N2a neuroblastoma cells (68). Elevated LDH also entirely consistent with that observed by Kalender *et al.* (69). Increased LDH in the malathion-treated group in the present study may indicate toxic effects of MTN in the heart. Multiple studies have reported an association

between ATP synthase and exposure to the OPs pesticide (31, 33 and 34). Decreases in ATP synthase level in OPs treated groups compared to control supported by several studies (70, 71). Acute exposure to MTN in a rat model induced cardiac failure, diminished ATP production and affected ADP/ATP ratio in heart tissue cell line (32). Decreased ATP synthase as observed in response to MTN may reflect the inability of the cells to maintain a normal energy level (72). Down-regulation of aconitate hydratase suggesting overproduction of superoxide anion and hydrogen peroxide in rat heart mitochondria in subacute intoxication with MTN (58).

Upon exposure to MTN, the level of malate dehydrogenase- a key enzyme in the Krebs cycle- in rat heart tissue was significantly reduced. In the present study, this reduction in malate dehydrogenase indicates a decrease in the respiration rate due to toxicity induced by MTN (73). Decreased level of malate dehydrogenase results in lesser production of ATP in the mitochondria which may prevent numerous important metabolic functions in the animals (74). Calreticulin, a Ca²⁺-binding chaperone in the endoplasmic reticulum (ER), is a regulator of calcium homeostasis through modulation of ER Ca²⁺ storage and transport (75). Furthermore, calreticulin affects many aspects of cellular processes, including cardiac development, muscle contraction, cell adhesion, adipocyte differentiation, innate immunity, steroid-sensitive gene expression, apoptosis, and stress responses (76, 77). It is possible that decreased calreticulin in the heart tissue following MTN treatment could be responsible for lipid production, previously proposed (69). In our proteomic study, reduced-expression of calreticulin was detected after exposure to MTN. Tropomyosin and desmin were found to decrease in the heart after MTN administration. Tropomyosins are actin filament binding proteins, possessing significant roles in regulating cardiovascular homeostasis and modulation of endothelial cell function (78). Desmin, as a muscle-specific protein and a crucial intermediate filament in the heart, involves inhibition of cardiac conduction, cell architecture, arrhythmias, mitochondrial function and behavior and restrictive heart failure (70, 79).

It has been reported a heart without desmin is characterized by cardiomyocyte degeneration and a dilated cardiomyopathy (71).

Conclusion

In conclusion, our data indicate that crocin has protective effects against MTN-induced cardiotoxicity through attenuating lipid peroxidation and increasing GSH content. Protein analysis identified proteins implicated in cardiac necrosis, tricarboxylic acid cycle, cellular energy homeostasis, arrhythmias, heart development, heart failure and cardiovascular homeostasis to be affected by MTN.

Acknowledgments

We thank Mashhad University of Medical Sciences for financial supports.

References

- (1) Kalender S, Uzun F G, Durak D, Demir F and Kalender Y. Malathion-induced hepatotoxicity in rats: the effects of vitamins C and E. *Food Chem. Toxicol.* (2010) 48: 633-8.
- (2) Lasram MM, Annabi AB, El Elj N, Selmi S, Kamoun A, El-Fazaa S and Gharbi N. Metabolic disorders of acute exposure to malathion in adult Wistar rats. *J. Hazard. Mat.* (2009) 163: 1052-5.
- (3) Kalender S, Ogutcu A, Uzunhisarcikli M, Açikgoz F, Durak D, Ulusoy Y and Kalender Y. Diazinon-induced hepatotoxicity and protective effect of vitamin E on some biochemical indices and ultrastructural changes. *Toxicology* (2005) 211: 197-206.
- (4) Bajgar J. Organophosphates/nerve agent poisoning: mechanism of action, diagnosis, prophylaxis, and treatment. *Adv. Clin. Chem.* (2004) 38: 151-216.
- (5) Akhgari M, Abdollahi M, Kebryaezadeh A, Hosseini R and Sabzevari O. Biochemical evidence for free radical induced lipid peroxidation as a mechanism for subchronic toxicity of malathion in blood and liver of rats. *Human Exp. Toxicol.* (2003) 22: 205-11.
- (6) Madhavan Nampoothiri G, Varkey J and Noufal C. Sub-acute malathion induced cardiotoxicity in wistar rats: protective effect of vitamin c. *Int. J. Biol. Pharm. Res.* (2012) 3: 779-83
- (7) Abdollahi M, Ranjbar A, Shadnia S, Nikfar S and Rezaiee A. Pesticides and oxidative stress: a review. *Med. Sci. Monit.* (2004) 10: 141-7.
- (8) Assini FL, Zanette KD, Brocardo PS, Pandolfo P, Rodrigues ALS and Takahashi RN. Behavioral effects and ChE measures after acute and repeated

- administration of malathion in rats. *Env. Toxicol. Pharmacol.* (2005) 20: 443-9.
- (9) Gangemi S, Gofita E and Fenga C. Occupational and environmental exposure to pesticides and cytokine pathways in chronic diseases (Review). *Int. J. Mol. Med.* (2016) 34: 1012–20.
- (10) Edwards D. Reregistration eligibility decision for malathion. *US Environmental Protection Agency-Prevention, Pesticides and Toxic Substances EPA* (2006) 9: 738.
- (11) Newhart K. Environmental fate of malathion. *California Environmental Protection Agency.* (2006) 11: 1-20.
- (12) Simoneschi D, Simoneschi F and Todd NE. Assessment of cardiotoxicity and effects of malathion on the early development of zebrafish (*Danio rerio*) using computer vision for heart rate quantification. *Zebrafish* (2014) 11: 275-80.
- (13) Rezg R, Mornagui B, El-Fazaa S and Gharbi N. Biochemical evaluation of hepatic damage in subchronic exposure to malathion in rats: effect on superoxide dismutase and catalase activities using native PAGE. *Comptes. Rendus. Biol.* (2008) 331: 655-62.
- (14) Vosough-Ghanbari S, Sayyar P, Pournourmohammadi S, Aliahmadi A, Ostad S N and Abdollahi M. Stimulation of insulin and glucagon synthesis in rat Langerhans islets by malathion *in-vitro*: Evidence for mitochondrial interaction and involvement of subcellular non-cholinergic mechanisms. *Pesticide. Biochem. Physiol.* (2007) 89: 130-6.
- (15) Bustos-Obregón E and González-Hormazabal P. Effect of a single dose of malathion on spermatogenesis in mice. *Asian. J. Androl.* (2003) 5: 105-7.
- (16) Mohammadi H, Karimi G, Mahdi Rezayat S, Reza A, Shafiee H, Nikfar S, Baeri M, Sabzevari O and Abdollahi M. Benefit of nanocarrier of magnetic magnesium in rat malathion-induced toxicity and cardiac failure using non-invasive monitoring of electrocardiogram and blood pressure. *Toxicol. Indust. Health* (2011) 27: 417-29.
- (17) Razavi M, Hosseinzadeh H, Abnous K, Motamedshariaty VS and Imenshahidi M. Crocin restores hypotensive effect of subchronic administration of diazinon in rats. *Iran. J. Basic Med. Sci.* (2013) 16: 64-72.
- (18) Rezaee R and Hosseinzadeh H. Safranal: from an aromatic natural product to a rewarding pharmacological agent. *Iran. J. Basic Med. Sci.* (2013) 16: 12-26.
- (19) Potnuri A, Allakonda L and Lahkar M. Crocin attenuates cyclophosphamide induced testicular toxicity by preserving glutathione redox system. *Biomed. Pharmacother.* (2018) 101: 174-80.
- (20) Hosseinzadeh H and Younesi HV. Antinociceptive and anti-inflammatory effects of *Crocus sativus* L. stigma and petal extracts in mice. *BMC Pharmacol.* (2002) 2: 7.
- (21) Mansoori P, Akhondzadeh S, Raisi F, Ghaeli P, Jamshidi A and Nasehi A. A randomized, double-blind, placebo-controlled study of safety of the adjunctive saffron on sexual dysfunction induced by a selective serotonin reuptake inhibitor. *J. Med. Plants.* (2011) 1: 121–30.
- (22) Shahmansouri N, Farokhnia M, Abbasi SH, Kassaian SE, Noorbala Tafti AA and Gougol A. A randomized, double-blind, clinical trial comparing the efficacy and safety of *Crocus sativus* L. with fluoxetine for improving mild to moderate depression in post percutaneous coronary intervention patients. *J. Affect. Disord.* (2014) 155: 216–22.
- (23) Hassani FV, Naseri V, Razavi BM, Mehri S, Abnous K and Hosseinzadeh H. Antidepressant effects of crocin and its effects on transcript and protein levels of CREB, BDNF, and VGF in rat hippocampus. *DARU J. Pharm. Sci.* (2014) 22: 16.
- (24) Mazidi M, Shemshian M, Mousavi S H, Norouzy A, Kermani T and Moghiman T. A double-blind, randomized and placebo-controlled trial of saffron (*Crocus sativus* L.) in the treatment of anxiety and depression. *J. Complement. Integr. Med.* (2016) 13: 195–9.
- (25) Akhondzadeh S, Sabet MS, Harirchian MH, Togha M, Cheraghmakani H and Razeghi S. Saffron in the treatment of patients with mild to moderate Alzheimer's disease: a 16-week, randomized and placebo-controlled trial. *J. Clin. Pharm. Ther.* (2010) 35: 581–8.
- (26) Sepahi S, Mohajeri SA, Hosseini SM, Khodaverdi E, Shoebi N, Namdari M and Tabassi SAS. Effects of crocin on diabetic maculopathy: a placebo-controlled randomized clinical trial. *Am. J. Ophthalmol.* (2018) 190: 89-98.
- (27) Hosseini A, Razavi BM and Hosseinzadeh H. Saffron (*Crocus sativus*) petal as a new pharmacological target: a review. *Iran. J. Basic Med. Sci.* (2018) 21: 1091-9.
- (28) Hariri AT, Moallem SA, Mahmoudi M, Memar B and Hosseinzadeh H. Sub-acute effects of diazinon on biochemical indices and specific biomarkers in rats: protective effects of crocin and safranal. *Food Chem. Toxicol.* (2010) 48: 2803-8.
- (29) Razavi BM, Hosseinzadeh H, Movassaghi AR, Imenshahidi M and Abnous K. Protective effect of crocin on diazinon induced cardiotoxicity in rats in subchronic exposure. *Chem-biol. Inter.* (2013) 203: 547-55.

- (30) Hosseinzadeh H, Shariaty VM, Sameni AK and Vahabzadeh M. Acute and sub-acute toxicity of crocin, a constituent of *Crocus sativus* L. (saffron), in mice and rats. *Pharmacologyonline* (2010) 2: 943-51.
- (31) Lari P, Rashedinia M, Abnous K and Hosseinzadeh H. Alteration of protein profile in rat liver of animals exposed to subacute diazinon: a proteomic approach. *Electrophoresis*. (2014) 35: 1419-27.
- (32) Yazdian-Robati R, Pourtaji A, Rashedinia M, Hosseinzadeh H, Ghorbani M, Razavi BM, Ramezani M and Abnous K. Screening and identification of SUMP-proteins in sub-acute treatment with diazinon. *Iran. J. Basic Med. Sci.* (2015) 18: 1240-4.
- (33) Bonfanti P, Colombo A, Orsi F, Nizzetto I, Andrioletti M, Bacchetta R, Mantecca P, Fascio U, Vailati G and Vismara C. Comparative teratogenicity of chlorpyrifos and malathion on *Xenopus laevis* development. *Aqua. Toxicol.* (2004) 70: 189-200.
- (34) Budischak SA, Belden LK and Hopkins WA. Effects of malathion on embryonic development and latent susceptibility to trematode parasites in ranid tadpoles. *Env. Toxicol. Chem: Int. J.* (2008) 27: 2496-500.
- (35) Hadizadeh F, Mohajeri S and Seifi M. Extraction and purification of crocin from saffron stigmas employing a simple and efficient crystallization method. *Pakistan J. Biol. Sci.* (2010) 13: 691-8.
- (36) Mohammadzadeh L, Abnous K, Razavi BM and Hosseinzadeh H. Crocin-protected malathion-induced spatial memory deficits by inhibiting TAU protein hyperphosphorylation and antiapoptotic effects. *Nut. Neurosci.* (2019) 23: 221-36.
- (37) Rashedinia M, Lari P, Abnous K and Hosseinzadeh H. Proteomic analysis of rat cerebral cortex following subchronic acrolein toxicity. *Toxicol. Appl. Pharmacol.* (2013) 272: 199-207.
- (38) Fernández J, Pérez-Álvarez JA and Fernández-López JA. Thiobarbituric acid test for monitoring lipid oxidation in meat. *Food Chem.* (1997) 59: 345-53.
- (39) Moron MS, Depierre JW and Mannervik B. Levels of glutathione, glutathione reductase and glutathione S-transferase activities in rat lung and liver. *Biochem. Biophysica Acta-Gen. Subj.* (1979) 582: 67-78.
- (40) Yazdian-Robati R, Ahmadi H, Riahi MM, Lari P, Aledavood SA, Rashedinia M, Abnous K and Ramezani M. Comparative proteome analysis of human esophageal cancer and adjacent normal tissues. *Iran. J. Basic Med. Sci.* (2017) 20: 265-71.
- (41) Ahmadi H, Ramezani M, Yazdian-Robati R, Behnam B, Azarkhiavi KR, Nia AH, Mokhtarzadeh A, Riahi MM, Razavi BM and Abnous K. Acute toxicity of functionalized single wall carbon nanotubes: a biochemical, histopathologic and proteomics approach. *Chem-biol. Inter.* (2017) 275: 196-209.
- (42) Hassani FV, Abnous K, Mehri S, Jafarian A, Birner-Gruenberger R, Robati RY and Hosseinzadeh H. Proteomics and phosphoproteomics analysis of liver in male rats exposed to bisphenol A: Mechanism of hepatotoxicity and biomarker discovery. *Food Chem. Toxicol.* (2018) 112: 26-38.
- (43) Kowsari R, Yazdian-Robati R, Razavi B M, Pourtaji A, Ghorbani M, Moghadam-Omranipour H, Hosseinzadeh H, Lari P and Abnous K. Recognition and characterization of Erythropoietin binding-proteins in the brain of mice. *Iran. J. Basic Med. Sci.* (2016) 19: 946-52.
- (44) Thomas PD, Campbell MJ, Kejariwal A, Mi H, Karlak B, Daverman R, Diemer K, Muruganujan A and Narechania A. PANTHER: a library of protein families and subfamilies indexed by function. *Genome Res.* (2003) 13: 2129-41.
- (45) Flessel P, Quintana P and Hooper K. Genetic toxicity of malathion: a review. *Env. Mol. Mutagen.* (1993) 22: 7-17.
- (46) Delgado EH, Streck EL, Quevedo JL and Dal-Pizzol F. Mitochondrial respiratory dysfunction and oxidative stress after chronic malathion exposure. *Neurochem. Res.* (2006) 31: 1021-5.
- (47) Ogutcu A, Uzunhisarcikli M, Kalender S, Durak D, Bayrakdar F and Kalender Y. The effects of organophosphate insecticide diazinon on malondialdehyde levels and myocardial cells in rat heart tissue and protective role of vitamin E. *Pest. Biochem. Physiol.* (2006) 86: 93-8.
- (48) Calore E, Perez N and Herman M. Morphometric studies of cardiac myocytes of rats chronically treated with an organophosphate. *Ecotoxicol. Env. Safety* (2007) 66: 447-50.
- (49) Abdollahi M, Mostafalou S, Pournourmohammadi S and Shadnia S. Oxidative stress and cholinesterase inhibition in saliva and plasma of rats following subchronic exposure to malathion. *Compar. Biochem. Physiol. Part C: Toxicol. Pharmacol.* (2004) 137: 29-34.
- (50) Possamai F, Fortunato J, Feier G, Agostinho F, Quevedo J, Wilhelm Filho D and Dal-Pizzol F. Oxidative stress after acute and sub-chronic malathion intoxication in Wistar rats. *Env. Toxicol. Pharmacol.* (2007) 23: 198-204.
- (51) Pournourmohammadi S, Farzami B, Ostad SN, Azizi E and Abdollahi M. Effects of malathion subchronic exposure on rat skeletal muscle glucose metabolism. *Env. Toxicol. Pharmacol.* (2005) 19: 191-6.
- (52) Mahboob M, Rahman M and Grover P. Serum lipid peroxidation and antioxidant enzyme levels in male and female diabetic patients. *Singapore Med. J.*

- (2005) 46: 322-4.
- (53) Kumar CT, Reddy VK, Prasad M, Thyagaraju K and Reddanna P. Dietary supplementation of vitamin E protects heart tissue from exercise-induced oxidant stress. *Mol. Cell. Biochem.* (1992) 111: 109-15.
- (54) Yavuz T, Delibas N, Yildirim B, Altuntas I, Candir O, Cora A, Karahan N, Ibrism E and Kutsal A. Vascular wall damage in rats induced by organophosphorus insecticide methidathion. *Toxicol. Lett.* (2005) 155: 59-64.
- (55) Goyal S, Arora S, Sharma A, Joshi S, Ray R, Bhatia J, Kumari S and Arya D. Preventive effect of crocin of *Crocus sativus* on hemodynamic, biochemical, histopathological and ultrastructural alterations in isoproterenol-induced cardiotoxicity in rats. *Phytomedicine* (2010) 17: 227-32.
- (56) Soeda S, Ochiai T, Shimeno H, Saito H, Abe K, Tanaka H and Shoyama Y. Pharmacological activities of crocin in saffron. *J. Nat. Med.* (2007) 61: 102-11.
- (57) Nam KN, Park YM, Jung HJ, Lee JY, Min BD, Park SU, Jung WS, Cho KH, Park JH and Kang I. Anti-inflammatory effects of crocin and crocetin in rat brain microglial cells. *Eur. J. Pharmacol.* (2010) 648: 110-6.
- (58) Vásquez-Vivar J, Kalyanaraman B and Kennedy MC. Mitochondrial aconitase is a source of hydroxyl radical an electron spin resonance investigation. *J. Biol. Chem.* (2000) 275: 14064-9.
- (59) Potnuri AG, Allakonda L and Lahkar M. Crocin attenuates cyclophosphamide induced testicular toxicity by preserving glutathione redox system. *Biomed. Pharmacother.* (2018) 101: 174-80.
- (60) Lasram MM, Lamine AJ, Dhoub IB, Bouzid K, Annabi A, Belhadjhmida N, Ahmed MB, El Faza S, Abdelmoula J and Gharbi N. Antioxidant and anti-inflammatory effects of N-acetylcysteine against malathion-induced liver damages and immunotoxicity in rats. *Life Sci.* (2014) 107: 50-8.
- (61) Zhang Y, Geng J, Hong Y, Jiao L, Li S, Sun R, Xie Y, Yan C, Aa J and Wang G. Orally administered crocin protects against cerebral ischemia/reperfusion injury through the metabolic transformation of crocetin by gut microbiota. *Front. Pharmacol.* (2019) 10: 440-53.
- (62) Feidantsis K, Mellidis K, Galatou E, Sinakos Z and Lazou A. Treatment with crocin improves cardiac dysfunction by normalizing autophagy and inhibiting apoptosis in STZ-induced diabetic cardiomyopathy. *Nut. Met. Cardiovasc. Dis.* (2018) 28: 952-61.
- (63) Ribeiro S, Guilhermino L, Sousa J and Soares A. Novel bioassay based on acetylcholinesterase and lactate dehydrogenase activities to evaluate the toxicity of chemicals to soil isopods. *Ecotoxicol. Env. Safety* (1999) 44: 287-93.
- (64) Diamantino TC, Almeida E, Soares A M and Guilhermino L. Lactate dehydrogenase activity as an effect criterion in toxicity tests with *Daphnia magna* straus. *Chemosphere* (2001) 45: 553-60.
- (65) Hassoun EA and Stohs SJ. Cadmium-induced production of superoxide anion and nitric oxide, DNA single strand breaks and lactate dehydrogenase leakage in J774A. 1 cell cultures. *Toxicology* (1996) 112: 219-26.
- (66) Wu RS and Lam PK. Glucose-6-phosphate dehydrogenase and lactate dehydrogenase in the green-lipped mussel (*Perna viridis*): possible biomarkers for hypoxia in the marine environment. *Water Res.* (1997) 31: 2797-801.
- (67) Wilson Tang W, Francis GS, Morrow DA, Newby LK, Cannon CP, Jesse RL, Storrow AB, Christenson RH, Members C and Christenson RH. National academy of clinical biochemistry laboratory medicine practice guidelines: clinical utilization of cardiac biomarker testing in heart failure. *Circulation* (2007) 116: e99-e109.
- (68) Venkatesan R, Park YU, Ji E, Yeo EJ and Kim SY. Malathion increases apoptotic cell death by inducing lysosomal membrane permeabilization in N2a neuroblastoma cells: a model for neurodegeneration in Alzheimer's disease. *Cell. Death Discov.* (2017) 3: 1-10.
- (69) Vig S, Talwar P, Kaur K, Srivastava R, Srivastava AK and Datta M. Transcriptome profiling identifies p53 as a key player during calreticulin deficiency: Implications in lipid accumulation. *Cell Cycle* (2015) 14: 2274-84.
- (70) Agnetti G, Halperin VL, Kirk JA, Chakir K, Guo Y, Lund L, Nicolini F, Gherli T, Guarnieri C and Calderera CM. Desmin modifications associate with amyloid-like oligomers deposition in heart failure. *Cardiovasc. Res.* (2014) 102: 24-34.
- (71) Paulin D and Li Z. Desmin: a major intermediate filament protein essential for the structural integrity and function of muscle. *Exp. Cell Res.* (2004) 301: 1-7.
- (72) Ruepp SU, Tonge RP, Shaw J, Wallis N and Pognan F. Genomics and proteomics analysis of acetaminophen toxicity in mouse liver. *Toxicol. Sci.* (2002) 65: 135-50.
- (73) Nath BS. Changes in carbohydrate metabolism in hemolymph and fat body of the silkworm, *Bombyx mori* L., exposed to organophosphorus insecticides. *Pest. Bioch. Physiol.* (2000) 68: 127-37.
- (74) Bose C, Agarwal S, Chatterjee R and Srivastava V. Carboline antifilarials: effects on carbohydrate metabolising enzymes in *Litomosoides carinii* female. *Ind. Exp. Biol.* (1994) 32: 431-3.
- (75) Martinho-Dias D, Leite-Moreira A and Castro-

- Chaves P. Calreticulin in the heart: from embryological development to cardiac pathology. *Curr. Mol. Med.* (2016) 16: 12-22.
- (76) Jiang X, Ni Y, Zhang S, Zhang Y and Shen X. Identification of differentially expressed proteins in liver in response to subacute ruminal acidosis (SARA) induced by high-concentrate diet. *Asian-Aust. J. Animal Sci.* (2014) 27: 1181-8.
- (77) Michalak M, Corbett EF, Mesacli N, Nakamura K and Opas M. Calreticulin: one protein, one gene, many functions. *Biochem. J.* (1999) 344: 281-92.
- (78) Houle F, Poirier A, Dumaresq J and Huot J. DAP kinase mediates the phosphorylation of tropomyosin-1 downstream of the ERK pathway, which regulates the formation of stress fibers in response to oxidative stress. *J. Cell Sci.* (2007) 120: 3666-77.
- (79) Goldfarb L G, Park KY, Cervenáková L, Gorokhova S, Lee HS, Vasconcelos O, Nagle JW, Semino-Mora C, Sivakumar K and Dalakas MC. Missense mutations in desmin associated with familial cardiac and skeletal myopathy. *Nat. Gene.* (1998) 19: 402-3.

This article is available online at <http://www.ijpr.ir>
