

Original Article

## Effects of Lichenic Extracts (*Hypogymnia physodes*, *Ramalina polymorpha* and *Usnea florida*) on Human Blood Cells: Cytogenetic and Biochemical Study

Hasan Türkez<sup>a</sup>, Elanur Aydın<sup>a\*</sup> and Ali Aslan<sup>b</sup>

<sup>a</sup>Department of Molecular Biology and Genetics, Faculty of Science, Erzurum Technical University, Erzurum, Turkey. <sup>b</sup>Atatürk University, Kazım Karabekir Education Faculty, Department of Biology, 25240, Erzurum, Turkey.

---

### Abstract

Several lichen species have been used for medicinal purposes throughout the ages, and they were reported to be effective in the treatment of different disorders including tuberculosis, hemorrhoids, ulcer, dysentery and cancer. It is revealed that they may be easily accessible sources of natural drugs that could be used as a possible food supplement or in pharmaceutical industry after their safety evaluations. However, so far, the nature and/or biological roles of plenty of lichenes have not been elucidated exactly. The aim of this study was to investigate the genetic and oxidative effects of water extracts of three different lichen species; *Hypogymnia physodes*, *Ramalina polymorpha* and *Usnea florida* in cultured human blood cells (n = 5) for the first time. All lichen species were collected from the Erzurum and Artvin provinces (in Turkey) during August 2010. The lichen extracts were added into culture tubes at various concentrations (0 to 2000 mg/L). Chromosome aberrations (CA) and micronucleus (MN) tests were used for genotoxic influences estimation. In addition, biochemical parameters (total antioxidant capacity (TAC) and total oxidant stress (TOS)) were examined to determine oxidative effects. In our in-vitro test systems, it was observed that all tested lichen extracts had no mutagenic effects on human lymphocytes. Furthermore, these extracts exhibited antioxidant properties due to the type of lichen species added to the cultures. In conclusion, these lichens can be a new resource of therapeutics as recognized in this study with their non-mutagenic and antioxidant features.

**Keywords:** Chromosomal aberrations; *Hypogymnia physodes*; Micronucleus assay; *Ramalina polymorpha*; Total antioxidant capacity; Total oxidant status; *Usnea florida*.

---

### Introduction

Lichens are composite organisms consisting of a symbiotic association of a fungus (the mycobiont) with a photosynthetic partner (the photobiont), usually either a green algae or

cyanobacterium (1). Lichens were effective in the treatment of diseases such as hemorrhoids, bronchitis, dysentery, and tuberculosis (2). Again, lichen species have been used as stomachic, antidiabetic, and hemostatic drug (3). Some recent studies have also revealed that secondary metabolites from lichens induce apoptosis in colon (4, 5) and prostate (6) cancers. Lichens have long been investigated

---

\* Corresponding author:

E-mail: elanuraydinn@gmail.com

for biological activities; mainly antimicrobial but also antitumor, antiviral, allergenic, plant growth inhibitory, antiherbivore, and enzyme inhibitory (7), more recently, antioxidant and anti-inflammatory activities (8-10).

Antioxidants could inhibit or delay the oxidation process by blocking the initiation or propagation of oxidizing chain reactions. A variety of synthetic antioxidants like butylated hydroxyanisole, butylated hydroxytoluene, and tert-butylhydroquinone are commonly used within the food industry although restrictions on the use of synthetic antioxidants are being imposed because of their toxicity (11-13). Much attention has recently been focused on the development of safe and effective antioxidants (14), because toxic free radicals play a role in the etiology of many disorders including neurodegenerative and cardiovascular diseases (15, 16), diabetes (18) and certain cancers (19). Therefore, the development and utilization of more effective and less harmful antioxidants of natural origins are reported to be very desirable (13).

Lichen species are very common in Turkey. It is pointed that they may be easily accessible sources of natural drugs that could be used as a possible food supplement or in pharmaceutical industry (20). In this investigation, it was aimed to describe the cytogenetic and oxidative effects of three lichen species; *Hypogymnia physodes*, *Ramalina polymorpha* and *Usnea florida* on cultured human blood cells for utilization as a possible food supplement or within the pharmaceutical industry. And, safety screening of these lichen extracts could also serve to explore new natural therapeutics. With this aim, not only CA frequencies but also MN formations have been established as related to the dose of aqueous extracts of lichenes on human lymphocytes. In addition, important oxidative parameters, TAC and TOS were used to monitor their antioxidant or pro-oxidant activities *in-vitro*.

## Experimental

### *Plant materials*

Lichen species, *Hypogymnia physodes* (L.) Nyl., *Ramalina polymorpha* (Lilj.) Ach. and *Usnea florida* (L.) Wigg. ex Web. em Clerc.

were collected from the Erzurum and Artvin provinces during August 2010. After drying at the room temperature a stereo microscope (since it produces a three-dimensional visualization of the sample being examined) and the usual spot tests were used in the identification of the samples with the reference books (20-23). The specimens are stored in the herbarium of Kazım Karabekir, Faculty of Education, Atatürk University, Erzurum.

### *Extraction*


For water extraction out of lichenes, 20 g sample was mixed with 400 mL distilled and boiling water using magnetic stirrer for 15 min. Then the extracts were filtered over Whatmann No. 1 paper.

### *Experimental design*

Whole heparinized human blood from five healthy non-smoking female donors between the ages 27 and 28 with no history of exposure to any genotoxic agent was used in our experiments. Questionnaires were obtained for each blood donor to evaluate exposure history, and in addition, informed consent forms were signed by each donor. In all the volunteers involved in this study, hematological and biochemical parameters were analyzed and no pathology was detected. A various concentrations (0, 1, 5, 10, 25, 50, 100, 200, 250, 500, 1000 and 2000 mg/L) of aqueous extracts of lichenes were tested in blood cultures. CA and MN rates were assessed in peripheral lymphocytes. Experiments conformed to the guidelines of the World Medical Assembly (Declaration of Helsinki) (24). The cultures without extracts were studied as control group. Mitomycin C ( $10^{-7}$ M) was used as the positive control in CA and MN assays. Likewise, ascorbic acid (10  $\mu$ M) and hydrogen peroxide (25  $\mu$ M) were also used as the positive controls in TAC and TOS analysis, respectively.

### *CA assay*

Human peripheral blood lymphocyte cultures were set up according to a slight modification of the protocol described by Evans and O'Riordan (25). A 0.5 mL aliquot of heparinized blood was cultured in 6 mL of culture medium (Chromosome



**Figure 1.** A sample binucleated cells (A) and micronucleus (arrow) formation (B) in lymphocyte cells.

Medium B; Biochrom®, Berlin) with 5 mg/mL of phytohemagglutinin (Biochrom®). The cultures were incubated in complete darkness for 72 h at 37°C. Two hours prior to harvesting, 0.1 mL of colchicine (0.2 mg/mL, Sigma®) was added to the culture flask. Hypotonic treatment and fixation were performed. To prepare slides, 3–5 drops of the fixed cell suspension were dropped on a clean slide and air-dried. The slides were stained in 3% Giemsa solution in phosphate buffer (pH 6.8) for 15 min. For each treatment, 30 well-spread metaphases were analyzed to detect the presence of chromosomal aberrations. Criteria to classify the different types of aberrations (chromatid or chromosome gap and chromatid or chromosome break) were in accordance with the recommendation of EHC (Environmental Health Criteria) 46 for environmental monitoring of human populations (26).

#### *MN assay*

The micronucleus test was performed by adding cytochalasin B (Sigma®; final concentration of 6 µg/mL) after 44 h of culture. At the end of the 72-h incubation period, the lymphocytes were fixed with ice-cold methanol: acetic acid (1:1). The fixed cells were put directly on slides using a cytospin and stained with Giemsa. All slides were coded before scoring. The criteria for scoring micronuclei were as

described by Fenech (27). A sample binucleated lymphocyte cells and micronucleus formation was shown in Figure 1A and B, respectively. At least 1000 binucleated lymphocytes were examined per concentration for the presence of one, two, or more micronuclei.

#### *TAC and TOS analysis*

The automated total antioxidant capacity and total oxidant status assays were carried out in plasma samples obtained from blood cultures for 2 h by commercially available kits (Rel Assay Diagnostics®, Turkey) (27).

#### *Statistical analysis*

Statistical analysis was performed using SPSS software (version 13.0, SPSS, Chicago, IL, USA). The Duncan's was used to determine whether any treatment significantly differed from controls or each other. Statistical decisions were made with a significance level of 0.05.

## **Results and Discussion**

Tables 1, 2, and 3 show the genetic and biochemical data obtained with various concentrations of *Hypogymnia physodes*, *Ramalina polymorpha* and *Usnea florida*, on cultured blood cells, respectively. All the lichen extracts at tested concentrations did not induce significant ( $p < 0,05$ ) number of CAs

**Table 1.** The genetic and biochemical effects of aqueous extracts of *Hypogymnia physodes in-vitro*.

Treatments	TAC (mmol Trolox Equiv./L)	TOS ( $\mu\text{mol H}_2\text{O}_2$ Equiv./L)	MN/1000 cell	CA/cell
Control <sup>-</sup>	6.24 $\pm$ 0.62 <sup>b</sup>	11.58 $\pm$ 2.64 <sup>a</sup>	3.18 $\pm$ 0.72 <sup>a</sup>	0.20 $\pm$ 0.02 <sup>a</sup>
Control <sup>+</sup>	13.71 $\pm$ 0.94 <sup>d</sup>	39.25 $\pm$ 4.63 <sup>c</sup>	8.26 $\pm$ 1.18 <sup>b</sup>	0.72 $\pm$ 0.09 <sup>b</sup>
1 mg/L	6.25 $\pm$ 0.43 <sup>b</sup>	11.27 $\pm$ 2.76 <sup>a</sup>	3.35 $\pm$ 0.48 <sup>a</sup>	0.20 $\pm$ 0.03 <sup>a</sup>
5 mg/L	6.27 $\pm$ 0.42 <sup>b</sup>	11.02 $\pm$ 2.54 <sup>a</sup>	3.40 $\pm$ 0.55 <sup>a</sup>	0.21 $\pm$ 0.02 <sup>a</sup>
10 mg/L	6.25 $\pm$ 0.39 <sup>b</sup>	11.01 $\pm$ 3.01 <sup>a</sup>	3.37 $\pm$ 0.62 <sup>a</sup>	0.23 $\pm$ 0.02 <sup>a</sup>
25 mg/L	6.26 $\pm$ 0.48 <sup>b</sup>	11.60 $\pm$ 2.74 <sup>a</sup>	3.39 $\pm$ 0.64 <sup>a</sup>	0.21 $\pm$ 0.03 <sup>a</sup>
50 mg/L	7.22 $\pm$ 0.41 <sup>c</sup>	11.63 $\pm$ 3.02 <sup>a</sup>	3.36 $\pm$ 0.59 <sup>a</sup>	0.22 $\pm$ 0.01 <sup>a</sup>
100 mg/L	6.27 $\pm$ 0.38 <sup>b</sup>	11.62 $\pm$ 2.94 <sup>a</sup>	3.34 $\pm$ 0.63 <sup>a</sup>	0.18 $\pm$ 0.03 <sup>a</sup>
200 mg/L	6.26 $\pm$ 0.45 <sup>b</sup>	11.64 $\pm$ 2.85 <sup>a</sup>	3.37 $\pm$ 0.61 <sup>a</sup>	0.21 $\pm$ 0.02 <sup>a</sup>
250 mg/L	6.28 $\pm$ 0.44 <sup>b</sup>	12.03 $\pm$ 3.10 <sup>b</sup>	3.32 $\pm$ 0.58 <sup>a</sup>	0.18 $\pm$ 0.03 <sup>a</sup>
500 mg/L	6.27 $\pm$ 0.47 <sup>b</sup>	12.27 $\pm$ 3.04 <sup>b</sup>	3.36 $\pm$ 0.51 <sup>a</sup>	0.22 $\pm$ 0.04 <sup>a</sup>
1000 mg/L	5.59 $\pm$ 0.43 <sup>a</sup>	13.35 $\pm$ 3.12 <sup>b</sup>	-	-
2000 mg/L	5.12 $\pm$ 0.46 <sup>a</sup>	14.02 $\pm$ 3.40 <sup>b</sup>	-	-

Values are presented as mean $\pm$ S.D.; n = 5, means shown by the same letter are not significantly different from each other at a level of 5%, using Duncan's test. For example, the mean shown by the letter b is different from a, c and d means, but not different from b mean. In other words, the means having the different letters are different from each other.

and MNs. However, the Mitomycin C applied culture (positive control) showed about three fold increases of both parameters as compared to control group.

In the present study, it was established that the extracts of *H. physodes*, *R. polymorpha* and *U. florida* lichen species were non-genotoxic. The results obtained through the present study did not indicate any significant increases in the ratios of the CAs and MNs in lymphocytes exposed to lichen extracts as compared to control values. In fact, CA test is regarded as a very important and useful indicator of exposure to biological and chemical agents (29). MN assay provides a measure of both chromosome breakage and chromosome loss or non-disjunction in clastogenic and aneugenic events, respectively (30). And damaged DNA can lead to aneuploidy and/or chromosomal instability, which is believed to be major contributor to tumor progression (31). Our findings are in accordance with the previous reports. Koparal *et al.* (32) investigated cytotoxic and genotoxic activities of the lichen *Ramalina farinacea* and the lichen *Cladonia foliacea* and suggested that usnic acid (a main component of lichens) was non-genotoxic

shown by the absence of MN induction in human lymphocytes. Again, Zeytinoglu *et al.* (33) investigated the genotoxic/antigenotoxic activities of the extract from lichen *Cetraria aculeata* in TA98 and TA100 strains of Salmonella typhimurium in the presence or absence of metabolic activity and in human lymphocytes. They have reported that the lichen extract was not mutagenic in all systems. Again, the genotoxic effects of the water extracts of *Pseudevernia furfuracea*, *Dermotocarpon intestiniforme*, *Ramalina capitata*, *Parmelia pulla* and *Rhizoplaca melanophthalma* lichens were ascertained by sister-chromatid exchange (SCE) and MN tests in human whole blood cultures. According to results of this study, it was established that these lichen extracts had also no genotoxic effect (34). The separated components of Chinese lichen extract such as AMH-C, AMH-D and AMH-E were determined as non genotoxic in Ames test (35). Turkez *et al.* (10) studied the effects of methanol, acetone, n-hexane and ether extracts obtained from the lichen, *Pseudovernia furfuracea*, on genotoxicity in cultured human blood cells by SCE and MN tests. The researchers observed that *P.*

**Table 2.** The genetic and biochemical effects of aqueous extracts of *Ramalina polymorpha in-vitro*.

Treatments	TAC (mmol Trolox Equiv./L)	TOS ( $\mu\text{mol H}_2\text{O}_2$ Equiv./L)	MN/1000 cell	CA/cell
Control <sup>-</sup>	6.24 $\pm$ 0.62 <sup>b</sup>	11.58 $\pm$ 2.64 <sup>a</sup>	3.18 $\pm$ 0.72 <sup>a</sup>	0.20 $\pm$ 0.02 <sup>a</sup>
Control <sup>+</sup>	13.71 $\pm$ 0.94 <sup>d</sup>	39.25 $\pm$ 4.63 <sup>c</sup>	8.26 $\pm$ 1.18 <sup>b</sup>	0.72 $\pm$ 0.09 <sup>b</sup>
1 mg/L	6.22 $\pm$ 0.56 <sup>b</sup>	11.54 $\pm$ 2.83 <sup>a</sup>	3.22 $\pm$ 0.45 <sup>a</sup>	0.21 $\pm$ 0.02 <sup>a</sup>
5 mg/L	6.29 $\pm$ 0.61 <sup>b</sup>	11.63 $\pm$ 3.05 <sup>a</sup>	3.25 $\pm$ 0.49 <sup>a</sup>	0.24 $\pm$ 0.03 <sup>a</sup>
10 mg/L	6.31 $\pm$ 0.49 <sup>b</sup>	11.49 $\pm$ 2.75 <sup>a</sup>	3.27 $\pm$ 0.43 <sup>a</sup>	0.22 $\pm$ 0.02 <sup>a</sup>
25 mg/L	6.69 $\pm$ 0.54 <sup>b</sup>	11.13 $\pm$ 3.18 <sup>a</sup>	3.35 $\pm$ 0.41 <sup>a</sup>	0.20 $\pm$ 0.02 <sup>a</sup>
50 mg/L	9.29 $\pm$ 0.79 <sup>c</sup>	11.56 $\pm$ 2.86 <sup>a</sup>	3.29 $\pm$ 0.48 <sup>a</sup>	0.21 $\pm$ 0.03 <sup>a</sup>
100 mg/L	6.58 $\pm$ 0.62 <sup>b</sup>	11.49 $\pm$ 2.74 <sup>a</sup>	3.32 $\pm$ 0.46 <sup>a</sup>	0.23 $\pm$ 0.02 <sup>a</sup>
200 mg/L	6.63 $\pm$ 0.67 <sup>b</sup>	11.47 $\pm$ 3.07 <sup>a</sup>	3.37 $\pm$ 0.51 <sup>a</sup>	0.21 $\pm$ 0.02 <sup>a</sup>
250 mg/L	6.71 $\pm$ 0.48 <sup>b</sup>	11.72 $\pm$ 2.93 <sup>a</sup>	3.29 $\pm$ 0.58 <sup>a</sup>	0.20 $\pm$ 0.03 <sup>a</sup>
500 mg/L	6.30 $\pm$ 0.51 <sup>b</sup>	11.74 $\pm$ 3.16 <sup>a</sup>	3.36 $\pm$ 0.54 <sup>a</sup>	0.22 $\pm$ 0.02 <sup>a</sup>
1000 mg/L	6.09 $\pm$ 0.45 <sup>b</sup>	11.81 $\pm$ 2.88 <sup>a</sup>	3.24 $\pm$ 0.61 <sup>a</sup>	0.21 $\pm$ 0.02 <sup>a</sup>
2000 mg/L	5.71 $\pm$ 0.47 <sup>a</sup>	11.86 $\pm$ 3.03 <sup>a</sup>	-	-

Values are presented as mean $\pm$ S.D.; n=5, means shown by the same letter are not significantly different from each other at a level of 5%, using Duncan's test. For example, the mean shown by the letter b is different from a, c and d means, but not different from b mean. In other words, the means having the different letters are different from each other.

*furfuracea* extracts exhibited non-mutagenic properties in both test systems.

Different concentrations of *H. physodes* and *R. polymorpha* (50 mg/L) and *U. florida* (100 mg/L) caused significant increases of TAC level when compared to control<sup>-</sup> value. In contrast, *H. physodes* (at concentrations of 1000 and 2000 mg/L), *R. polymorpha* (at concentration of 2000 mg/L) and *U. florida* (at concentrations of 1000 and 2000 mg/L) caused significant decreases of TAC level. As shown from the results presented in Tables 1, 2 and 3, the TOS levels increased at higher concentrations of *H. physodes* (250, 500, 1000 and 2000 mg/L) and *U. florida* (2000 mg/L). However, *R. Polymorpha* did not cause any significant increases of TOS levels. Besides, the cultures found to be sterile at concentrations of 1000, 2000 and 1000 mg/L for *H. Physodes*, *R. polymorpha* and *U. florida*, respectively.

The results of the present study reveal that treatment with aqueous extracts of *H. physodes*, *R. polymorpha* and *U. florida* lichen species provide antioxidant effects at different degree. Similarly to our findings, various reports, especially published in last ten years, indicated the antioxidant properties of several

lichen species. According to the previous studies, the extracts of *Cladonia clathrata* (8), *Pseudovernia furfuracea* (10), *Xanthoparmelia spp.* (36), *Lethariella sernanderi*, *L. cashmeriana*, and *L. sinensis* (37), *Lobaria pulmonaria* (38), *Usnea ghattensis* (39), *Usnea longissima* (40), Graphidaceae (41), *Lethariella canariensis* (42), *Cetraria islandica* (19), *Parmelia caperata* and *P. soredians* (43), *Dermatocarpon miniatum* (44), *Parmotrema stuppeum* (45) were found to have antioxidant properties.

The results of the present study also indicated that the aqueous extracts of *H. physodes* and *U. florida* lichen species caused sterility of human blood cultures for 72 h at concentration of 2000 mg/L. Similarly, *R. polymorpha* caused sterility at concentration of 1000 mg/L. On the other hand, *H. physodes* (at concentrations of 250, 500, 1000 and 2000 mg/L) and *U. florida* (at concentration of 2000 mg/L) caused significant increases of TOS levels at increasing doses. However, *R. Polymorpha* did not cause significant increases of TOS levels. Hence, the cytotoxic effects of this lichen species could be, at least in part, attributed to oxidative stress-induced by

**Table 3.** The genetic and biochemical effects of aqueous extracts of *Usnea florida in-vitro*.

Treatments	TAC (mmol Trolox Equiv./L)	TOS ( $\mu\text{mol H}_2\text{O}_2$ Equiv./L)	MN/1000 cell	CA/cell
Control <sup>-</sup>	6.24 $\pm$ 0.62 <sup>b</sup>	11.58 $\pm$ 2.64 <sup>a</sup>	3.18 $\pm$ 0.72 <sup>a</sup>	0.20 $\pm$ 0.02 <sup>a</sup>
Control <sup>+</sup>	13.71 $\pm$ 0.94 <sup>d</sup>	39.25 $\pm$ 4.63 <sup>c</sup>	8.26 $\pm$ 1.18 <sup>b</sup>	0.72 $\pm$ 0.09 <sup>b</sup>
1 mg/L	6.27 $\pm$ 0.53 <sup>b</sup>	11.97 $\pm$ 2.14 <sup>a</sup>	3.43 $\pm$ 0.63 <sup>a</sup>	0.22 $\pm$ 0.03 <sup>a</sup>
5 mg/L	6.30 $\pm$ 0.62 <sup>b</sup>	11.73 $\pm$ 3.13 <sup>a</sup>	3.27 $\pm$ 0.34 <sup>a</sup>	0.20 $\pm$ 0.04 <sup>a</sup>
10 mg/L	6.25 $\pm$ 0.57 <sup>b</sup>	11.58 $\pm$ 2.42 <sup>a</sup>	3.18 $\pm$ 0.22 <sup>a</sup>	0.20 $\pm$ 0.02 <sup>a</sup>
25 mg/L	6.38 $\pm$ 0.61 <sup>b</sup>	11.33 $\pm$ 3.07 <sup>a</sup>	2.87 $\pm$ 0.39 <sup>a</sup>	0.20 $\pm$ 0.02 <sup>a</sup>
50 mg/L	6.69 $\pm$ 0.67 <sup>b</sup>	11.56 $\pm$ 2.88 <sup>a</sup>	3.32 $\pm$ 0.55 <sup>a</sup>	0.18 $\pm$ 0.03 <sup>a</sup>
100 mg/L	7.33 $\pm$ 0.49 <sup>c</sup>	11.14 $\pm$ 2.43 <sup>a</sup>	3.23 $\pm$ 0.35 <sup>a</sup>	0.15 $\pm$ 0.03 <sup>a</sup>
200 mg/L	6.51 $\pm$ 0.55 <sup>b</sup>	11.63 $\pm$ 3.17 <sup>a</sup>	3.25 $\pm$ 0.38 <sup>a</sup>	0.22 $\pm$ 0.03 <sup>a</sup>
250 mg/L	6.35 $\pm$ 0.57 <sup>b</sup>	11.72 $\pm$ 2.43 <sup>a</sup>	3.39 $\pm$ 0.57 <sup>a</sup>	0.24 $\pm$ 0.02 <sup>a</sup>
500 mg/L	6.18 $\pm$ 0.59 <sup>b</sup>	11.82 $\pm$ 2.77 <sup>a</sup>	3.41 $\pm$ 0.69 <sup>a</sup>	0.24 $\pm$ 0.02 <sup>a</sup>
1000 mg/L	5.43 $\pm$ 0.43 <sup>a</sup>	11.88 $\pm$ 3.14 <sup>a</sup>	-	-
2000 mg/L	5.22 $\pm$ 0.51 <sup>a</sup>	12.76 $\pm$ 3.27 <sup>b</sup>	-	-

Values are presented as mean $\pm$ S.D.; n=5, means shown by the same letter are not significantly different from each other at a level of 5%, using Duncan's test. For example, the mean shown by the letter b is different from a, c and d means, but not different from b mean. In other words, the means having the different letters are different from each other.

high lichenic contents. The multidisciplinary toxicity studies can be of help here as well. But cytotoxic actions of these lichens at higher concentrations were also could be due to initiation of cell death, inhibition of functional enzymes, antimitochondrial actions, binding to plasma membrane and changes in osmotic potential. Therefore, further investigations are necessary to find out the definite mechanisms in lichen toxicity.

In conclusion, our results clearly indicated that the water extracts of *H. physodes*, *R. polymorpha* and *U. florida* lichen species collected from Erzurum and Artvin provinces had no mutagenic effects on human lymphocytes. Furthermore, these extracts exhibited antioxidant properties due to the applied dose and the type of lichen species added to the cultures. On the other hand, the extracts caused sterility of cultures due to oxidative stress at higher concentrations above than 1000 mg/L. Generally speaking, our results have indicated that several North East Anatolian lichens have the potential of being utilized as novel bioresources for naturally occurring antioxidant therapies.

#### Acknowledgment

The authors are grateful to all volunteers of the blood samples.

#### References

- (1) Nash I and Thomas H. *Lichen Biology*. Cambridge University Press, UK (1996) 304.
- (2) Dulger B, Gucin F and Aslan A. Antimicrobial effect of lichen *Cetraria islandica* (L) Ach. *Turk. J. Biol.* (1998) 22: 111-18.
- (3) Baytop T. *Therapy with Medicinal Plants in Turkey (Past and Present)*. Istanbul University Press, Istanbul (1999) 1-233.
- (4) Bezivin C, Tomasi M, Lohezic-Le Devehat F and Boustie J. Cytotoxic activity of some lichen extracts on murine and human cancer cell lines. *Phytomed.* (2003) 10: 499-503.
- (5) Ren MR, Hur JS, Kim JY, Park KW, Park SC and Seong CN. Anti-proliferative effects of *Lethariella zahlbruckneri* extracts in human HT-29 human colon cancer cells. *Food Chem. Toxicol.* (2009) 47: 2157-62.
- (6) Russo A, Piovano M, Lombardo L, Vanella L, Cardile V and Garbarino J. Pannarin inhibits cell growth and induces cell death in human prostate carcinoma DU-145 cells. *Anticancer Drugs* (2006) 17: 1163-69.
- (7) Huneck S. The significance of lichens and their metabolites. *Naturwissenschaften* (1999) 86: 559-70.
- (8) Silva JA, Bomfim RR, Estevam Cdos S, Antonioli AR, Araújo AA and Thomazzi SM. Pharmacological properties of lichen *Cladonia clathrata*. *Pharm. Biol.* (2010) 48: 745-52.
- (9) Javadzadeh A, Vatanpour H, Delavarian Z, Momajed A, Esmaily H, Vatanpour M and Shirazian SH. Efficacy of Clobetasol, Ketoconazole and Amitriptyline Mouthwash on Oral Lichen Planus. *Iranian J. Pharm. Res.* (2008) 7: 171-178.
- (10) Türkez H, Geyikoglu F, Aslan A, Karagöz Y, Türkez

- O and Anar M. Antimutagenic effects of lichen *Pseudovernia furfuracea* (L.) Zoph. extracts against the mutagenicity of aflatoxin B1 *in-vitro*. *Toxicol. Ind. Health*. (2010) 26: 625-31.
- (11) Grice HC. Safety evaluation of butylated hydroxytoluene (BHT) in the liver, lung and gastrointestinal tract. *Food Chem. Toxicol.* (1986) 24: 1127-30.
  - (12) Wichl HP. Enhanced tumor development by butylated hydroxyanisole (BHA) from the prospective of effect on forestomach and oesophageal squamous epithelium. *Food Chem. Toxicol.* (1988) 26: 717-23.
  - (13) Luo H, Yoshikazu Y, Liu Y, Jung JS, Kahng HY, Koh YJ and Hur JS. The *in-vitro* antioxidant properties of Chinese highland lichens. *J. Microbiol. Biotechnol.* (2010) 20: 1524-28.
  - (14) Choi JS, Chung HY, Jung HA, Park HJ and Yokozawa T. Comparative evaluation of antioxidant potential of alaternin (2-hydroxyemodin) and emodin. *J. Agric. Food Chem.* (2000) 48: 6347-51.
  - (15) Dumont M and Beal MF. Neuroprotective strategies involving ROS in Alzheimer disease. *Free Radic. Biol. Med.* (2010) 51: 1014-26.
  - (16) Lorgis L, Zeller M, Dentan G, Sicard P, Richard C, Buffet P, L'Huillier I, Beer JC, Cottin Y, Rochette L and Vergely C. The free oxygen radicals test (FORT) to assess circulating oxidative stress in patients with acute myocardial infarction. *Atherosclerosis* (2010) 213: 616-21.
  - (17) Giacco F and Brownlee M. Oxidative stress and diabetic complications. *Circ. Res.* (2010) 107: 1058-70.
  - (18) Ziech D, Franco R, Georgakilas AG, Georgakila S, Malamou-Mitsi V, Schoneveld O, Pappa A and Panayiotidis MI. The role of reactive oxygen species and oxidative stress in environmental carcinogenesis and biomarker development. *Chem. Biol. Interact.* (2010) 188: 334-39.
  - (19) Gülçin İ, Oktay M, Küfrevioğlu Öİ and Aslan A. Determination of antioxidant activity of lichen *Cetraria islandica* (L.) Ach. *J. Ethnopharmacol.* (2002) 79: 325-29.
  - (20) Aslan A and Yazici K. Contribution to the lichen flora of Giresun province of Turkey. *Acta. Bot. Hung.* (2006) 48: 231-45.
  - (21) Poelt J. *Bestimmungsschlüssel Europäischer Flechten*. Cramer, Lehre (1974) 757.
  - (22) Purvis OW, Coppins BJ, Hawksworth DL, James PW and Moore DM. *The Lichen Flora of Great Britain and Ireland*. Natural History Museum, London (1992) 710. Wirth V. *Die Flechten Baden Württenbergs*. Teil 1-2 Ulmer, Stuttgart (1995) 1006.
  - (23) WMA Medical Ethics Committee. Updating the WMA Declaration of Helsinki. *Wld. Med. J.* (1999) 45: 11-13.
  - (24) Evans HJ and O'Riordan ML. Human peripheral blood lymphocytes for the analysis of chromosome aberrations in mutagen tests. *Mutat. Res.* (1975) 31: 135-48.
  - (25) International Program on Chemical Safety (IPCS). *Environmental Health Criteria 46. Guidelines for the Study of Genetic Effects in Human Populations*. WHO, Geneva (1985) 45-54.
  - (26) Fenech M. The cytokinesis blocks micronucleus technique. A detailed description on the method and its application to genotoxicity studies in human population. *Mutat. Res.* (1993) 285: 35-44.
  - (27) Erel O. A novel automated direct measurement method for total antioxidant capacity using a new generation, more stable ABTS radical cation. *Clin. Biochem.* (2004) 37: 277-85.
  - (28) Padovani L, Tronati L, Mauro F, Testa A, Appolloni M, Azzidei P, Caporossi D, Tedeschi B and Vernole P. Cytogenetic effects in lymphocytes from children exposed to radiation fall-out after the chernobyl accident. *Mutat. Res.* (1997) 395: 249-54.
  - (29) Karaman A, Kadı M and Kara F. Sister chromatid exchange and micronucleus studies in patients with Behçet's disease. *J. Cutan. Pathol.* (2009) 36: 831-37.
  - (30) Erol A. Systemic DNA damage response and metabolic syndrome as a premalignant state. *Curr. Mol. Med.* (2010) 10: 321-34.
  - (31) Koparal AT, Tüylü BA and Türk H. *In-vitro* cytotoxic activities of (+)-usnic acid and (-)-usnic acid on V79, A549, and human lymphocyte cells and their non-genotoxicity on human lymphocytes. *Nat. Prod. Res.* (2006) 20: 1300-07.
  - (32) Zeytinoglu H, Incesu Z, Tuylu BA, Turk AO and Barutca B. Determination of genotoxic, antigenotoxic and cytotoxic potential of the extract from lichen *Cetraria aculeata* (Schreb.) Fr. *In-vitro. Phytother. Res.* (2008) 22: 118-23.
  - (33) Geyikoglu F, Turkez H and Aslan A. The protective roles of some lichen species on colloidal bismuth subcitrate genotoxicity. *Toxicol. Ind. Health* (2007) 23: 487-92.
  - (34) He X, Hu Y, Winter J and Young GP. Antimutagenic lichen extract has double-edged effect on azoxymethane-induced colorectal oncogenesis in C57BL/6J mice. *Toxicol. Mech. Meth.* (2010) 20: 31-35.
  - (35) de Paz GA, Raggio J, Gómez-Serranillos MP, Palomino OM, González-Burgos E, Carretero ME and Crespo A. HPLC isolation of antioxidant constituents from *Xanthoparmelia* spp. *J. Pharmaceut. Med.* (2010) 53: 165-71.
  - (36) Kinoshita K, Togawa T, Hiraishi A, Nakajima Y, Koyama K, Narui T, Wang LS and Takahashi K. Antioxidant activity of red pigments from the lichens *Lethariella sernanderi*, *L. cashmeriana*, and *L. sinensis*. *J. Nat. Med.* (2010) 64: 85-88.
  - (37) Karakus B, Odabasoglu F, Cakir A, Halici Z, Bayir Y, Halici M, Aslan A and Suleyman H. The effects of methanol extract of *Lobaria pulmonaria*, a lichen species, on indometacin-induced gastric mucosal damage, oxidative stress and neutrophil infiltration. *Phytother. Res.* (2009) 23: 635-39.
  - (38) Verma N, Behera BC and Makhija U. Antioxidant and hepatoprotective activity of a lichen *Usnea ghattensis*

- in-vitro. Appl. Biochem. Biotech.* (2008) 151: 167-81.
- (39) Halici M, Odabasoglu F, Suleyman H, Cakir A, Aslan A and Bayir Y. Effects of water extract of *Usnea longissima* on antioxidant enzyme activity and mucosal damage caused by indomethacin in rats. *Phytomed.* (2005) 12: 656-62.
- (40) Behera BC, Adawadkar B and Makhija U. Inhibitory activity of xanthine oxidase and superoxide-scavenging activity in some taxa of the lichen family Graphidaceae. *Phytomed.* (2003) 10: 536-43.
- (41) Toledo Marante FJ, García Castellano A, Estévez Rosas F, Quintana Aguiar J and Bermejo Barrera J. Identification and quantitation of allelochemicals from the lichen *Lethariella canariensis*: phytotoxicity and antioxidative activity. *J. Chem. Ecol.* (2003) 29: 2049-71.
- (42) Caviglia AM, Nicora P, Giordani P, Brunialti G and Modenesi P. Oxidative stress and usnic acid content in *Parmelia caperata* and *Parmelia soledians* (Lichenes). *Farmaco.* (2001) 56: 379-82.
- (43) Jin J, Bian X, Ge P, Jing H, Ding L and Ding D. The effects of the polysaccharides from *Dermatocarpon miniatum* on oxygen radicals and lipid peroxidation. *Zhong Yao. Cai.* (2001) 24: 660-1.
- (44) Jayaprakasha GK and Rao LJ. Phenolic constituents from the lichen *Parmotrema stippeum* (Nyl.) Hale and their antioxidant activity. *Z. Naturforsch C.* (2000) 55: 1018-22.
- 
- This article is available online at <http://www.ijpr.ir>